



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 2, 2023 – 08:36 PM JST

PDB ID : 8IVU  
Title : Crystal Structure of Human NAMPT in complex with A4276  
Authors : Kang, B.G.; Cha, S.S.  
Deposited on : 2023-03-28  
Resolution : 2.09 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	491	 8% 90% 5% 5%
1	B	491	 8% 91% 5% 5%
1	C	491	 6% 90% 5% 5%
1	D	491	 13% 88% 7% 5%

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 15794 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nicotinamide phosphoribosyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	466	Total 3730	C 2399	N 617	O 707	S 7	0	0	0
1	B	466	Total 3730	C 2399	N 617	O 707	S 7	0	0	0
1	C	466	Total 3730	C 2399	N 617	O 707	S 7	0	0	0
1	D	466	Total 3730	C 2399	N 617	O 707	S 7	0	0	0

- Molecule 2 is N-[[4-(6-methyl-1,3-benzoxazol-2-yl)phenyl]methyl]pyridine-3-carboxamide (three-letter code: SJ6) (formula: C<sub>21</sub>H<sub>17</sub>N<sub>3</sub>O<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	Total 26	C 21	N 3	O 2	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	N	O	0	0
			26	21	3	2		
2	C	1	Total	C	N	O	0	0
			26	21	3	2		
2	D	1	Total	C	N	O	0	0
			26	21	3	2		

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	A	1	Total	O	P	0	0
			5	4	1		
3	A	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		
3	C	1	Total	O	P	0	0
			5	4	1		
3	C	1	Total	O	P	0	0
			5	4	1		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total	O	P	0	0
			5	4	1		
3	D	1	Total	O	P	0	0
			5	4	1		
3	D	1	Total	O	P	0	0
			5	4	1		
3	D	1	Total	O	P	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	206	Total	O	0	0
			206	206		
4	B	216	Total	O	0	0
			216	216		
4	C	180	Total	O	0	0
			180	180		
4	D	108	Total	O	0	0
			108	108		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Nicotinamide phosphoribosyltransferase




- Molecule 1: Nicotinamide phosphoribosyltransferase




- Molecule 1: Nicotinamide phosphoribosyltransferase



- Molecule 1: Nicotinamide phosphoribosyltransferase





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	97.56Å 116.69Å 195.41Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.71 – 2.09 29.71 – 2.09	Depositor EDS
% Data completeness (in resolution range)	100.0 (29.71-2.09) 100.0 (29.71-2.09)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.53 (at 2.10Å)	Xtrriage
Refinement program	PHENIX 1.11.1-2575-000	Depositor
R, $R_{free}$	0.204 , 0.231 0.204 , 0.231	Depositor DCC
$R_{free}$ test set	6612 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	40.4	Xtrriage
Anisotropy	0.255	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 38.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	15794	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.70% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SJ6, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.26	0/3818	0.45	0/5172
1	B	0.26	0/3818	0.45	0/5172
1	C	0.25	0/3818	0.45	0/5172
1	D	0.25	0/3818	0.44	0/5172
All	All	0.26	0/15272	0.45	0/20688

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3730	0	3707	13	0
1	B	3730	0	3707	10	0
1	C	3730	0	3707	12	0
1	D	3730	0	3707	18	0
2	A	26	0	0	0	0
2	B	26	0	0	0	0
2	C	26	0	0	0	0
2	D	26	0	0	0	0
3	A	15	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	15	0	0	0	0
3	C	15	0	0	0	0
3	D	15	0	0	0	0
4	A	206	0	0	1	0
4	B	216	0	0	1	0
4	C	180	0	0	2	0
4	D	108	0	0	1	0
All	All	15794	0	14828	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:179:THR:HG21	1:D:375:ILE:H	1.40	0.84
1:B:357:ASP:OD2	1:B:389:LYS:NZ	2.21	0.73
1:C:288:GLU:HG3	1:C:331:LYS:HE3	1.70	0.72
1:B:391:THR:OG1	1:B:393:ASP:OD1	2.08	0.71
1:A:234:LYS:HD3	1:D:234:LYS:HA	1.74	0.69
1:C:429:ARG:NH2	1:C:449:ASP:OD2	2.29	0.64
1:D:429:ARG:NH2	1:D:449:ASP:OD2	2.30	0.63
1:A:237:VAL:HG21	1:B:90:HIS:HA	1.81	0.63
1:D:41:GLU:HG3	1:D:128:GLY:HA3	1.81	0.63
1:C:357:ASP:OD2	1:C:389:LYS:NZ	2.33	0.60
1:D:322:LEU:HD13	1:D:371:LYS:HD2	1.82	0.60
1:D:291:TRP:HE3	1:D:299:ILE:HD11	1.68	0.58
1:D:304:THR:HG23	1:D:346:PRO:HB2	1.85	0.58
1:C:234:LYS:HG2	1:C:474:ASP:OD2	2.04	0.56
1:C:93:ASP:OD1	1:C:94:ASP:N	2.39	0.56
1:A:321:VAL:HG23	1:A:352:GLN:HE21	1.72	0.54
1:D:391:THR:OG1	1:D:393:ASP:OD1	2.17	0.53
1:A:234:LYS:HB2	1:D:233:THR:O	2.09	0.53
1:A:234:LYS:HG3	1:A:474:ASP:OD2	2.11	0.51
1:B:85:ASP:OD2	1:D:469:LYS:NZ	2.43	0.51
1:B:321:VAL:HG23	1:B:352:GLN:HE21	1.75	0.50
1:A:12:LEU:HD23	1:A:96:PHE:HZ	1.77	0.50
1:A:304:THR:HG23	1:A:346:PRO:HB2	1.93	0.50
1:C:321:VAL:HG23	1:C:352:GLN:HE21	1.76	0.50
1:A:429:ARG:NH2	1:A:449:ASP:OD2	2.43	0.49
1:C:391:THR:OG1	1:C:393:ASP:OD1	2.22	0.48

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:151:ILE:HG13	4:D:633:HOH:O	2.14	0.48
1:A:151:ILE:HG13	4:A:689:HOH:O	2.14	0.47
1:C:23:TYR:OH	1:C:24:LYS:HE3	2.15	0.47
1:C:37:PHE:CZ	1:C:397:CYS:HB3	2.50	0.46
1:C:151:ILE:HG13	4:C:676:HOH:O	2.14	0.46
1:C:8:GLU:N	4:C:613:HOH:O	2.48	0.45
1:D:12:LEU:HD23	1:D:96:PHE:HZ	1.83	0.44
1:A:414:PHE:HB3	1:A:427:LYS:HG2	2.00	0.43
1:B:23:TYR:CE1	1:B:97:ASN:HB2	2.53	0.43
1:A:23:TYR:CE1	1:A:97:ASN:HB2	2.53	0.43
1:D:318:LEU:HD13	1:D:364:ILE:HA	2.00	0.43
1:D:169:LYS:HG2	1:D:482:LEU:HD11	1.99	0.43
1:B:233:THR:HG23	1:B:235:ASP:O	2.19	0.42
1:D:37:PHE:CZ	1:D:397:CYS:HB3	2.54	0.42
1:D:185:GLY:O	1:D:189:LYS:HG2	2.19	0.42
1:D:179:THR:OG1	1:D:374:SER:HA	2.20	0.42
1:A:37:PHE:CZ	1:A:397:CYS:HB3	2.54	0.42
1:D:23:TYR:CE1	1:D:97:ASN:HB2	2.55	0.42
1:C:23:TYR:CE1	1:C:97:ASN:HB2	2.56	0.41
1:A:172:LEU:O	1:A:176:LEU:HB2	2.21	0.41
1:B:37:PHE:CZ	1:B:397:CYS:HB3	2.56	0.41
1:B:400:LYS:NZ	4:B:620:HOH:O	2.54	0.41
1:B:443:LEU:HD12	1:B:448:GLY:HA2	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	462/491 (94%)	452 (98%)	10 (2%)	0	100 100
1	B	462/491 (94%)	455 (98%)	7 (2%)	0	100 100

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	462/491 (94%)	452 (98%)	10 (2%)	0	100	100
1	D	462/491 (94%)	453 (98%)	9 (2%)	0	100	100
All	All	1848/1964 (94%)	1812 (98%)	36 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	410/430 (95%)	404 (98%)	6 (2%)	65	71
1	B	410/430 (95%)	407 (99%)	3 (1%)	84	88
1	C	410/430 (95%)	406 (99%)	4 (1%)	76	82
1	D	410/430 (95%)	407 (99%)	3 (1%)	84	88
All	All	1640/1720 (95%)	1624 (99%)	16 (1%)	76	82

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	TYR
1	A	176	LEU
1	A	177	LEU
1	A	195	TYR
1	A	234	LYS
1	A	255	LYS
1	B	18	TYR
1	B	53	LYS
1	B	195	TYR
1	C	18	TYR
1	C	195	TYR
1	C	219	ASP
1	C	234	LYS
1	D	18	TYR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	195	TYR
1	D	343	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	352	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	PO4	C	502	-	4,4,4	0.95	0	6,6,6	0.44	0
3	PO4	B	502	-	4,4,4	0.94	0	6,6,6	0.45	0
3	PO4	D	502	-	4,4,4	0.91	0	6,6,6	0.44	0
3	PO4	A	502	-	4,4,4	0.95	0	6,6,6	0.47	0
2	SJ6	C	501	-	26,29,29	1.37	3 (11%)	35,40,40	1.27	3 (8%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SJ6	B	501	-	26,29,29	1.35	3 (11%)	35,40,40	1.31	4 (11%)
3	PO4	A	504	-	4,4,4	0.85	0	6,6,6	0.41	0
3	PO4	D	504	-	4,4,4	0.89	0	6,6,6	0.42	0
3	PO4	C	504	-	4,4,4	0.85	0	6,6,6	0.43	0
3	PO4	C	503	-	4,4,4	0.93	0	6,6,6	0.44	0
2	SJ6	A	501	-	26,29,29	1.38	3 (11%)	35,40,40	1.27	3 (8%)
3	PO4	B	504	-	4,4,4	0.82	0	6,6,6	0.44	0
2	SJ6	D	501	-	26,29,29	1.40	3 (11%)	35,40,40	1.20	3 (8%)
3	PO4	D	503	-	4,4,4	0.97	0	6,6,6	0.47	0
3	PO4	A	503	-	4,4,4	0.91	0	6,6,6	0.46	0
3	PO4	B	503	-	4,4,4	0.92	0	6,6,6	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SJ6	A	501	-	-	0/11/13/13	0/4/4/4
2	SJ6	D	501	-	-	0/11/13/13	0/4/4/4
2	SJ6	C	501	-	-	0/11/13/13	0/4/4/4
2	SJ6	B	501	-	-	0/11/13/13	0/4/4/4

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	501	SJ6	C7-N9	5.49	1.45	1.33
2	C	501	SJ6	C7-N9	5.37	1.45	1.33
2	A	501	SJ6	C7-N9	5.36	1.45	1.33
2	B	501	SJ6	C7-N9	5.14	1.45	1.33
2	D	501	SJ6	C22-C19	2.36	1.42	1.37
2	A	501	SJ6	C22-C19	2.35	1.42	1.37
2	B	501	SJ6	C22-C19	2.32	1.42	1.37
2	C	501	SJ6	C22-C19	2.26	1.42	1.37
2	B	501	SJ6	O8-C7	-2.17	1.18	1.23
2	A	501	SJ6	O8-C7	-2.09	1.19	1.23
2	C	501	SJ6	O8-C7	-2.08	1.19	1.23
2	D	501	SJ6	O8-C7	-2.05	1.19	1.23

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	SJ6	C10-C11-N9	3.57	120.70	113.05
2	A	501	SJ6	C10-C11-N9	3.43	120.40	113.05
2	C	501	SJ6	C10-C11-N9	2.87	119.20	113.05
2	D	501	SJ6	C10-C11-N9	2.85	119.15	113.05
2	B	501	SJ6	C1-C6-C5	-2.73	117.11	120.34
2	C	501	SJ6	C1-C6-C5	-2.63	117.23	120.34
2	A	501	SJ6	C1-C6-C5	-2.55	117.33	120.34
2	D	501	SJ6	C2-N3-C4	2.22	120.68	116.85
2	A	501	SJ6	C2-N3-C4	2.16	120.59	116.85
2	C	501	SJ6	C2-N3-C4	2.15	120.57	116.85
2	B	501	SJ6	C17-N21-C20	2.08	107.89	103.78
2	D	501	SJ6	C1-C6-C5	-2.03	117.95	120.34
2	B	501	SJ6	C2-N3-C4	2.01	120.33	116.85


There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	466/491 (94%)	0.36	37 (7%) <b>12</b> <b>16</b>	26, 38, 55, 87	0
1	B	466/491 (94%)	0.51	40 (8%) <b>10</b> <b>13</b>	28, 42, 63, 88	0
1	C	466/491 (94%)	0.54	31 (6%) <b>17</b> <b>22</b>	29, 44, 66, 91	0
1	D	466/491 (94%)	0.90	63 (13%) <b>3</b> <b>4</b>	32, 55, 79, 94	0
All	All	1864/1964 (94%)	0.58	171 (9%) <b>9</b> <b>11</b>	26, 44, 73, 94	0

All (171) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	452	GLU	6.3
1	B	409	LEU	5.9
1	D	340	GLY	5.8
1	D	436	PRO	5.7
1	C	453	TYR	5.7
1	D	437	ALA	5.3
1	D	271	SER	5.3
1	B	338	SER	5.2
1	D	450	LEU	5.1
1	A	305	GLN	4.9
1	C	53	LYS	4.9
1	D	438	GLY	4.8
1	D	453	TYR	4.8
1	D	183	LEU	4.8
1	B	449	ASP	4.7
1	D	454	GLY	4.6
1	C	450	LEU	4.5
1	B	446	GLY	4.4
1	D	272	VAL	4.4
1	B	148	ILE	4.4
1	D	337	ASN	4.4

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	D	434	ARG	4.3
1	A	452	GLU	4.3
1	D	326	GLU	4.2
1	A	148	ILE	4.1
1	B	453	TYR	4.1
1	D	455	GLN	4.0
1	B	18	TYR	4.0
1	B	448	GLY	4.0
1	A	234	LYS	4.0
1	D	452	GLU	3.9
1	D	184	ASP	3.8
1	B	450	LEU	3.8
1	C	451	GLU	3.7
1	D	178	GLU	3.7
1	B	451	GLU	3.6
1	C	437	ALA	3.6
1	D	305	GLN	3.6
1	D	435	THR	3.6
1	A	450	LEU	3.5
1	D	351	ILE	3.5
1	D	236	PRO	3.4
1	B	152	LEU	3.4
1	A	152	LEU	3.3
1	C	448	GLY	3.3
1	A	134	VAL	3.3
1	D	257	HIS	3.3
1	D	256	ASP	3.3
1	C	454	GLY	3.1
1	A	233	THR	3.1
1	B	147	TRP	3.1
1	A	235	ASP	3.1
1	A	53	LYS	3.1
1	C	93	ASP	3.0
1	C	94	ASP	3.0
1	A	454	GLY	3.0
1	D	451	GLU	3.0
1	C	436	PRO	2.9
1	C	455	GLN	2.9
1	B	19	LYS	2.9
1	D	152	LEU	2.9
1	B	58	VAL	2.9
1	C	8	GLU	2.9

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	D	339	LYS	2.9
1	B	153	VAL	2.9
1	B	59	PHE	2.9
1	D	187	GLU	2.9
1	C	435	THR	2.9
1	D	179	THR	2.9
1	A	236	PRO	2.9
1	D	245	ALA	2.9
1	D	295	LEU	2.9
1	A	303	SER	2.8
1	B	151	ILE	2.8
1	B	452	GLU	2.8
1	D	234	LYS	2.8
1	A	399	PHE	2.8
1	B	42	LYS	2.8
1	D	94	ASP	2.8
1	D	367	GLY	2.8
1	C	99	LYS	2.8
1	B	351	ILE	2.8
1	C	159	ILE	2.8
1	C	152	LEU	2.8
1	A	237	VAL	2.7
1	A	350	VAL	2.7
1	D	255	LYS	2.7
1	D	409	LEU	2.7
1	A	451	GLU	2.7
1	B	132	PHE	2.7
1	A	351	ILE	2.7
1	D	148	ILE	2.7
1	C	449	ASP	2.7
1	C	92	GLN	2.7
1	D	270	SER	2.6
1	D	304	THR	2.6
1	B	146	ASN	2.6
1	B	399	PHE	2.6
1	A	94	ASP	2.6
1	D	233	THR	2.6
1	C	414	PHE	2.6
1	B	28	PRO	2.6
1	C	88	LYS	2.5
1	C	424	ARG	2.5
1	D	182	ASN	2.5

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	D	185	GLY	2.5
1	A	271	SER	2.5
1	B	339	LYS	2.5
1	C	190	LEU	2.5
1	B	150	THR	2.5
1	C	418	VAL	2.5
1	D	235	ASP	2.5
1	B	447	LYS	2.4
1	A	116	ILE	2.4
1	C	411	ILE	2.4
1	B	455	GLN	2.4
1	B	29	ASN	2.4
1	A	42	LYS	2.4
1	D	439	ASN	2.4
1	A	144	LEU	2.4
1	D	343	LEU	2.4
1	C	18	TYR	2.4
1	B	234	LYS	2.4
1	A	145	THR	2.4
1	D	147	TRP	2.3
1	D	298	LEU	2.3
1	D	275	SER	2.3
1	B	379	ALA	2.3
1	B	93	ASP	2.3
1	A	193	PHE	2.3
1	D	407	ASN	2.3
1	D	118	ALA	2.3
1	C	336	GLU	2.3
1	D	92	GLN	2.3
1	C	439	ASN	2.3
1	B	62	LEU	2.2
1	C	150	THR	2.2
1	D	309	ILE	2.2
1	D	368	MET	2.2
1	D	35	SER	2.2
1	A	276	VAL	2.2
1	A	453	TYR	2.2
1	B	419	ALA	2.2
1	C	145	THR	2.2
1	A	238	PRO	2.2
1	D	244	ALA	2.2
1	A	310	ILE	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	89	GLU	2.1
1	D	267	THR	2.1
1	B	8	GLU	2.1
1	D	483	ASN	2.1
1	D	466	LYS	2.1
1	A	35	SER	2.1
1	A	304	THR	2.1
1	D	259	LYS	2.1
1	D	449	ASP	2.1
1	B	37	PHE	2.1
1	B	118	ALA	2.1
1	D	293	GLU	2.1
1	A	245	ALA	2.1
1	B	309	ILE	2.1
1	A	449	ASP	2.1
1	A	18	TYR	2.1
1	A	381	GLY	2.1
1	A	455	GLN	2.0
1	B	145	THR	2.0
1	D	190	LEU	2.0
1	A	244	ALA	2.0
1	C	338	SER	2.0
1	D	338	SER	2.0
1	D	134	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	PO4	B	504	5/5	0.90	0.19	55,57,58,64	0
3	PO4	A	504	5/5	0.91	0.19	53,63,64,73	0
2	SJ6	D	501	26/26	0.91	0.20	47,54,60,62	0
3	PO4	C	504	5/5	0.92	0.23	53,59,67,72	0
2	SJ6	A	501	26/26	0.93	0.21	34,41,44,46	0
3	PO4	C	503	5/5	0.93	0.16	54,55,55,60	0
3	PO4	B	502	5/5	0.93	0.16	39,43,48,49	0
2	SJ6	B	501	26/26	0.95	0.20	27,34,40,41	0
3	PO4	D	503	5/5	0.95	0.21	46,47,57,59	0
3	PO4	D	504	5/5	0.95	0.26	69,71,75,79	0
2	SJ6	C	501	26/26	0.96	0.15	30,35,40,43	0
3	PO4	D	502	5/5	0.97	0.10	45,47,54,57	0
3	PO4	A	502	5/5	0.98	0.15	34,38,42,42	0
3	PO4	A	503	5/5	0.98	0.09	35,35,37,39	0
3	PO4	C	502	5/5	0.99	0.14	38,38,39,40	0
3	PO4	B	503	5/5	0.99	0.16	27,28,34,35	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around SJ6 A 501:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around SJ6 B 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [i](#)

There are no such residues in this entry.