



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 17, 2023 – 04:08 PM JST

PDB ID : 8IVX  
Title : Crystal structure of NRP2 in complex with aNRP2-14 Fab fragment  
Authors : Geng, Y.; Zhai, L.  
Deposited on : 2023-03-29  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	583	 7% 65% 5% 30%
2	H	231	 90% 7%
3	L	215	 93% 6%

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6954 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called aNRP2-14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	407	3251	2069	564	602	16	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	13	HIS	-	expression tag	UNP O60462
A	14	HIS	-	expression tag	UNP O60462
A	15	HIS	-	expression tag	UNP O60462
A	16	HIS	-	expression tag	UNP O60462
A	17	HIS	-	expression tag	UNP O60462
A	18	HIS	-	expression tag	UNP O60462
A	19	GLU	-	expression tag	UNP O60462
A	20	ASN	-	expression tag	UNP O60462
A	21	LEU	-	expression tag	UNP O60462
A	22	TYR	-	expression tag	UNP O60462
A	23	PHE	-	expression tag	UNP O60462
A	24	GLN	-	expression tag	UNP O60462

- Molecule 2 is a protein called Heavy chain of antibody 14V4 Fab fragment.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	215	1621	1030	263	321	7	0	0	0

- Molecule 3 is a protein called Light chain of antibody 14V4 Fab fragment.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	L	213	1662	1034	279	341	8	0	1	0

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	H	1	Total C O 4 2 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	102	Total O 102 102	0	0
5	H	178	Total O 178 178	0	0
5	L	136	Total O 136 136	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: aNRP2-14



- Molecule 2: Heavy chain of antibody 14V4 Fab fragment



- Molecule 3: Light chain of antibody 14V4 Fab fragment



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	89.50Å 89.41Å 130.88Å 90.00° 94.18° 90.00°	Depositor
Resolution (Å)	46.65 – 1.90 46.61 – 1.90	Depositor EDS
% Data completeness (in resolution range)	97.1 (46.65-1.90) 97.1 (46.61-1.90)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.44 (at 1.90Å)	Xtrriage
Refinement program	REFMAC 5.8.0352	Depositor
R, $R_{free}$	0.191 , 0.225 0.199 , 0.231	Depositor DCC
$R_{free}$ test set	4067 reflections (5.16%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.7	Xtrriage
Anisotropy	0.393	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 49.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6954	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	45.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.74% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.27	0/3335	0.56	0/4522
2	H	0.30	0/1666	0.56	0/2278
3	L	0.28	0/1699	0.54	0/2306
All	All	0.28	0/6700	0.56	0/9106

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	239	ARG	Sidechain
1	A	263	ARG	Sidechain

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3251	0	3155	14	0
2	H	1621	0	1575	6	0
3	L	1662	0	1586	8	0
4	H	4	0	6	0	0
5	A	102	0	0	2	0
5	H	178	0	0	0	0
5	L	136	0	0	0	0
All	All	6954	0	6322	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:336:LEU:HD11	1:A:468:VAL:HG21	1.74	0.70
1:A:399:HIS:N	1:A:399:HIS:ND1	2.44	0.63
1:A:381:HIS:N	5:A:602:HOH:O	2.37	0.57
2:H:1:GLN:OE1	2:H:1:GLN:N	2.38	0.55
3:L:36:TYR:OH	3:L:89[A]:GLN:NE2	2.43	0.52
2:H:107:MET:CE	3:L:89[B]:GLN:OE1	2.58	0.51
2:H:107:MET:HE2	3:L:89[B]:GLN:OE1	2.11	0.50
3:L:196:GLU:HG3	3:L:207:VAL:HG22	1.95	0.48
1:A:284:GLU:OE1	1:A:312:HIS:HD2	1.96	0.48
1:A:173:LEU:HD12	1:A:175:CYS:SG	2.53	0.48
3:L:138:ASN:HB3	3:L:139:ASN:HD22	1.80	0.47
2:H:184:LEU:HD12	2:H:184:LEU:C	2.36	0.47
3:L:80:SER:HA	3:L:107:ILE:CD1	2.45	0.46
1:A:585:MET:HB2	1:A:585:MET:HE3	1.80	0.44
1:A:338:MET:O	1:A:426:GLY:HA3	2.18	0.43
3:L:137:LEU:HD12	3:L:137:LEU:N	2.33	0.43
1:A:523:LYS:HD2	1:A:548:LEU:HD11	2.00	0.43
1:A:373:MET:HB2	1:A:554:HIS:CE1	2.54	0.42
1:A:218:GLY:HA2	1:A:244:ILE:HG23	2.01	0.42
1:A:494:THR:CG2	1:A:495:PRO:HD2	2.50	0.42
3:L:108:LYS:HA	3:L:141:TYR:OH	2.20	0.42
1:A:455:THR:OG1	1:A:456:GLN:N	2.51	0.41
1:A:195:ASP:C	1:A:196:LEU:HD23	2.40	0.41
2:H:199:THR:HG23	2:H:216:LYS:HE3	2.01	0.41
2:H:50:LEU:C	2:H:50:LEU:HD12	2.42	0.40
1:A:149:CYS:N	5:A:607:HOH:O	2.54	0.40



There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	391/583 (67%)	372 (95%)	18 (5%)	1 (0%)	41	31
2	H	211/231 (91%)	209 (99%)	2 (1%)	0	100	100
3	L	212/215 (99%)	211 (100%)	1 (0%)	0	100	100
All	All	814/1029 (79%)	792 (97%)	21 (3%)	1 (0%)	51	42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	484	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	355/506 (70%)	349 (98%)	6 (2%)	60	57
2	H	188/200 (94%)	188 (100%)	0	100	100
3	L	190/191 (100%)	188 (99%)	2 (1%)	73	73
All	All	733/897 (82%)	725 (99%)	8 (1%)	73	73

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	196	LEU
1	A	253	MET
1	A	399	HIS
1	A	428	ARG
1	A	566	ILE
1	A	585	MET
3	L	77	ASN
3	L	200	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	307	GLN
1	A	312	HIS
2	H	13	GLN
3	L	53	ASN
3	L	139	ASN
3	L	213	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	EDO	H	301	-	3,3,3	0.18	0	2,2,2	0.16	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	H	301	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	407/583 (69%)	0.50	39 (9%) 8 9	27, 50, 89, 120	0
2	H	215/231 (93%)	0.06	1 (0%) 91 92	22, 31, 55, 91	0
3	L	213/215 (99%)	-0.06	0 100 100	26, 35, 54, 78	0
All	All	835/1029 (81%)	0.25	40 (4%) 30 33	22, 40, 79, 120	0

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	456	GLN	7.2
1	A	476	PRO	6.9
1	A	460	TRP	6.4
1	A	474	TRP	5.8
1	A	253	MET	5.6
1	A	255	VAL	5.2
1	A	594	TRP	5.0
1	A	544	GLN	4.4
1	A	535	TRP	4.0
1	A	484	GLY	3.6
1	A	322	LEU	3.6
1	A	541	PRO	3.5
1	A	488	LEU	3.4
1	A	595	THR	3.3
1	A	275	PHE	3.3
1	A	542	ARG	3.3
1	A	171	HIS	3.3
1	A	538	ILE	3.2
1	A	453	SER	3.2
1	A	475	PHE	3.1
1	A	210	TYR	3.1
1	A	230	CYS	3.1
1	A	455	THR	3.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	583	ILE	2.9
1	A	377	HIS	2.8
1	A	172	ASN	2.7
1	A	530	LEU	2.6
1	A	539	GLN	2.6
1	A	428	ARG	2.5
1	A	169	TYR	2.4
1	A	170	PRO	2.4
1	A	536	GLU	2.4
1	A	566	ILE	2.2
1	A	457	GLU	2.2
1	A	299	TYR	2.2
1	A	485	GLU	2.2
2	H	221	ASP	2.1
1	A	537	TYR	2.1
1	A	256	ALA	2.1
1	A	454	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	EDO	H	301	4/4	0.88	0.13	37,38,51,52	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.