



Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2020 – 07:27 pm BST

PDB ID : 2IYD
Title : SENP1 covalent complex with SUMO-2
Authors : Dong, C.; Naismith, J.H.
Deposited on : 2006-07-14
Resolution : 3.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1133 (3.20-3.20)
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)
RSRZ outliers	127900	1095 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	226	 57% 38% 5%
2	B	81	 52% 36% 9% ..

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2524 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SENTRIN-SPECIFIC PROTEASE 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	226	1891	1213	327	337	14	0	0	0


- Molecule 2 is a protein called SMALL UBIQUITIN-RELATED MODIFIER 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	79	633	390	115	124	4	0	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: SENTRIN-SPECIFIC PROTEASE 1



- Molecule 2: SMALL UBIQUITIN-RELATED MODIFIER 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	143.38 Å 143.38 Å 71.89 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.98 – 3.20 39.30 – 3.20	Depositor EDS
% Data completeness (in resolution range)	99.0 (46.98-3.20) 99.1 (39.30-3.20)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.55 (at 3.18 Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.272 , 0.288 0.249 , 0.273	Depositor DCC
R_{free} test set	714 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	108.4	Xtrriage
Anisotropy	0.210	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 117.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.028 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2524	wwPDB-VP
Average B, all atoms (Å ²)	107.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.34% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.41	0/1934	0.55	0/2599
2	B	0.47	1/642 (0.2%)	0.60	0/861
All	All	0.43	1/2576 (0.0%)	0.57	0/3460

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	15	ASP	CG-OD2	5.04	1.36	1.25

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1891	0	1889	60	0
2	B	633	0	618	20	0
All	All	2524	0	2507	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:53:SER:O	2:B:54:MET:HB2	1.74	0.88
1:A:557:ASN:HD22	1:A:557:ASN:N	1.77	0.82
1:A:594:ILE:HB	1:A:595:PRO:HD2	1.59	0.81
1:A:534:TRP:HB2	1:A:603:CYS:HB2	1.64	0.77
1:A:457:ILE:HD11	1:A:635:VAL:HG13	1.68	0.74
1:A:634:MET:O	1:A:638:ILE:HG12	1.88	0.74
1:A:444:LEU:O	1:A:445:SER:HB2	1.88	0.73
1:A:622:PHE:HA	1:A:626:HIS:HD2	1.56	0.71
1:A:594:ILE:HB	1:A:595:PRO:CD	2.20	0.71
1:A:557:ASN:HD22	1:A:557:ASN:H	1.41	0.69
2:B:84:ASP:HB3	2:B:86:PHE:CZ	2.30	0.65
1:A:595:PRO:HD3	1:A:620:ILE:HG21	1.78	0.64
2:B:57:ILE:O	2:B:58:ARG:HG2	1.98	0.63
1:A:545:ASN:HB3	1:A:585:GLN:HB3	1.81	0.62
2:B:59:PHE:HB3	2:B:66:ILE:HD12	1.81	0.61
1:A:430:GLU:O	1:A:434:VAL:HG23	1.99	0.61
1:A:524:LEU:HB2	1:A:539:VAL:HB	1.83	0.61
1:A:616:LYS:C	1:A:618:ARG:H	2.04	0.60
1:A:605:MET:HE2	1:A:631:ARG:HA	1.84	0.59
1:A:490:VAL:HG21	1:A:615:THR:HG22	1.85	0.58
1:A:482:SER:C	1:A:484:GLU:H	2.08	0.56
1:A:557:ASN:N	1:A:557:ASN:ND2	2.50	0.56
2:B:35:ARG:HG3	2:B:36:HIS:CD2	2.41	0.55
1:A:559:ALA:HA	1:A:562:ILE:HD12	1.89	0.54
1:A:526:VAL:O	1:A:536:LEU:HD12	2.08	0.53
1:A:481:ARG:O	1:A:481:ARG:HG2	2.09	0.53
2:B:84:ASP:HB3	2:B:86:PHE:CE2	2.43	0.53
1:A:457:ILE:O	1:A:457:ILE:HD13	2.09	0.52
1:A:484:GLU:HB3	1:A:487:LEU:HD12	1.90	0.52
2:B:47:CYS:O	2:B:51:GLY:HA2	2.10	0.52
1:A:557:ASN:ND2	1:A:557:ASN:H	2.07	0.51
2:B:57:ILE:C	2:B:58:ARG:HG2	2.31	0.51
1:A:512:TRP:CD1	2:B:62:ASP:HA	2.46	0.50
1:A:459:THR:HG21	1:A:467:ASN:HB3	1.93	0.50
2:B:53:SER:O	2:B:54:MET:CB	2.55	0.50
1:A:634:MET:HG2	1:A:638:ILE:HD11	1.93	0.49
1:A:482:SER:O	1:A:484:GLU:N	2.44	0.49
2:B:39:LEU:O	2:B:43:MET:HG3	2.12	0.49
1:A:454:ARG:HA	1:A:457:ILE:HG22	1.93	0.49
1:A:482:SER:OG	1:A:483:LYS:N	2.45	0.49
1:A:424:THR:H	1:A:427:MET:HG3	1.78	0.49
1:A:627:MET:N	1:A:628:PRO:HD2	2.28	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:88:GLN:O	2:B:89:GLN:HB2	2.12	0.48
1:A:490:VAL:HG22	1:A:523:ILE:HB	1.95	0.48
1:A:558:GLU:OE1	1:A:562:ILE:HD11	2.14	0.48
1:A:614:ILE:O	1:A:617:ASP:N	2.46	0.48
2:B:16:HIS:HA	2:B:33:ILE:O	2.14	0.47
1:A:558:GLU:O	1:A:562:ILE:HG13	2.14	0.47
1:A:533:HIS:NE2	1:A:552:MET:SD	2.88	0.46
2:B:37:THR:O	2:B:72:PRO:HD3	2.15	0.46
1:A:424:THR:N	1:A:427:MET:HG3	2.29	0.46
1:A:622:PHE:HA	1:A:626:HIS:CD2	2.44	0.46
1:A:629:TYR:OH	1:A:633:ARG:HD3	2.17	0.45
1:A:468:ASP:HB3	2:B:89:GLN:HB2	1.99	0.45
1:A:553:GLY:N	1:A:596:GLN:HE22	2.15	0.44
1:A:555:ILE:HG23	1:A:588:SER:OG	2.18	0.43
1:A:635:VAL:O	1:A:639:LEU:HD12	2.18	0.43
1:A:590:LYS:HB2	1:A:593:GLU:HG3	2.00	0.42
1:A:616:LYS:C	1:A:618:ARG:N	2.71	0.42
2:B:33:ILE:HG13	2:B:34:LYS:O	2.20	0.42
1:A:525:LEU:HD23	1:A:525:LEU:HA	1.93	0.42
1:A:641:ARG:O	1:A:641:ARG:HD2	2.20	0.42
2:B:56:GLN:O	2:B:88:GLN:HB2	2.20	0.42
1:A:539:VAL:HG22	1:A:546:ILE:HG12	2.01	0.41
1:A:471:ILE:HG13	1:A:608:CYS:SG	2.60	0.41
1:A:455:LYS:NZ	2:B:88:GLN:HE22	2.18	0.41
1:A:564:LEU:HD22	1:A:586:LEU:HD11	2.02	0.41
1:A:622:PHE:HB3	1:A:630:PHE:HE2	1.85	0.41
1:A:421:PRO:HG3	1:A:644:LEU:HB3	2.02	0.41
1:A:595:PRO:HD2	1:A:610:TYR:HE2	1.85	0.41
1:A:439:ASN:HD21	1:A:441:ASP:HB2	1.86	0.41
1:A:448:PHE:C	1:A:450:LEU:H	2.24	0.40
2:B:79:ASP:C	2:B:81:ASP:H	2.24	0.40
1:A:488:PRO:HD2	1:A:615:THR:O	2.21	0.40
1:A:605:MET:CE	1:A:631:ARG:HA	2.50	0.40
1:A:478:LEU:HD21	1:A:612:ASP:N	2.36	0.40
2:B:24:GLN:H	2:B:24:GLN:HG3	1.72	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	224/226 (99%)	193 (86%)	19 (8%)	12 (5%)	2	14
2	B	77/81 (95%)	60 (78%)	14 (18%)	3 (4%)	3	22
All	All	301/307 (98%)	253 (84%)	33 (11%)	15 (5%)	2	16

All (15) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	445	SER
1	A	483	LYS
1	A	494	ASN
2	B	54	MET
1	A	516	VAL
1	A	531	GLY
1	A	617	ASP
1	A	510	LYS
2	B	53	SER
1	A	439	ASN
1	A	482	SER
1	A	449	ARG
2	B	89	GLN
1	A	625	GLN
1	A	438	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	209/209 (100%)	187 (90%)	22 (10%)	7	28
2	B	70/72 (97%)	54 (77%)	16 (23%)	1	4
All	All	279/281 (99%)	241 (86%)	38 (14%)	3	17

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	427	MET
1	A	437	ASN
1	A	443	VAL
1	A	444	LEU
1	A	451	THR
1	A	457	ILE
1	A	466	LEU
1	A	467	ASN
1	A	482	SER
1	A	514	LYS
1	A	518	VAL
1	A	547	THR
1	A	551	SER
1	A	557	ASN
1	A	569	GLN
1	A	581	THR
1	A	587	PHE
1	A	592	GLN
1	A	603	CYS
1	A	613	CYS
1	A	618	ARG
1	A	641	ARG
2	B	14	ASN
2	B	18	ASN
2	B	24	GLN
2	B	35	ARG
2	B	40	SER
2	B	49	ARG
2	B	50	GLN
2	B	54	MET
2	B	58	ARG
2	B	64	GLN
2	B	74	GLN
2	B	75	LEU
2	B	77	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	78	GLU
2	B	81	ASP
2	B	86	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	439	ASN
1	A	440	GLN
1	A	494	ASN
1	A	557	ASN
1	A	592	GLN
1	A	596	GLN
1	A	626	HIS
2	B	14	ASN
2	B	16	HIS
2	B	18	ASN
2	B	64	GLN
2	B	88	GLN
2	B	89	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	226/226 (100%)	0.01	1 (0%) 92 89	107, 107, 107, 107	0
2	B	79/81 (97%)	0.00	0 100 100	107, 107, 107, 107	0
All	All	305/307 (99%)	0.01	1 (0%) 94 92	107, 107, 107, 107	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	622	PHE	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.