

# wwPDB X-ray Structure Validation Summary Report (i)

#### Jun 24, 2024 – 11:28 PM EDT

PDB ID	:	6IY0
Title	:	Crystal structure of conserved hypothetical protein SAV0927 from Staphylo-
		coccus aureus subsp. aureus Mu50
Authors	:	Jeong, S.; Ha, NC.
Deposited on	:	2018-12-12
Resolution	:	2.50  Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
$\mathrm{EDS}$	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
$R_{free}$	130704	$4661 \ (2.50-2.50)$
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length		Quality of chain				
			23%					
1	А	96		56%	17%	27%		
			22%					
1	В	96		56%	19%	25%		
			2%					
1	С	96		78%		10% 11%		
			6%					
1	D	96		72%		16% 12%		
1	Ε	96		81%		10% 8%		



Continued from previous page...

Mol	Chain	Length	Quality of chain				
1	F	96	.% •	71%		21%	8%
1	G	96	8%	74%		16%	10%
1	Н	96	6%	82	%	8%	9%
1	Ι	96	16% 27%	8% •	64%		
1	J	96	10%	11%	67%		



# 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 5933 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	А	70	Total	С	Ν	Ο	Se	0	0	0
	A	10	564	357	93	112	2	0	0	0
1	В	72	Total	С	Ν	Ο	Se	0	0	0
1	D	12	595	379	94	119	3	0	0	0
1	С	85	Total	С	Ν	Ο	Se	0	0	0
1	U	00	689	434	111	141	3	0	0	0
1	D	84	Total	С	Ν	Ο	Se	0	0	0
T	D	04	678	428	108	138	4	0	0	0
1	Е	88	Total	С	Ν	Ο	Se	0	1	0
		00	722	456	117	145	4	0	1	0
1	F	88	Registration Total C N	Ν	Ο	Se	0	0	0	
	1	00	715	451	116	144	4	0	0	0
1	G	86	Total	С	Ν	Ο	Se	0	0	0
	ŭ	00	697	440	112	142	3	0	0	0
1	Н	87	Total	С	Ν	Ο	Se	0	0	0
	11	01	705	445	113	143	4	0	0	0
1	Ι	35	Total	С	Ν	Ο	Se	0	0	0
	1	00	289	188	46	54	1		U	0
1	J	32	Total	С	Ν	Ο	Se	0	0	0
		02	266	170	45	49	2		U	U

• Molecule 1 is a protein called SAV0927.

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	89	LEU	-	expression tag	UNP A0A0H3JQV2
А	90	GLU	-	expression tag	UNP A0A0H3JQV2
А	91	HIS	-	expression tag	UNP A0A0H3JQV2
А	92	HIS	-	expression tag	UNP A0A0H3JQV2
А	93	HIS	-	expression tag	UNP A0A0H3JQV2
А	94	HIS	-	expression tag	UNP A0A0H3JQV2
А	95	HIS	-	expression tag	UNP A0A0H3JQV2
А	96	HIS	-	expression tag	UNP A0A0H3JQV2
В	89	LEU	-	expression tag	UNP A0A0H3JQV2



Chain	Residue	Modelled	Actual	Comment	Reference
В	90	GLU	-	expression tag	UNP A0A0H3JQV2
В	91	HIS	-	expression tag	UNP A0A0H3JQV2
В	92	HIS	-	expression tag	UNP A0A0H3JQV2
В	93	HIS	-	expression tag	UNP A0A0H3JQV2
В	94	HIS	-	expression tag	UNP A0A0H3JQV2
В	95	HIS	-	expression tag	UNP A0A0H3JQV2
В	96	HIS	-	expression tag	UNP A0A0H3JQV2
С	89	LEU	-	expression tag	UNP A0A0H3JQV2
С	90	GLU	-	expression tag	UNP A0A0H3JQV2
С	91	HIS	-	expression tag	UNP A0A0H3JQV2
С	92	HIS	-	expression tag	UNP A0A0H3JQV2
С	93	HIS	-	expression tag	UNP A0A0H3JQV2
С	94	HIS	-	expression tag	UNP A0A0H3JQV2
С	95	HIS	-	expression tag	UNP A0A0H3JQV2
С	96	HIS	-	expression tag	UNP A0A0H3JQV2
D	89	LEU	-	expression tag	UNP A0A0H3JQV2
D	90	GLU	-	expression tag	UNP A0A0H3JQV2
D	91	HIS	-	expression tag	UNP A0A0H3JQV2
D	92	HIS	-	expression tag	UNP A0A0H3JQV2
D	93	HIS	-	expression tag	UNP A0A0H3JQV2
D	94	HIS	-	expression tag	UNP A0A0H3JQV2
D	95	HIS	-	expression tag	UNP A0A0H3JQV2
D	96	HIS	-	expression tag	UNP A0A0H3JQV2
Ε	89	LEU	-	expression tag	UNP A0A0H3JQV2
Е	90	GLU	-	expression tag	UNP A0A0H3JQV2
Ε	91	HIS	-	expression tag	UNP A0A0H3JQV2
Е	92	HIS	-	expression tag	UNP A0A0H3JQV2
Ε	93	HIS	-	expression tag	UNP A0A0H3JQV2
Ε	94	HIS	-	expression tag	UNP A0A0H3JQV2
Ε	95	HIS	-	expression tag	UNP A0A0H3JQV2
Ε	96	HIS	-	expression tag	UNP A0A0H3JQV2
F	89	LEU	-	expression tag	UNP A0A0H3JQV2
F	90	GLU	-	expression tag	UNP A0A0H3JQV2
F	91	HIS	-	expression tag	UNP A0A0H3JQV2
F	92	HIS	-	expression tag	UNP A0A0H3JQV2
F	93	HIS	-	expression tag	UNP A0A0H3JQV2
F	94	HIS	-	expression tag	UNP A0A0H3JQV2
F	95	HIS	-	expression tag	UNP A0A0H3JQV2
F	96	HIS	-	expression tag	UNP A0A0H3JQV2
G	89	LEU	-	expression tag	UNP A0A0H3JQV2
G	90	GLU	-	expression tag	UNP A0A0H3JQV2
G	91	HIS	-	expression tag	UNP A0A0H3JQV2

Continued from previous page...



Chain	Residue	Modelled	Actual	Comment	Reference
G	92	HIS	-	expression tag	UNP A0A0H3JQV2
G	93	HIS	_	expression tag	UNP A0A0H3JQV2
G	94	HIS	_	expression tag	UNP A0A0H3JQV2
G	95	HIS	-	expression tag	UNP A0A0H3JQV2
G	96	HIS	-	expression tag	UNP A0A0H3JQV2
Н	89	LEU	-	expression tag	UNP A0A0H3JQV2
Н	90	GLU	-	expression tag	UNP A0A0H3JQV2
Н	91	HIS	-	expression tag	UNP A0A0H3JQV2
Н	92	HIS	-	expression tag	UNP A0A0H3JQV2
Н	93	HIS	-	expression tag	UNP A0A0H3JQV2
Н	94	HIS	-	expression tag	UNP A0A0H3JQV2
Н	95	HIS	-	expression tag	UNP A0A0H3JQV2
Н	96	HIS	-	expression tag	UNP A0A0H3JQV2
Ι	89	LEU	-	expression tag	UNP A0A0H3JQV2
Ι	90	GLU	-	expression tag	UNP A0A0H3JQV2
Ι	91	HIS	-	expression tag	UNP A0A0H3JQV2
Ι	92	HIS	-	expression tag	UNP A0A0H3JQV2
Ι	93	HIS	-	expression tag	UNP A0A0H3JQV2
Ι	94	HIS	-	expression tag	UNP A0A0H3JQV2
Ι	95	HIS	-	expression tag	UNP A0A0H3JQV2
Ι	96	HIS	-	expression tag	UNP A0A0H3JQV2
J	89	LEU	-	expression tag	UNP A0A0H3JQV2
J	90	GLU	-	expression tag	UNP A0A0H3JQV2
J	91	HIS	-	expression tag	UNP A0A0H3JQV2
J	92	HIS	-	expression tag	UNP A0A0H3JQV2
J	93	HIS	-	expression tag	UNP A0A0H3JQV2
J	94	HIS	-	expression tag	UNP A0A0H3JQV2
J	95	HIS	-	expression tag	UNP A0A0H3JQV2
J	96	HIS	-	expression tag	UNP A0A0H3JQV2

Continued from previous page...

• Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	Ε	1	Total 1	Cl 1	0	0

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	С	3	Total O 3 3	0	0




Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	D	1	Total O 1 1	0	0
3	Е	5	Total O 5 5	0	0
3	F	1	Total O 1 1	0	0
3	Н	2	Total O 2 2	0	0



# 3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: SAV0927









# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	128.18Å 76.75Å $43.57$ Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.35^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	28.80 - 2.50	Depositor
Resolution (A)	28.80 - 2.49	EDS
% Data completeness	98.2 (28.80-2.50)	Depositor
(in resolution range)	98.2 (28.80-2.49)	EDS
R <sub>merge</sub>	0.08	Depositor
R <sub>sym</sub>	(Not available)	Depositor
$< I/\sigma(I) > 1$	$5.55 (at 2.48 \text{\AA})$	Xtriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
D D.	0.237 , $0.278$	Depositor
$R, R_{free}$	0.236 , $0.278$	DCC
R <sub>free</sub> test set	2012 reflections $(6.88\%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	24.9	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.36 , $61.5$	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.50, < L^2 > = 0.33$	Xtriage
Estimated twinning fraction	0.000 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	5933	wwPDB-VP
Average B, all atoms $(Å^2)$	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 34.72 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.5309e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond	angles
	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	А	0.25	0/570	0.43	0/764
1	В	0.25	0/600	0.42	0/802
1	С	0.25	0/698	0.43	0/938
1	D	0.25	0/685	0.42	0/918
1	Е	0.25	0/732	0.43	0/984
1	F	0.25	0/725	0.42	0/974
1	G	0.25	0/706	0.45	0/949
1	Н	0.25	0/714	0.47	0/959
1	Ι	0.24	0/291	0.42	0/387
1	J	0.29	0/268	0.48	0/356
All	All	0.25	0/5989	0.44	0/8031

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	564	0	531	10	0
1	В	595	0	559	12	0
1	С	689	0	644	8	0



6IY0	
------	--

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	678	0	642	11	0
1	Е	722	0	682	9	0
1	F	715	0	674	14	0
1	G	697	0	655	9	0
1	Н	705	0	667	7	0
1	Ι	289	0	275	6	0
1	J	266	0	256	7	0
2	Ε	1	0	0	0	0
3	С	3	0	0	0	0
3	D	1	0	0	0	0
3	Ε	5	0	0	0	0
3	F	1	0	0	0	0
3	Н	2	0	0	0	0
All	All	5933	0	5585	77	0

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 77 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:J:28:ASP:OD2	1:J:47:GLN:NE2	2.15	0.79
1:B:34:THR:HG22	1:B:36:ARG:H	1.54	0.72
1:E:17:PHE:HE1	1:E:30:MSE:HE2	1.59	0.66
1:H:23:GLU:HB3	1:H:78:GLU:OE2	1.97	0.65
1:I:32:VAL:HG13	1:I:42:LEU:HB2	1.77	0.65

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



(	6IY0

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	64/96~(67%)	64 (100%)	0	0	100	100
1	В	64/96~(67%)	61~(95%)	3~(5%)	0	100	100
1	$\mathbf{C}$	83/96~(86%)	83 (100%)	0	0	100	100
1	D	80/96~(83%)	80 (100%)	0	0	100	100
1	Ε	87/96~(91%)	86~(99%)	1 (1%)	0	100	100
1	F	86/96~(90%)	83~(96%)	3~(4%)	0	100	100
1	G	84/96~(88%)	84 (100%)	0	0	100	100
1	Η	85/96~(88%)	85 (100%)	0	0	100	100
1	Ι	27/96~(28%)	25~(93%)	2~(7%)	0	100	100
1	J	28/96~(29%)	28 (100%)	0	0	100	100
All	All	688/960~(72%)	679~(99%)	9~(1%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	61/82~(74%)	61~(100%)	0	100 100
1	В	66/82~(80%)	66~(100%)	0	100 100
1	С	75/82~(92%)	74 (99%)	1 (1%)	69 87
1	D	74/82~(90%)	74 (100%)	0	100 100
1	Ε	79/82~(96%)	79~(100%)	0	100 100
1	F	78/82~(95%)	76~(97%)	2(3%)	46 72
1	G	76/82~(93%)	74 (97%)	2(3%)	46 72
1	Η	77/82~(94%)	77~(100%)	0	100 100
1	Ι	32/82~(39%)	30~(94%)	2~(6%)	18 34
1	J	30/82~(37%)	30~(100%)	0	100 100
All	All	648/820~(79%)	641 (99%)	7 (1%)	73 89



6IY0

5 of 7 residues with a non-rotameric side chain are listed below:

Mol	Chain	$\mathbf{Res}$	Type
1	G	4	MSE
1	G	84	ASN
1	Ι	42	LEU
1	Ι	36	ARG
1	F	77	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	$\mathbf{Res}$	Type
1	А	16	GLN
1	В	47	GLN
1	С	47	GLN
1	D	10	ASN
1	D	14	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.



## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	# RSRZ > 2	$OWAB(Å^2)$	Q<0.9
1	А	68/96~(70%)	1.58	22 (32%) 0 0	39, 58, 82, 92	0
1	В	69/96~(71%)	1.56	21 (30%) 0 0	34, 65, 92, 102	0
1	С	82/96~(85%)	0.05	2 (2%) 59 62	8, 26, 57, 73	0
1	D	80/96~(83%)	0.26	6 (7%) 14 14	9, 39, 68, 74	0
1	Е	84/96~(87%)	0.00	0 100 100	7, 26, 56, 65	0
1	F	84/96~(87%)	0.21	1 (1%) 79 80	15, 36, 62, 71	0
1	G	83/96~(86%)	0.72	8 (9%) 8 7	20, 54, 82, 96	0
1	Н	83/96~(86%)	0.41	6 (7%) 15 16	18,  40,  73,  87	0
1	Ι	34/96~(35%)	1.83	15 (44%) 0 0	56, 76, 92, 113	0
1	J	30/96~(31%)	1.74	10 (33%) 0 0	58, 78, 93, 101	0
All	All	697/960~(72%)	0.67	91 (13%) 3 3	7, 45, 84, 113	0

The worst 5 of 91 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	77	ASP	6.6
1	А	76	GLY	6.3
1	G	24	HIS	6.3
1	В	86	VAL	5.8
1	В	5	TYR	5.7

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
2	CL	Е	101	1/1	0.97	0.10	36, 36, 36, 36	0

### 6.5 Other polymers (i)

There are no such residues in this entry.

