



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 15, 2023 – 01:42 PM JST

PDB ID : 6J4Q  
Title : Structural basis of tubulin detyrosination by vasohibins-SVBP enzyme complex and functional implications  
Authors : Wang, N.; Bao, H.; Huang, H.  
Deposited on : 2019-01-10  
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	251	
1	C	251	
1	F	251	
1	J	251	
2	B	66	
2	D	66	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	G	66	<p>2% 35% 11% 52%</p>
2	K	66	<p>36% 8% 56%</p>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	303	-	X	-	-

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 9379 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tubuliny1-Tyr carboxypeptidase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	247	2005	1297	348	345	15	0	0	0
1	C	245	1993	1288	346	345	14	0	0	0
1	F	246	1990	1285	346	344	15	0	0	0
1	J	247	1970	1276	336	345	13	0	0	0

- Molecule 2 is a protein called Small vasohibin-binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	33	276	170	51	53	2	0	0	0
2	D	34	268	163	51	52	2	0	0	0
2	G	32	239	148	43	46	2	0	0	0
2	K	29	225	139	42	43	1	0	0	0

- Molecule 3 is N-[(2S)-4-chloro-3-oxo-1-phenyl-butan-2-yl]-4-methyl-benzenesulfonamide (three-letter code: TQ8) (formula: C<sub>17</sub>H<sub>18</sub>ClNO<sub>3</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
3	A	1	Total	C	N	O	S	0	0
			22	17	1	3	1		
3	B	1	Total	C	N	O	S	0	0
			22	17	1	3	1		
3	C	1	Total	C	N	O	S	0	0
			22	17	1	3	1		
3	F	1	Total	C	N	O	S	0	0
			22	17	1	3	1		
3	G	1	Total	C	N	O	S	0	0
			22	17	1	3	1		
3	J	1	Total	C	N	O	S	0	0
			22	17	1	3	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	C O	0	0
			6	3 3		
4	A	1	Total	C O	0	0
			6	3 3		
4	C	1	Total	C O	0	0
			6	3 3		
4	F	1	Total	C O	0	0
			6	3 3		
4	J	1	Total	C O	0	0
			6	3 3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	67	Total	O	0	0
			67	67		
5	B	2	Total	O	0	0
			2	2		
5	C	84	Total	O	0	0
			84	84		
5	D	3	Total	O	0	0
			3	3		
5	F	48	Total	O	0	0
			48	48		
5	G	4	Total	O	0	0
			4	4		
5	J	43	Total	O	0	0
			43	43		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tubuliny-Tyr carboxypeptidase 2




- Molecule 1: Tubuliny-Tyr carboxypeptidase 2




- Molecule 1: Tubuliny-Tyr carboxypeptidase 2



- Molecule 1: Tubuliny-Tyr carboxypeptidase 2






- Molecule 2: Small vasohibin-binding protein

Chain B:




- Molecule 2: Small vasohibin-binding protein

Chain D:




- Molecule 2: Small vasohibin-binding protein

Chain G:



- Molecule 2: Small vasohibin-binding protein

Chain K:





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	117.18Å 117.18Å 209.04Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.31 – 2.70 49.31 – 2.70	Depositor EDS
% Data completeness (in resolution range)	93.8 (49.31-2.70) 93.8 (49.31-2.70)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.91 (at 2.69Å)	Xtrriage
Refinement program	PHENIX (1.12_2829: ???)	Depositor
R, $R_{free}$	0.183 , 0.247 0.183 , 0.247	Depositor DCC
$R_{free}$ test set	2213 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	40.4	Xtrriage
Anisotropy	0.089	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 45.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.024 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	9379	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	50.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.24% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TQ8, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.57	1/2060 (0.0%)	0.66	1/2782 (0.0%)
1	C	0.56	0/2048	0.68	1/2766 (0.0%)
1	F	0.49	0/2045	0.63	0/2766
1	J	0.45	0/2025	0.61	0/2744
2	B	0.64	0/278	0.65	0/371
2	D	0.46	0/269	0.56	0/360
2	G	0.69	1/240 (0.4%)	0.73	1/323 (0.3%)
2	K	0.41	0/226	0.54	0/304
All	All	0.53	2/9191 (0.0%)	0.64	3/12416 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	193	HIS	C-O	-6.48	1.11	1.23
2	G	38	GLU	CD-OE2	5.17	1.31	1.25

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	220	LEU	CA-CB-CG	-9.62	93.16	115.30
1	A	136	MET	CG-SD-CE	5.10	108.36	100.20
2	G	38	GLU	OE1-CD-OE2	5.02	129.32	123.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2005	0	2007	42	0
1	C	1993	0	1991	49	0
1	F	1990	0	1972	51	0
1	J	1970	0	1929	30	0
2	B	276	0	260	9	0
2	D	268	0	249	11	0
2	G	239	0	212	6	0
2	K	225	0	201	3	0
3	A	22	0	16	1	0
3	B	22	0	16	1	0
3	C	22	0	16	1	0
3	F	22	0	16	0	0
3	G	22	0	16	0	0
3	J	22	0	16	1	0
4	A	12	0	16	3	0
4	C	6	0	8	1	0
4	F	6	0	8	0	0
4	J	6	0	8	0	0
5	A	67	0	0	3	0
5	B	2	0	0	1	0
5	C	84	0	0	7	0
5	D	3	0	0	0	0
5	F	48	0	0	6	0
5	G	4	0	0	0	0
5	J	43	0	0	2	0
All	All	9379	0	8957	185	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 185 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:67:TRP:NE1	5:F:401:HOH:O	1.92	0.92
1:A:59:ASP:OD1	1:A:62:THR:HG22	1.77	0.85

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:151:ARG:NH1	5:C:401:HOH:O	2.06	0.84
1:F:210:SER:HB3	1:F:216:MET:HG2	1.60	0.83
1:F:93:PRO:HB2	2:G:36:ARG:HH12	1.44	0.82

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	245/251 (98%)	234 (96%)	10 (4%)	1 (0%)	34	60
1	C	243/251 (97%)	231 (95%)	11 (4%)	1 (0%)	34	60
1	F	244/251 (97%)	233 (96%)	9 (4%)	2 (1%)	19	43
1	J	245/251 (98%)	232 (95%)	13 (5%)	0	100	100
2	B	31/66 (47%)	29 (94%)	2 (6%)	0	100	100
2	D	32/66 (48%)	31 (97%)	1 (3%)	0	100	100
2	G	30/66 (46%)	28 (93%)	1 (3%)	1 (3%)	4	8
2	K	27/66 (41%)	27 (100%)	0	0	100	100
All	All	1097/1268 (86%)	1045 (95%)	47 (4%)	5 (0%)	29	54

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	276	ARG
1	F	88	ALA
1	F	153	SER
2	G	47	THR
1	A	153	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	216/225 (96%)	209 (97%)	7 (3%)	39	68
1	C	215/225 (96%)	210 (98%)	5 (2%)	50	78
1	F	213/225 (95%)	203 (95%)	10 (5%)	26	54
1	J	208/225 (92%)	200 (96%)	8 (4%)	33	62
2	B	28/60 (47%)	27 (96%)	1 (4%)	35	64
2	D	26/60 (43%)	25 (96%)	1 (4%)	33	62
2	G	21/60 (35%)	19 (90%)	2 (10%)	8	20
2	K	20/60 (33%)	19 (95%)	1 (5%)	24	51
All	All	947/1140 (83%)	912 (96%)	35 (4%)	34	63

5 of 35 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	J	135	LYS
1	J	177	ARG
1	J	237	LYS
1	C	284	GLU
1	C	196	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
2	G	35	GLN
1	J	122	GLN
1	C	129	GLN
1	C	241	HIS
2	D	53	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	GOL	J	302	-	5,5,5	1.38	1 (20%)	5,5,5	0.69	0
3	TQ8	B	101	2	23,23,24	2.42	1 (4%)	28,32,33	1.82	4 (14%)
3	TQ8	J	301	1	23,23,24	2.28	1 (4%)	28,32,33	2.72	7 (25%)
3	TQ8	A	301	1	23,23,24	2.26	2 (8%)	28,32,33	3.49	8 (28%)
4	GOL	A	303	-	5,5,5	1.64	2 (40%)	5,5,5	0.74	0
4	GOL	F	302	-	5,5,5	1.28	0	5,5,5	0.75	0
3	TQ8	F	301	1	23,23,24	2.19	1 (4%)	28,32,33	2.74	5 (17%)
3	TQ8	C	301	1	23,23,24	2.17	2 (8%)	28,32,33	2.80	5 (17%)
4	GOL	A	302	-	5,5,5	1.56	2 (40%)	5,5,5	0.78	0
4	GOL	C	302	-	5,5,5	1.16	1 (20%)	5,5,5	0.91	0
3	TQ8	G	101	2	23,23,24	2.13	1 (4%)	28,32,33	2.02	5 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	J	302	-	-	2/4/4/4	-

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TQ8	B	101	2	-	0/18/19/21	0/2/2/2
3	TQ8	J	301	1	-	2/18/19/21	0/2/2/2
3	TQ8	A	301	1	-	11/18/19/21	0/2/2/2
4	GOL	A	303	-	-	4/4/4/4	-
4	GOL	F	302	-	-	2/4/4/4	-
3	TQ8	F	301	1	-	3/18/19/21	0/2/2/2
3	TQ8	C	301	1	-	8/18/19/21	0/2/2/2
4	GOL	A	302	-	-	2/4/4/4	-
4	GOL	C	302	-	-	1/4/4/4	-
3	TQ8	G	101	2	-	0/18/19/21	0/2/2/2

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	101	TQ8	C18-S12	-11.10	1.59	1.76
3	J	301	TQ8	C18-S12	-10.27	1.60	1.76
3	F	301	TQ8	C18-S12	-9.97	1.61	1.76
3	A	301	TQ8	C18-S12	-9.89	1.61	1.76
3	G	101	TQ8	C18-S12	-9.47	1.61	1.76

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	301	TQ8	O14-S12-O13	-9.87	107.42	119.55
3	F	301	TQ8	O14-S12-O13	-9.80	107.50	119.55
3	A	301	TQ8	C3-C1-N2	9.61	122.25	110.59
3	A	301	TQ8	C18-S12-N2	9.51	120.94	107.78
3	C	301	TQ8	C3-C1-N2	9.36	121.95	110.59

There are no chirality outliers.

5 of 35 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	301	TQ8	C3-C1-N2-S12
3	C	301	TQ8	C3-C1-N2-S12
4	A	303	GOL	C1-C2-C3-O3
4	A	303	GOL	O2-C2-C3-O3
4	F	302	GOL	C1-C2-C3-O3

There are no ring outliers.



6 monomers are involved in 8 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	101	TQ8	1	0
3	J	301	TQ8	1	0
3	A	301	TQ8	1	0
3	C	301	TQ8	1	0
4	A	302	GOL	3	0
4	C	302	GOL	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	247/251 (98%)	-0.50	0 <b>100</b> <b>100</b>	22, 40, 74, 90	0
1	C	245/251 (97%)	-0.49	2 (0%) 86 87	21, 41, 70, 118	0
1	F	246/251 (98%)	-0.37	6 (2%) 59 60	27, 47, 103, 127	0
1	J	247/251 (98%)	-0.41	2 (0%) 86 87	29, 51, 80, 90	0
2	B	33/66 (50%)	0.01	0 <b>100</b> <b>100</b>	36, 57, 97, 102	0
2	D	34/66 (51%)	-0.23	1 (2%) 51 52	32, 63, 106, 111	0
2	G	32/66 (48%)	-0.04	1 (3%) 49 49	62, 73, 95, 99	0
2	K	29/66 (43%)	-0.30	0 <b>100</b> <b>100</b>	44, 62, 89, 96	0
All	All	1113/1268 (87%)	-0.41	12 (1%) 80 82	21, 47, 86, 127	0

The worst 5 of 12 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	256	PRO	6.0
1	C	257	HIS	5.6
1	F	89	PHE	2.8
1	F	90	LEU	2.8
1	J	75	PRO	2.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	GOL	A	302	6/6	0.86	0.19	44,55,55,56	0
3	TQ8	B	101	22/23	0.87	0.22	55,73,88,95	0
4	GOL	J	302	6/6	0.87	0.27	58,77,81,81	0
4	GOL	C	302	6/6	0.93	0.27	58,61,61,64	0
4	GOL	A	303	6/6	0.93	0.24	47,53,55,56	0
3	TQ8	F	301	22/23	0.94	0.20	45,66,80,87	0
4	GOL	F	302	6/6	0.94	0.21	55,59,66,67	0
3	TQ8	G	101	22/23	0.94	0.20	67,75,85,95	0
3	TQ8	J	301	22/23	0.96	0.17	39,57,72,80	0
3	TQ8	A	301	22/23	0.96	0.16	35,50,63,69	0
3	TQ8	C	301	22/23	0.98	0.16	32,52,63,64	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around TQ8 F 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around TQ8 G 101:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







**Electron density around TQ8 J 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around TQ8 A 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.