



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 18, 2024 – 10:08 AM JST

PDB ID : 8J4C
Title : YeeE(TsuA)-YeeD(TsuB) complex for thiosulfate uptake
Authors : Ikei, M.; Miyazaki, R.; Monden, K.; Naito, Y.; Takeuchi, A.; Takahashi, Y.S.;
Tanaka, Y.; Ichikawa, M.; Tsukazaki, T.
Deposited on : 2023-04-19
Resolution : 3.34 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1060 (3.38-3.30)
Clashscore	141614	1111 (3.38-3.30)
Ramachandran outliers	138981	1090 (3.38-3.30)
Sidechain outliers	138945	1089 (3.38-3.30)
RSRZ outliers	127900	1028 (3.38-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	460	 10% 65% 22% 11%
1	B	460	 13% 62% 26% 11%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	THJ	A	501	-	-	-	X
2	THJ	B	501	-	-	-	X
3	OLC	A	504	-	-	-	X
3	OLC	A	505	-	-	-	X
3	OLC	A	506	-	-	-	X
3	OLC	A	507	-	-	-	X
3	OLC	A	508	-	-	-	X
3	OLC	A	511	-	-	-	X
3	OLC	B	503	-	-	-	X
3	OLC	B	506	-	-	-	X
3	OLC	B	507	-	-	-	X
3	OLC	B	508	-	-	-	X
3	OLC	B	512	-	-	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6675 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Spirochaeta thermophila YeeE(TsuA)-YeeD(TsuB),UPF0033 domain-containing protein, SirA-like domain-containing protein (chimera).


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	408	3119	2065	505	537	12	0	0	0
1	A	408	3119	2065	505	537	12	0	0	0

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	22	ALA	CYS	engineered mutation	UNP G0GAP6
B	451	GLY	-	expression tag	UNP G0GAP7
B	452	SER	-	expression tag	UNP G0GAP7
B	453	SER	-	expression tag	UNP G0GAP7
B	454	GLY	-	expression tag	UNP G0GAP7
B	455	GLU	-	expression tag	UNP G0GAP7
B	456	ASN	-	expression tag	UNP G0GAP7
B	457	LEU	-	expression tag	UNP G0GAP7
B	458	TYR	-	expression tag	UNP G0GAP7
B	459	PHE	-	expression tag	UNP G0GAP7
B	460	GLN	-	expression tag	UNP G0GAP7
A	22	ALA	CYS	engineered mutation	UNP G0GAP6
A	451	GLY	-	expression tag	UNP G0GAP7
A	452	SER	-	expression tag	UNP G0GAP7
A	453	SER	-	expression tag	UNP G0GAP7
A	454	GLY	-	expression tag	UNP G0GAP7
A	455	GLU	-	expression tag	UNP G0GAP7
A	456	ASN	-	expression tag	UNP G0GAP7
A	457	LEU	-	expression tag	UNP G0GAP7
A	458	TYR	-	expression tag	UNP G0GAP7
A	459	PHE	-	expression tag	UNP G0GAP7
A	460	GLN	-	expression tag	UNP G0GAP7


- Molecule 2 is THIOSULFATE (three-letter code: THJ) (formula: O₃S₂) (labeled as "Ligand

of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	S	0	0
			5	3	2		
2	A	1	Total	O	S	0	0
			5	3	2		

- Molecule 3 is (2R)-2,3-dihydroxypropyl (9Z)-octadec-9-enoate (three-letter code: OLC) (formula: C₂₁H₄₀O₄).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total C O 25 21 4	0	0
3	B	1	Total C O 25 21 4	0	0
3	B	1	Total C O 20 16 4	0	0
3	B	1	Total C 17 17	0	0
3	B	1	Total C O 17 13 4	0	0
3	B	1	Total C O 25 21 4	0	0
3	B	1	Total C 12 12	0	0
3	B	1	Total C O 25 21 4	0	0
3	B	1	Total C 8 8	0	0
3	B	1	Total C O 25 21 4	0	0
3	B	1	Total C O 25 21 4	0	0
3	A	1	Total C O 25 21 4	0	0
3	A	1	Total C O 25 21 4	0	0
3	A	1	Total C 14 14	0	0
3	A	1	Total C 16 16	0	0
3	A	1	Total C O 25 21 4	0	0
3	A	1	Total C O 25 21 4	0	0
3	A	1	Total C O 25 21 4	0	0
3	A	1	Total C O 25 21 4	0	0
3	A	1	Total C O 11 7 4	0	0
3	A	1	Total C O 12 8 4	0	0


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Spirochaeta thermophila YeeE(TsuA)-YeeD(TsuB),UPF0033 domain-containing protein, SirA-like domain-containing protein (chimera)



- Molecule 1: Spirochaeta thermophila YeeE(TsuA)-YeeD(TsuB),UPF0033 domain-containing protein, SirA-like domain-containing protein (chimera)





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 2 21	Depositor
Cell constants a, b, c, α , β , γ	73.98Å 102.44Å 186.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.39 – 3.34 49.39 – 3.34	Depositor EDS
% Data completeness (in resolution range)	99.1 (49.39-3.34) 77.1 (49.39-3.34)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.78 (at 3.33Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.265 , 0.311 0.265 , 0.308	Depositor DCC
R_{free} test set	2057 reflections (9.68%)	wwPDB-VP
Wilson B-factor (Å ²)	34.0	Xtrriage
Anisotropy	1.441	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 81.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	6675	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 43.88 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.6554e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: OLC, THJ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/3197	0.45	0/4344
1	B	0.26	0/3197	0.45	0/4344
All	All	0.26	0/6394	0.45	0/8688

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3119	0	3211	69	0
1	B	3119	0	3211	80	0
2	A	5	0	0	1	0
2	B	5	0	0	0	0
3	A	203	0	318	13	0
3	B	224	0	356	15	0
All	All	6675	0	7096	158	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 158 close contacts within the same asymmetric unit are listed below, sorted by

their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:201:LEU:HB3	3:A:502:OLC:H13A	1.62	0.82
1:A:435:GLU:HB3	1:A:443:ASN:HB2	1.71	0.72
1:B:424:VAL:HG22	1:B:429:SER:HB2	1.71	0.72
1:B:172:ALA:HB1	3:B:507:OLC:H15	1.72	0.70
1:A:120:THR:HA	1:A:125:PHE:HB2	1.74	0.69

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	404/460 (88%)	385 (95%)	18 (4%)	1 (0%)	47 78
1	B	404/460 (88%)	381 (94%)	22 (5%)	1 (0%)	47 78
All	All	808/920 (88%)	766 (95%)	40 (5%)	2 (0%)	47 78

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	376	VAL
1	A	376	VAL

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	321/359 (89%)	313 (98%)	8 (2%)	47	74
1	B	321/359 (89%)	316 (98%)	5 (2%)	62	81
All	All	642/718 (89%)	629 (98%)	13 (2%)	55	78

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	118	TYR
1	A	156	SER
1	A	374	TYR
1	A	263	LYS
1	A	268	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	24	ASN
1	B	320	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

23 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	OLC	A	502	-	24,24,24	0.24	0	25,25,25	0.24	0
3	OLC	B	503	-	24,24,24	0.25	0	25,25,25	0.29	0
3	OLC	A	506	-	24,24,24	0.25	0	25,25,25	0.26	0
3	OLC	A	509	-	24,24,24	0.24	0	25,25,25	0.24	0
3	OLC	B	502	-	24,24,24	0.24	0	25,25,25	0.28	0
3	OLC	B	504	-	19,19,24	0.23	0	20,20,25	0.30	0
3	OLC	A	504	-	13,13,24	0.19	0	12,12,25	0.20	0
3	OLC	B	505	-	16,16,24	0.19	0	15,15,25	0.19	0
3	OLC	B	510	-	7,7,24	0.09	0	6,6,25	0.11	0
3	OLC	A	511	-	11,11,24	0.24	0	12,12,25	0.33	0
3	OLC	A	507	-	24,24,24	0.24	0	25,25,25	0.23	0
3	OLC	B	512	-	24,24,24	0.26	0	25,25,25	0.26	0
2	THJ	A	501	-	2,4,4	1.26	0	2,6,6	0.22	0
2	THJ	B	501	-	2,4,4	1.22	0	2,6,6	0.18	0
3	OLC	A	503	-	24,24,24	0.25	0	25,25,25	0.27	0
3	OLC	A	505	-	15,15,24	0.22	0	14,14,25	0.20	0
3	OLC	B	509	-	24,24,24	0.23	0	25,25,25	0.27	0
3	OLC	A	508	-	24,24,24	0.23	0	25,25,25	0.27	0
3	OLC	B	507	-	24,24,24	0.25	0	25,25,25	0.27	0
3	OLC	A	510	-	10,10,24	0.24	0	11,11,25	0.33	0
3	OLC	B	511	-	24,24,24	0.23	0	25,25,25	0.28	0
3	OLC	B	506	-	16,16,24	0.27	0	17,17,25	0.34	0
3	OLC	B	508	-	11,11,24	0.18	0	9,10,25	0.16	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	OLC	A	502	-	-	3/24/24/24	-
3	OLC	B	503	-	-	6/24/24/24	-
3	OLC	A	506	-	-	5/24/24/24	-
3	OLC	A	509	-	-	1/24/24/24	-
3	OLC	B	502	-	-	3/24/24/24	-
3	OLC	B	504	-	-	3/19/19/24	-
3	OLC	A	504	-	-	1/11/11/24	-
3	OLC	B	505	-	-	1/14/14/24	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	OLC	B	510	-	-	1/5/5/24	-
3	OLC	A	511	-	-	0/11/11/24	-
3	OLC	A	507	-	-	2/24/24/24	-
3	OLC	B	512	-	-	3/24/24/24	-
3	OLC	A	503	-	-	3/24/24/24	-
3	OLC	A	505	-	-	5/13/13/24	-
3	OLC	B	509	-	-	4/24/24/24	-
3	OLC	A	508	-	-	5/24/24/24	-
3	OLC	B	507	-	-	7/24/24/24	-
3	OLC	A	510	-	-	0/10/10/24	-
3	OLC	B	511	-	-	4/24/24/24	-
3	OLC	B	506	-	-	3/16/16/24	-
3	OLC	B	508	-	-	1/9/9/24	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 61 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	506	OLC	O20-C21-C22-O23
3	B	507	OLC	C2-C1-O20-C21
3	B	507	OLC	O19-C1-O20-C21
3	B	509	OLC	C2-C1-O20-C21
3	B	509	OLC	O19-C1-O20-C21

There are no ring outliers.

15 monomers are involved in 28 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	502	OLC	5	0
3	A	506	OLC	3	0
3	A	509	OLC	4	0
3	B	502	OLC	2	0
3	B	504	OLC	1	0
3	B	505	OLC	1	0
3	B	510	OLC	2	0
3	B	512	OLC	2	0




Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	THJ	1	0
3	A	503	OLC	2	0
3	B	509	OLC	3	0
3	A	508	OLC	2	0
3	B	507	OLC	3	0
3	B	511	OLC	1	0
3	B	506	OLC	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


























5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	408/460 (88%)	0.56	47 (11%) 4 4	21, 47, 155, 176	0
1	B	408/460 (88%)	0.65	58 (14%) 2 2	24, 48, 165, 186	0
All	All	816/920 (88%)	0.60	105 (12%) 3 3	21, 48, 161, 186	0

The worst 5 of 105 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	377	SER	6.4
1	B	377	SER	6.3
1	B	391	ASP	6.1
1	B	426	LYS	5.5
1	B	441	GLU	5.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	OLC	B	508	12/25	0.52	0.55	29,40,55,63	0
2	THJ	A	501	5/5	0.53	0.49	42,43,96,115	0
3	OLC	A	504	14/25	0.64	0.42	33,47,61,63	0
3	OLC	A	507	25/25	0.68	0.61	24,54,93,101	0
2	THJ	B	501	5/5	0.69	0.45	47,57,76,114	0
3	OLC	B	509	25/25	0.69	0.37	32,54,90,98	0
3	OLC	A	508	25/25	0.69	0.54	20,59,114,116	0
3	OLC	B	506	17/25	0.71	0.44	20,31,47,51	0
3	OLC	A	505	16/25	0.71	0.44	8,22,44,57	0
3	OLC	B	511	25/25	0.72	0.37	24,41,88,111	0
3	OLC	B	507	25/25	0.73	0.59	37,51,70,75	0
3	OLC	A	506	25/25	0.74	0.49	10,39,82,90	0
3	OLC	B	512	25/25	0.76	0.53	9,35,69,75	0
3	OLC	A	511	12/25	0.77	0.44	18,43,69,75	0
3	OLC	A	510	11/25	0.78	0.33	6,37,56,65	0
3	OLC	B	503	25/25	0.79	0.45	17,33,46,48	0
3	OLC	B	505	17/25	0.80	0.36	13,34,69,76	0
3	OLC	B	504	20/25	0.81	0.53	31,50,62,71	0
3	OLC	A	509	25/25	0.81	0.47	13,26,80,109	0
3	OLC	A	503	25/25	0.85	0.41	13,24,57,69	0
3	OLC	B	502	25/25	0.86	0.42	13,25,46,49	0
3	OLC	A	502	25/25	0.90	0.42	12,30,49,58	0
3	OLC	B	510	8/25	0.90	0.36	3,14,15,17	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around OLC B 508:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around THJ A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around OLC A 504:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around OLC A 507:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around THJ B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around OLC B 509:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around OLC A 508:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around OLC B 506:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around OLC A 505:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around OLC B 511:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around OLC B 507:**



$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)








Electron density around OLC A 511:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around OLC B 503:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around OLC B 505:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around OLC B 504:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around OLC A 509:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around OLC A 503:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around OLC B 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around OLC A 502:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.