



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2024 – 05:05 AM EST

PDB ID : 4J6V
Title : Crystal Structure of Tyrosinase from Bacillus megaterium N205D mutant
Authors : Kanteev, M.; Goldfeder, M.; Adir, N.; Fishman, A.
Deposited on : 2013-02-12
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	303	
1	B	303	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 5037 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	286	Total	C	N	O	S	0	0	0
			2346	1492	422	424	8			
1	B	287	Total	C	N	O	S	0	0	0
			2354	1498	423	425	8			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	GLY	SER	engineered mutation	UNP B2ZB02
A	205	ASP	ASN	engineered mutation	UNP B2ZB02
A	298	HIS	-	expression tag	UNP B2ZB02
A	299	HIS	-	expression tag	UNP B2ZB02
A	300	HIS	-	expression tag	UNP B2ZB02
A	301	HIS	-	expression tag	UNP B2ZB02
A	302	HIS	-	expression tag	UNP B2ZB02
A	303	HIS	-	expression tag	UNP B2ZB02
B	2	GLY	SER	engineered mutation	UNP B2ZB02
B	205	ASP	ASN	engineered mutation	UNP B2ZB02
B	298	HIS	-	expression tag	UNP B2ZB02
B	299	HIS	-	expression tag	UNP B2ZB02
B	300	HIS	-	expression tag	UNP B2ZB02
B	301	HIS	-	expression tag	UNP B2ZB02
B	302	HIS	-	expression tag	UNP B2ZB02
B	303	HIS	-	expression tag	UNP B2ZB02

- Molecule 2 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cu	0	0
			1	1		
2	B	1	Total	Cu	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	165	Total 165	O 165	0	0
3	B	170	Total 170	O 170	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosinase



- Molecule 1: Tyrosinase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	54.90Å 78.33Å 82.16Å 90.00° 105.50° 90.00°	Depositor
Resolution (Å)	39.39 – 1.90 43.84 – 1.90	Depositor EDS
% Data completeness (in resolution range)	97.3 (39.39-1.90) 97.3 (43.84-1.90)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.19 (at 1.89Å)	Xtrriage
Refinement program	PHENIX 1.7.3_928	Depositor
R, R_{free}	0.193 , 0.210 0.192 , 0.210	Depositor DCC
R_{free} test set	2619 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	18.7	Xtrriage
Anisotropy	0.374	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 58.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5037	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
CU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.72	1/2424 (0.0%)	0.68	2/3303 (0.1%)
1	B	0.72	0/2432	0.68	1/3314 (0.0%)
All	All	0.72	1/4856 (0.0%)	0.68	3/6617 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	274	GLU	CD-OE2	5.77	1.31	1.25

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	147	GLY	N-CA-C	6.25	128.73	113.10
1	B	16	ASP	N-CA-CB	-5.54	100.62	110.60
1	A	215	MET	CG-SD-CE	5.46	108.94	100.20

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	4	LYS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2346	0	2225	45	0
1	B	2354	0	2236	59	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	165	0	0	5	0
3	B	170	0	0	2	0
All	All	5037	0	4461	103	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (103) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:4:LYS:N	1:B:4:LYS:HD2	1.56	1.17
1:A:158:GLU:N	1:A:158:GLU:OE2	1.79	1.15
1:A:61:MET:SD	1:A:184:MET:CE	2.38	1.11
1:A:4:LYS:N	1:A:4:LYS:HD3	1.50	1.09
1:B:4:LYS:HG3	1:B:280:ARG:HB3	1.30	1.07
1:B:99:GLN:HE22	1:B:165:ARG:NH1	1.50	1.07
1:A:62:SER:HB3	3:A:664:HOH:O	1.61	1.01
1:A:253:MET:SD	3:A:665:HOH:O	2.20	0.96
1:A:4:LYS:N	1:A:4:LYS:CD	2.29	0.95
1:B:99:GLN:NE2	1:B:165:ARG:HD2	1.82	0.95
1:A:61:MET:SD	1:A:184:MET:HE2	2.06	0.94
1:B:4:LYS:N	1:B:4:LYS:CD	2.32	0.88
1:B:253:MET:SD	3:B:580:HOH:O	2.30	0.88
1:B:61:MET:HE3	3:B:669:HOH:O	1.75	0.86
1:B:99:GLN:HE22	1:B:165:ARG:CZ	1.91	0.84
1:B:99:GLN:HE22	1:B:165:ARG:HH11	1.25	0.82
1:A:61:MET:CG	1:A:184:MET:HE2	2.12	0.80
1:A:61:MET:HG2	1:A:184:MET:HE2	1.63	0.79
1:B:122:LYS:O	1:B:123:ASP:HB2	1.83	0.78
1:A:158:GLU:H	1:A:158:GLU:CD	1.87	0.78
1:B:99:GLN:NE2	1:B:165:ARG:NH1	2.29	0.77

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:61:MET:SD	1:A:184:MET:HE1	2.24	0.77
1:B:4:LYS:HG3	1:B:280:ARG:CB	2.15	0.72
1:B:99:GLN:NE2	1:B:165:ARG:CZ	2.52	0.72
1:A:254:LYS:HG2	1:A:255:ASN:H	1.56	0.70
1:B:99:GLN:NE2	1:B:165:ARG:CD	2.54	0.69
1:B:6:ARG:HG3	1:B:6:ARG:HH11	1.57	0.69
1:B:11:VAL:HA	1:B:14:LEU:HD12	1.75	0.69
1:B:278:ASN:HD22	1:B:281:LYS:H	1.41	0.67
1:A:194:LEU:HA	1:A:203:LEU:HD13	1.77	0.67
1:B:142:GLN:N	1:B:142:GLN:NE2	2.42	0.66
1:B:99:GLN:HE22	1:B:165:ARG:HD2	1.62	0.65
1:B:94:TRP:HE1	1:B:230:HIS:HD1	1.46	0.64
1:B:14:LEU:HD11	1:B:90:PRO:HB3	1.80	0.63
1:A:61:MET:CG	1:A:184:MET:CE	2.74	0.63
1:B:278:ASN:ND2	1:B:281:LYS:H	1.98	0.62
1:B:20:ARG:O	1:B:20:ARG:HD3	2.01	0.61
1:A:243:ILE:HD12	3:A:561:HOH:O	2.02	0.59
1:B:218:VAL:N	1:B:219:PRO:HD2	2.18	0.57
1:B:4:LYS:CG	1:B:280:ARG:HB3	2.21	0.57
1:B:93:GLU:HG2	1:B:96:THR:HG23	1.86	0.57
1:A:150:LYS:O	1:A:213:GLY:HA3	2.04	0.57
1:A:4:LYS:O	1:A:285:VAL:HG22	2.04	0.56
1:A:254:LYS:HG2	1:A:255:ASN:N	2.18	0.56
1:B:99:GLN:HE22	1:B:165:ARG:CD	2.18	0.56
1:B:108:ILE:HG12	1:B:108:ILE:O	2.06	0.56
1:A:94:TRP:CE3	1:A:163:PRO:HG2	2.41	0.56
1:B:278:ASN:ND2	1:B:281:LYS:HG3	2.21	0.56
1:B:196:GLY:HA3	1:B:202:GLN:O	2.07	0.54
1:B:99:GLN:HE21	1:B:165:ARG:HD2	1.70	0.54
1:A:4:LYS:HD3	1:A:280:ARG:HH21	1.73	0.54
1:B:20:ARG:HD3	1:B:20:ARG:C	2.27	0.54
1:B:99:GLN:OE1	1:B:165:ARG:CZ	2.57	0.53
1:B:96:THR:HB	1:B:165:ARG:HH21	1.74	0.53
1:B:102:ASP:OD1	1:B:104:SER:HB3	2.08	0.52
1:B:61:MET:O	1:B:62:SER:HB2	2.09	0.52
1:A:288:ILE:O	1:A:289:GLU:C	2.47	0.52
1:B:93:GLU:OE1	1:B:96:THR:HG21	2.09	0.52
1:B:142:GLN:N	1:B:142:GLN:HE21	2.08	0.52
1:A:61:MET:HG2	1:A:184:MET:CE	2.38	0.51
1:B:196:GLY:CA	1:B:202:GLN:O	2.59	0.50
1:A:243:ILE:CD1	3:A:561:HOH:O	2.56	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:99:GLN:NE2	1:B:165:ARG:NE	2.59	0.50
1:B:36:ASP:OD2	1:B:139:ILE:HD13	2.13	0.49
1:B:14:LEU:HD11	1:B:90:PRO:CB	2.42	0.49
1:B:106:SER:OG	1:B:108:ILE:HG22	2.13	0.49
1:A:119:ASN:HD22	1:A:122:LYS:HE3	1.79	0.48
1:A:39:ILE:HG12	1:A:222:PRO:HB2	1.96	0.48
1:A:81:GLN:HG3	1:A:85:PRO:HA	1.96	0.47
1:A:196:GLY:CA	1:A:202:GLN:O	2.62	0.47
1:A:4:LYS:HE3	1:A:280:ARG:HE	1.80	0.47
1:B:185:THR:O	1:B:187:GLN:HG2	2.15	0.46
1:B:246:ARG:HA	1:B:246:ARG:HD2	1.53	0.46
1:B:6:ARG:HG3	1:B:6:ARG:NH1	2.29	0.46
1:B:20:ARG:HG2	1:B:20:ARG:HH11	1.79	0.46
1:A:4:LYS:O	1:A:285:VAL:CG2	2.63	0.46
1:B:204:HIS:HD2	1:B:230:HIS:NE2	2.14	0.46
1:B:20:ARG:C	1:B:20:ARG:CD	2.85	0.45
1:B:93:GLU:OE1	1:B:96:THR:CG2	2.65	0.45
1:A:56:ARG:HD3	1:A:62:SER:OG	2.17	0.45
1:B:99:GLN:HE22	1:B:165:ARG:NE	2.15	0.45
1:B:122:LYS:O	1:B:123:ASP:CB	2.48	0.44
1:A:96:THR:O	1:A:99:GLN:HG2	2.17	0.44
1:B:92:TRP:CG	1:B:108:ILE:CD1	3.01	0.44
1:B:99:GLN:CD	1:B:165:ARG:CZ	2.85	0.44
1:A:178:ASP:HA	1:A:189:SER:HB2	1.99	0.44
1:A:196:GLY:HA3	1:A:202:GLN:O	2.17	0.43
1:A:183:ASP:HB2	3:A:520:HOH:O	2.19	0.43
1:B:200:GLY:HA2	1:B:202:GLN:HG2	2.00	0.43
1:B:176:GLN:NE2	1:B:251:GLN:HE22	2.17	0.42
1:A:73:LEU:HD22	1:A:89:LEU:HD11	2.00	0.42
1:A:47:LYS:HE3	1:B:141:GLU:O	2.19	0.42
1:A:201:PRO:HB3	1:A:205:ASP:OD2	2.20	0.42
1:B:21:ASP:OD1	1:B:24:ARG:NH2	2.52	0.42
1:A:119:ASN:ND2	1:A:122:LYS:HE3	2.35	0.41
1:B:218:VAL:N	1:B:219:PRO:CD	2.82	0.41
1:A:221:ALA:N	1:A:222:PRO:CD	2.84	0.41
1:A:280:ARG:HG2	1:A:285:VAL:HG12	2.01	0.41
1:A:61:MET:SD	1:A:184:MET:HE3	2.47	0.41
1:A:202:GLN:O	1:A:203:LEU:HB2	2.21	0.41
1:A:196:GLY:HA2	1:A:202:GLN:O	2.21	0.41
1:A:254:LYS:O	1:A:261:ASN:CG	2.60	0.40
1:A:193:GLN:O	1:A:203:LEU:HD12	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	284/303 (94%)	272 (96%)	12 (4%)	0	100	100
1	B	285/303 (94%)	277 (97%)	8 (3%)	0	100	100
All	All	569/606 (94%)	549 (96%)	20 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	249/265 (94%)	235 (94%)	14 (6%)	21	11
1	B	250/265 (94%)	239 (96%)	11 (4%)	28	19
All	All	499/530 (94%)	474 (95%)	25 (5%)	24	15

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	LYS
1	A	5	TYR
1	A	29	LEU
1	A	81	GLN
1	A	89	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	94	TRP
1	A	99	GLN
1	A	158	GLU
1	A	169	LEU
1	A	189	SER
1	A	246	ARG
1	A	254	LYS
1	A	277	MET
1	A	281	LYS
1	B	4	LYS
1	B	7	VAL
1	B	20	ARG
1	B	94	TRP
1	B	96	THR
1	B	141	GLU
1	B	142	GLN
1	B	169	LEU
1	B	226	VAL
1	B	246	ARG
1	B	278	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	270	ASN
1	B	99	GLN
1	B	142	GLN
1	B	176	GLN
1	B	204	HIS
1	B	278	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	286/303 (94%)	0.21	17 (5%) 22 25	10, 22, 46, 96	1 (0%)
1	B	287/303 (94%)	0.17	13 (4%) 33 36	8, 21, 45, 104	1 (0%)
All	All	573/606 (94%)	0.19	30 (5%) 27 30	8, 21, 47, 104	2 (0%)

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	5	TYR	7.2
1	A	121	ILE	6.5
1	A	5	TYR	6.3
1	A	254	LYS	5.7
1	B	290	LEU	5.3
1	B	7	VAL	4.5
1	B	16	ASP	4.1
1	B	15	THR	3.9
1	A	255	ASN	3.9
1	B	158	GLU	3.3
1	A	119	ASN	3.1
1	B	289	GLU	2.7
1	B	101	GLN	2.5
1	A	288	ILE	2.5
1	A	125	ILE	2.5
1	B	20	ARG	2.5
1	B	24	ARG	2.5
1	B	255	ASN	2.4
1	A	133	ALA	2.3
1	A	99	GLN	2.3
1	A	110	SER	2.2
1	A	166	ASP	2.2
1	A	41	TRP	2.1
1	A	4	LYS	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	105	GLN	2.1
1	B	86	GLU	2.1
1	A	280	ARG	2.1
1	B	17	THR	2.1
1	A	165	ARG	2.0
1	A	122	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CU	B	401	1/1	0.92	0.13	33,33,33,33	1
2	CU	A	401	1/1	0.95	0.10	27,27,27,27	1

6.5 Other polymers [i](#)

There are no such residues in this entry.