

# wwPDB X-ray Structure Validation Summary Report (i)

Nov 15, 2023 – 01:52 PM JST

PDB ID : 6J9S

Title : Penta mutant of Lactobacillus casei lactate dehydrogenase

Authors: Arai, K.; Miyanaga, A.; Uchikoba, H.; Fushinobu, S.; Taguchi, H.

Deposited on : 2019-01-24

Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : 2.36

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$ 

CCP4 : 7.0.044 (Gargrove) roteins) : Engh & Huber (2001)

Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\#  ext{Entries},  ext{ resolution range}( ext{Å}))$
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	326	83%	6% • 10%
	11	920	4%	070 1 1070
1	В	326	89%	8% •
1	С	326	85%	6% • 8%
1	D	326	5% 84%	6% • 9%
1	Е	326	90%	5% • •
1	F	326	5% 89%	6% • •



## 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 14730 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called L-lactate dehydrogenase.

Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf	Trace
1	A	294	Total	С	N	О	S	0	0	0
1	A	294	2263	1447	375	435	6	0	U	
1	В	315	Total	С	N	О	S	0	0	0
1	D	313	2422	1547	403	466	6		0	U
1	С	299	Total	С	N	О	S	0	0	0
1		299	2302	1472	382	442	6		U	
1	D	297	Total	С	N	О	S	0	0	0
1	D	291	2287	1463	379	439	6		0	
1	Е	315	Total	С	N	О	S	0	0	0
1	12	313	2422	1547	403	466	6		0	
1	F	315	Total	С	N	О	S	0	0	0
1	I.	319	2422	1547	403	466	6		U	

There are 30 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	55	GLU	SER	engineered mutation	UNP S6C1Z0
A	56	ASP	ASN	engineered mutation	UNP S6C1Z0
A	164	LYS	GLU	engineered mutation	UNP S6C1Z0
A	223	ASN	ASP	engineered mutation	UNP S6C1Z0
A	224	LYS	ALA	engineered mutation	UNP S6C1Z0
В	55	GLU	SER	engineered mutation	UNP S6C1Z0
В	56	ASP	ASN	engineered mutation	UNP S6C1Z0
В	164	LYS	GLU	engineered mutation	UNP S6C1Z0
В	223	ASN	ASP	engineered mutation	UNP S6C1Z0
В	224	LYS	ALA	engineered mutation	UNP S6C1Z0
С	55	GLU	SER	engineered mutation	UNP S6C1Z0
С	56	ASP	ASN	engineered mutation	UNP S6C1Z0
С	164	LYS	GLU	engineered mutation	UNP S6C1Z0
С	223	ASN	ASP	engineered mutation	UNP S6C1Z0
С	224	LYS	ALA	engineered mutation	UNP S6C1Z0
D	55	GLU	SER	engineered mutation	UNP S6C1Z0
D	56	ASP	ASN	engineered mutation	UNP S6C1Z0



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled	Actual	Comment	Reference
D	164	LYS	GLU	engineered mutation	UNP S6C1Z0
D	223	ASN	ASP	engineered mutation	UNP S6C1Z0
D	224	LYS	ALA	engineered mutation	UNP S6C1Z0
E	55	GLU	SER	engineered mutation	UNP S6C1Z0
E	56	ASP	ASN	engineered mutation	UNP S6C1Z0
E	164	LYS	GLU	engineered mutation	UNP S6C1Z0
E	223	ASN	ASP	engineered mutation	UNP S6C1Z0
E	224	LYS	ALA	engineered mutation	UNP S6C1Z0
F	55	GLU	SER	engineered mutation	UNP S6C1Z0
F	56	ASP	ASN	engineered mutation	UNP S6C1Z0
F	164	LYS	GLU	engineered mutation	UNP S6C1Z0
F	223	ASN	ASP	engineered mutation	UNP S6C1Z0
F	224	LYS	ALA	engineered mutation	UNP S6C1Z0

 $\bullet$  Molecule 2 is SULFATE ION (three-letter code: SO4) (formula:  $\mathrm{O_4S}).$ 




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	
2	A	1	Total O S	0	0	
	Λ	1	5 4 1	0	U	
2	Δ	1	Total O S	0	0	
	Λ	1	5 4 1	0		
2	В	1	Total O S	0	0	
2	Б	1	5 4 1	0	0	
2	В	1	Total O S	0	0	
2	Б	1	5 4 1	0	0	
2	С	1	Total O S	0	0	
		1	5   4   1			



 $Continued\ from\ previous\ page...$ 

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
2	С	1	Total	Ο	S	0	0
		1	5	4	1	0	U
2	D	1	Total	О	S	0	0
2	D	1	5	4	1	0	U
2	D	1	Total	Ο	S	0	0
	D	1	5	4	1	U	0
2	E	1	Total	Ο	S	0	0
	П	1	5	4	1	U	U
$\frac{1}{2}$	E	1	Total	Ο	S	0	0
	ш	1	5	4	1	U	U
$\frac{1}{2}$	F	1	Total	Ο	S	0	0
	I.	ı ı	5	4	1		U
2	F	1	Total	Ο	S	0	0
	I.	1	5	4	1		U

• Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	В	1	Total C O 6 3 3	0	0
3	В	1	Total C O 6 3 3	0	0
3	С	1	Total C O 6 3 3	0	0
3	D	1	Total C O 6 3 3	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	E	1	Total C O 6 3 3	0	0
3	Е	1	Total C O 6 3 3	0	0
3	F	1	Total C O 6 3 3	0	0

### • Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	52	Total O 52 52	0	0
4	В	111	Total O 111 111	0	0
4	С	69	Total O 69 69	0	0
4	D	82	Total O 82 82	0	0
4	Е	117	Total O 117 117	0	0
4	F	79	Total O 79 79	0	0




## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: L-lactate dehydrogenase








• Molecule 1: L-lactate dehydrogenase



• Molecule 1: L-lactate dehydrogenase





## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	163.57Å 82.89Å 180.83Å	Donogitor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $91.50^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	41.07 - 2.00	Depositor
Resolution (A)	41.03 - 2.00	EDS
% Data completeness	97.0 (41.07-2.00)	Depositor
(in resolution range)	97.0 (41.03-2.00)	EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.18  (at  2.00Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
D.D.	0.209 , 0.234	Depositor
$R, R_{free}$	0.216 , $0.242$	DCC
$R_{free}$ test set	7948 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	39.6	Xtriage
Anisotropy	0.173	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.39 , 52.7	EDS
L-test for twinning <sup>2</sup>	$< L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	0.014 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	14730	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.76% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bond angles		
IVIOI	Chain	RMSZ	SZ  = # Z  > 5		# Z >5	
1	A	0.85	0/2303	0.88	5/3120 (0.2%)	
1	В	0.94	3/2466 (0.1%)	0.92	6/3343~(0.2%)	
1	С	0.86	0/2342	0.89	7/3173 (0.2%)	
1	D	0.96	$1/2327 \ (0.0\%)$	0.90	4/3152 (0.1%)	
1	Е	0.95	0/2466	0.92	6/3343~(0.2%)	
1	F	0.88	1/2466 (0.0%)	0.82	2/3343 (0.1%)	
All	All	0.91	5/14370 (0.0%)	0.89	30/19474 (0.2%)	

#### All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\operatorname{Observed}(\text{\AA})$	Ideal(A)
1	В	84	ALA	N-CA	10.20	1.66	1.46
1	В	22	SER	CB-OG	-5.88	1.34	1.42
1	D	208	GLU	CB-CG	5.04	1.61	1.52
1	F	291	GLU	CD-OE2	-5.04	1.20	1.25
1	В	62	SER	CB-OG	-5.02	1.35	1.42

The worst 5 of 30 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
1	В	83	THR	C-N-CA	-12.35	90.82	121.70
1	A	112	ILE	CG1-CB-CG2	-8.45	92.81	111.40
1	В	83	THR	O-C-N	-8.28	109.46	122.70
1	Е	145	ARG	NE-CZ-NH1	-7.50	116.55	120.30
1	A	145	ARG	NE-CZ-NH2	7.06	123.83	120.30

There are no chirality outliers.

There are no planarity outliers.



### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2263	0	2288	12	0
1	В	2422	0	2451	11	0
1	С	2302	0	2334	8	0
1	D	2287	0	2319	15	0
1	Е	2422	0	2451	9	0
1	F	2422	0	2451	12	0
2	A	10	0	0	0	0
2	В	10	0	0	1	0
2	С	10	0	0	0	0
2	D	10	0	0	0	0
2	Е	10	0	0	1	0
2	F	10	0	0	1	0
3	В	12	0	16	0	0
3	С	6	0	8	0	0
3	D	6	0	8	0	0
3	${ m E}$	12	0	16	0	0
3	F	6	0	8	0	0
4	A	52	0	0	0	0
4	В	111	0	0	0	0
4	С	69	0	0	0	0
4	D	82	0	0	2	0
4	Е	117	0	0	0	0
4	F	79	0	0	0	0
All	All	14730	0	14350	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 63 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:B:235:THR:HG21	2:B:401:SO4:O3	1.71	0.89
1:F:235:THR:HG21	2:F:401:SO4:O2	1.77	0.84
1:D:262:VAL:HG12	1:D:286:ILE:CD1	2.10	0.82
1:D:262:VAL:HG12	1:D:286:ILE:HD13	1.71	0.71



Continued from previous page...

Atom-1	1100111 1		$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:D:262:VAL:HG12	1:D:286:ILE:HD11	1.77	0.65

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	$290/326\ (89\%)$	282 (97%)	8 (3%)	0	100	100
1	В	313/326~(96%)	303 (97%)	10 (3%)	0	100	100
1	С	295/326~(90%)	288 (98%)	7 (2%)	0	100	100
1	D	293/326 (90%)	285 (97%)	8 (3%)	0	100	100
1	Е	313/326 (96%)	305 (97%)	7 (2%)	1 (0%)	41	37
1	F	313/326 (96%)	304 (97%)	9 (3%)	0	100	100
All	All	1817/1956 (93%)	1767 (97%)	49 (3%)	1 (0%)	51	49

#### All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	Ε	84	ALA

### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	244/270 (90%)	237 (97%)	7 (3%)	42 43
1	В	261/270 (97%)	258 (99%)	3 (1%)	73 78
1	С	249/270 (92%)	243 (98%)	6 (2%)	49 51
1	D	247/270 (92%)	242 (98%)	5 (2%)	55 58
1	E	261/270 (97%)	258 (99%)	3 (1%)	73 78
1	F	261/270 (97%)	256 (98%)	5 (2%)	57 61
All	All	1523/1620 (94%)	1494 (98%)	29 (2%)	57 61

5 of 29 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	С	307	SER
1	F	300	GLU
1	D	166	VAL
1	F	106	LYS
1	D	50	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	В	223	ASN
1	D	118	ASN
1	F	126	ASN
1	F	223	ASN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



### 5.6 Ligand geometry (i)

19 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trino	Chain	Res	Link	В	ond leng	$_{ m gths}$	В	ond ang	gles
IVIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SO4	F	401	-	4,4,4	0.63	0	6,6,6	0.70	0
2	SO4	F	402	_	4,4,4	0.38	0	6,6,6	0.52	0
2	SO4	A	402	_	4,4,4	0.46	0	6,6,6	0.61	0
2	SO4	D	402	_	4,4,4	1.33	1 (25%)	6,6,6	1.11	1 (16%)
3	GOL	С	403	-	5,5,5	0.97	0	5,5,5	0.54	0
2	SO4	В	401	_	4,4,4	0.27	0	6,6,6	0.74	0
2	SO4	D	401	_	4,4,4	0.44	0	6,6,6	1.14	0
2	SO4	С	402	_	4,4,4	0.43	0	6,6,6	0.97	0
3	GOL	E	403	-	5,5,5	1.54	1 (20%)	5, 5, 5	1.41	1 (20%)
2	SO4	Е	401	-	4,4,4	0.59	0	6,6,6	0.99	0
3	GOL	D	403	-	5,5,5	1.58	0	5,5,5	2.22	2 (40%)
3	GOL	F	403	-	5,5,5	0.61	0	5,5,5	0.61	0
2	SO4	В	402	_	4,4,4	0.77	0	6,6,6	0.68	0
3	GOL	В	404	_	5,5,5	0.72	0	5, 5, 5	1.37	1 (20%)
2	SO4	С	401	-	4,4,4	0.40	0	6,6,6	1.17	0
2	SO4	A	401	-	4,4,4	0.71	0	6,6,6	0.57	0
2	SO4	Е	402	-	4,4,4	0.88	0	6,6,6	1.05	0
3	GOL	В	403	-	5,5,5	0.37	0	5,5,5	0.64	0
3	GOL	Е	404	-	5,5,5	0.77	0	5,5,5	1.47	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	В	404	-	-	2/4/4/4	_
3	GOL	D	403	-	-	3/4/4/4	-
3	GOL	F	403	-	-	2/4/4/4	-
3	GOL	Ε	403	-	-	2/4/4/4	-

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	С	403	-	-	0/4/4/4	-
3	GOL	В	403	-	-	2/4/4/4	-
3	GOL	Е	404	-	-	0/4/4/4	-

#### All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	Observed(A)	$\operatorname{Ideal}( ext{\AA})$
3	Е	403	GOL	O3-C3	2.20	1.51	1.42
2	D	402	SO4	O2-S	2.02	1.56	1.46

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
3	D	403	GOL	O1-C1-C2	3.37	126.35	110.20
3	D	403	GOL	O3-C3-C2	2.96	124.37	110.20
3	В	404	GOL	O2-C2-C1	2.63	120.72	109.12
3	Е	403	GOL	O1-C1-C2	2.58	122.57	110.20
3	Е	404	GOL	O1-C1-C2	-2.22	99.53	110.20

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	В	403	GOL	C1-C2-C3-O3
3	D	403	GOL	O1-C1-C2-O2
3	D	403	GOL	O1-C1-C2-C3
3	Е	403	GOL	C1-C2-C3-O3
3	F	403	GOL	C1-C2-C3-O3

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	401	SO4	1	0
2	В	401	SO4	1	0
2	Е	401	SO4	1	0

## 5.7 Other polymers (i)

There are no such residues in this entry.



## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	# RSRZ > 2	$OWAB(A^2)$	Q < 0.9
1	A	$294/326\ (90\%)$	0.30	20 (6%) 17 16	34, 52, 78, 124	0
1	В	315/326~(96%)	0.16	13 (4%) 37 36	31, 46, 67, 87	0
1	С	299/326~(91%)	0.10	9 (3%) 50 49	33, 50, 74, 94	0
1	D	$297/326 \ (91\%)$	0.20	16 (5%) 25 24	31, 44, 75, 95	0
1	Е	315/326 (96%)	0.09	17 (5%) 25 24	32, 42, 63, 88	0
1	F	315/326 (96%)	0.20	15 (4%) 30 29	33, 51, 70, 86	0
All	All	1835/1956 (93%)	0.17	90 (4%) 29 28	31, 47, 72, 124	0

The worst 5 of 90 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	103	LYS	7.5
1	В	212	ASP	4.9
1	A	311	LYS	4.8
1	Е	210	LYS	4.6
1	D	313	LEU	4.5

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathbf{A}}^2)$	Q<0.9
3	GOL	Е	403	6/6	0.77	0.26	52,55,57,61	0
3	GOL	D	403	6/6	0.84	0.16	44,59,62,63	0
3	GOL	В	403	6/6	0.85	0.18	47,51,53,59	0
3	GOL	В	404	6/6	0.86	0.18	44,50,55,55	0
3	GOL	С	403	6/6	0.86	0.20	49,60,62,62	0
3	GOL	F	403	6/6	0.89	0.27	47,62,64,66	0
3	GOL	Е	404	6/6	0.92	0.18	44,45,47,49	0
2	SO4	Е	401	5/5	0.96	0.09	51,55,68,81	0
2	SO4	F	401	5/5	0.97	0.09	54,54,63,67	0
2	SO4	С	401	5/5	0.97	0.09	55,57,69,81	0
2	SO4	D	401	5/5	0.97	0.07	49,51,66,70	0
2	SO4	A	401	5/5	0.97	0.07	58,59,72,75	0
2	SO4	D	402	5/5	0.98	0.09	41,42,49,49	0
2	SO4	A	402	5/5	0.98	0.07	47,48,50,51	0
2	SO4	Е	402	5/5	0.99	0.07	44,45,48,49	0
2	SO4	В	402	5/5	0.99	0.10	44,45,52,53	0
2	SO4	F	402	5/5	0.99	0.13	52,52,56,62	0
2	SO4	В	401	5/5	0.99	0.06	48,48,64,70	0
2	SO4	С	402	5/5	0.99	0.09	50,51,54,56	0

## 6.5 Other polymers (i)

There are no such residues in this entry.

