



wwPDB EM Validation Summary Report ⓘ

Nov 20, 2022 – 02:59 AM EST

PDB ID : 3JB1
EMDB ID : EMD-6375
Title : Atomic model of cytoplasmic polyhedrosis virus with SAM
Authors : Yu, X.K.; Jiang, J.S.; Sun, J.C.; Zhou, Z.H.
Deposited on : 2015-07-06
Resolution : 3.10 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4.02b-467
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.3


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1058	
2	B	1333	
2	C	1333	
3	D	448	
3	E	448	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 32271 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Structural protein VP3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	1057	8434	5345	1457	1587	45	0	0


- Molecule 2 is a protein called Capsid protein VP1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	1191	9397	5937	1634	1789	37	0	0
2	C	1250	9851	6219	1712	1882	38	0	0

- Molecule 3 is a protein called Viral structural protein 5.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	D	292	2281	1449	399	425	8	0	0
3	E	292	2281	1449	399	425	8	0	0

- Molecule 4 is S-ADENOSYLMETHIONINE (three-letter code: SAM) (formula: C₁₅H₂₂N₆O₅S).




Mol	Chain	Residues	Atoms					AltConf
			Total	C	N	O	S	
4	A	1	27	15	6	5	1	0

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Structural protein VP3







● Molecule 2: Capsid protein VP1






• Molecule 2: Capsid protein VP1





• Molecule 3: Viral structural protein 5



• Molecule 3: Viral structural protein 5



M1	A135	R247	THR	GLU	SER
L2	K136	R250	GLU	PHE	VAL
Q3	L137	V251	ASN	ALA	LEU
Q4	G138	R252	PHE	GLU	
I19	N139	L252	LYS	VAL	
D22	D148	E253	GLY	PRO	
T29	D151	Y254	LEU	THR	
Q30	I152	I256	SER	THR	
F31	Y153	V257	PRO	VAL	
L35	A154	T262	ILE	TYR	
A39	H155	T266	GLU	GLU	
L44	V156	D272	ALA	ARC	
V45	E159	L273	GLN	HIS	
T48	L160	S274	LYS	GLN	
E53	D164	R275	HIS	LYS	
T54	R167	K281	GLU	LEU	
H55	V171	F282	LEU	GLN	
L56	M172	L283	THR	THR	
V66	P173	T286	GLN	GLN	
D70	A177	A292	ILE	ILE	
Q74	D180	ALA	SER	SER	
A81	S181	SER	ALA	GLN	
A81	M182	ALA	ASP	ASP	
F90	S185	ILE	ILE	ALA	
S91	L186	GLY	ASP	VAL	
R92	L189	ARG	GLY	ILE	
L93	S190	SER	SER	VAL	
L96	V193	PRO	LYS	ILE	
T101	W196	THR	THR	GLU	
L105	L199	THR	THR	ASN	
I111	D204	ILE	ILE	GLU	
N113	L213	THR	THR	TYR	
S114	R225	ALA	ALA	LYS	
E115	M226	GLY	ASN	ASN	
T116	H227	GLU	GLY	SER	
I117	L228	ASN	ILE	VAL	
N118	F229	ALA	ASP	SER	
D123	I230	LEU	GLU	GLU	
P124	T233	VAL	LEU	THR	
T133	T233	ARG	THR	ILE	
Y134	V240	HIS	ALA	ASP	
	K246	ASN	PRO	THR	
		ARG	TRP	GLN	
		TYR	THR	SER	
		MET	GLU	ASP	

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, I	Depositor
Number of particles used	44908	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	Each particle	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	25	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	60535	Depositor
Image detector	KODAK SO-163 FILM	Depositor
Maximum map value	27.021	Depositor
Minimum map value	-18.914	Depositor
Average map value	0.066	Depositor
Map value standard deviation	1.604	Depositor
Recommended contour level	4.0	Depositor
Map size (\AA)	772.8, 772.8, 772.8	wwPDB
Map dimensions	700, 700, 700	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.104, 1.104, 1.104	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SAM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.28	0/8619	0.52	3/11737 (0.0%)
2	B	0.33	0/9590	0.54	0/13056
2	C	0.34	0/10052	0.56	0/13687
3	D	0.33	0/2327	0.54	0/3163
3	E	0.30	0/2327	0.52	0/3163
All	All	0.32	0/32915	0.54	3/44806 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	186	ALA	C-N-CD	-8.20	102.57	120.60
1	A	186	ALA	C-N-CA	5.69	145.91	122.00
1	A	890	LEU	CA-CB-CG	5.39	127.70	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8434	0	8399	219	0
2	B	9397	0	9315	255	0
2	C	9851	0	9762	224	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	2281	0	2282	62	0
3	E	2281	0	2282	47	0
4	A	27	0	22	5	0
All	All	32271	0	32062	785	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 785 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:709:MET:O	2:C:715:ASN:ND2	2.09	0.85
2:C:873:TYR:HB3	2:C:898:GLN:HB2	1.60	0.83
3:D:44:LEU:HG	3:D:174:VAL:HG22	1.64	0.79
1:A:752:VAL:HG12	1:A:781:VAL:HG23	1.65	0.78
2:C:462:LEU:HD13	2:C:680:THR:HG22	1.64	0.78

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1055/1058 (100%)	1004 (95%)	45 (4%)	6 (1%)	25	59
2	B	1187/1333 (89%)	1135 (96%)	49 (4%)	3 (0%)	41	73
2	C	1246/1333 (94%)	1181 (95%)	60 (5%)	5 (0%)	34	69
3	D	290/448 (65%)	281 (97%)	8 (3%)	1 (0%)	41	73
3	E	290/448 (65%)	281 (97%)	9 (3%)	0	100	100
All	All	4068/4620 (88%)	3882 (95%)	171 (4%)	15 (0%)	38	69

5 of 15 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	186	ALA
2	C	1145	ALA
2	C	1251	VAL
2	B	1123	PRO
3	D	244	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	942/943 (100%)	829 (88%)	113 (12%)	5 20
2	B	1038/1153 (90%)	963 (93%)	75 (7%)	14 44
2	C	1089/1153 (94%)	980 (90%)	109 (10%)	7 28
3	D	240/379 (63%)	215 (90%)	25 (10%)	7 27
3	E	240/379 (63%)	222 (92%)	18 (8%)	13 42
All	All	3549/4007 (89%)	3209 (90%)	340 (10%)	12 31

5 of 340 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	546	VAL
2	C	1188	VAL
2	C	637	TYR
2	C	892	VAL
3	D	30	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
2	C	1138	HIS
2	C	1203	HIS
2	B	1138	HIS
2	C	156	GLN
2	C	293	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SAM	A	1101	-	24,29,29	1.21	3 (12%)	23,42,42	1.59	4 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SAM	A	1101	-	-	6/12/33/33	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1101	SAM	C2-N3	3.85	1.38	1.32
4	A	1101	SAM	C2-N1	2.62	1.38	1.33
4	A	1101	SAM	OXT-C	-2.11	1.23	1.30

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1101	SAM	N3-C2-N1	-5.23	120.51	128.68
4	A	1101	SAM	C3'-C2'-C1'	3.12	105.67	100.98
4	A	1101	SAM	OXT-C-O	-2.52	118.36	124.09
4	A	1101	SAM	OXT-C-CA	2.22	120.95	113.38

There are no chirality outliers.

5 of 6 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	1101	SAM	N-CA-CB-CG
4	A	1101	SAM	C-CA-CB-CG
4	A	1101	SAM	CB-CG-SD-CE
4	A	1101	SAM	CB-CG-SD-C5'
4	A	1101	SAM	CA-CB-CG-SD

There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1101	SAM	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-6375. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 350



Y Index: 350




Z Index: 350


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 327



Y Index: 300




Z Index: 286


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views [i](#)


6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 4.0. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.5 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 9715 nm^3 ; this corresponds to an approximate mass of 8776 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)



*Reported resolution corresponds to spatial frequency of 0.323 Å⁻¹

8 Fourier-Shell correlation


This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-6375 and PDB model 3JB1. Per-residue inclusion information can be found in section 3 on page 5.

9.1 Map-model overlays

9.1.1 Map-model overlay [i](#)



9.1.2 Map-model assembly overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 4.0 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (4.0).













9.4 Atom inclusion [i](#)



At the recommended contour level, 79% of all backbone atoms, 68% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary [i](#)

The table lists the average atom inclusion at the recommended contour level (4.0) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.6821	 0.4790
A	 0.5307	 0.4390
B	 0.7221	 0.4880
C	 0.7447	 0.4980
D	 0.7409	 0.5010
E	 0.7481	 0.4930

