



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 27, 2023 – 06:12 PM JST

PDB ID : 8JB1
Title : Crystal structure of glyceraldehyde-3-phosphate dehydrogenase from *Corynebacterium glutamicum* ATCC13032 in complex with NADP
Authors : Son, H.F.; Kim, K.J.
Deposited on : 2023-05-07
Resolution : 2.44 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.44 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1564 (2.46-2.42)
Clashscore	141614	1631 (2.46-2.42)
Ramachandran outliers	138981	1617 (2.46-2.42)
Sidechain outliers	138945	1617 (2.46-2.42)
RSRZ outliers	127900	1547 (2.46-2.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	488	 6% 76% 19% .
1	B	488	 4% 77% 18% . .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	B	502	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7536 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Glyceraldehyde-3-phosphate dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	468	3651	2288	657	701	5	0	0	0
1	B	468	3641	2282	653	701	5	0	0	0

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	481	LEU	-	expression tag	UNP A0A8G0FZM1
A	482	GLU	-	expression tag	UNP A0A8G0FZM1
A	483	HIS	-	expression tag	UNP A0A8G0FZM1
A	484	HIS	-	expression tag	UNP A0A8G0FZM1
A	485	HIS	-	expression tag	UNP A0A8G0FZM1
A	486	HIS	-	expression tag	UNP A0A8G0FZM1
A	487	HIS	-	expression tag	UNP A0A8G0FZM1
A	488	HIS	-	expression tag	UNP A0A8G0FZM1
B	481	LEU	-	expression tag	UNP A0A8G0FZM1
B	482	GLU	-	expression tag	UNP A0A8G0FZM1
B	483	HIS	-	expression tag	UNP A0A8G0FZM1
B	484	HIS	-	expression tag	UNP A0A8G0FZM1
B	485	HIS	-	expression tag	UNP A0A8G0FZM1
B	486	HIS	-	expression tag	UNP A0A8G0FZM1
B	487	HIS	-	expression tag	UNP A0A8G0FZM1
B	488	HIS	-	expression tag	UNP A0A8G0FZM1

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: C₂₁H₂₈N₇O₁₇P₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	A	1	48	21	7	17	3	0	0
2	B	1	48	21	7	17	3	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	A	1	6	3	3	0	0
3	B	1	6	3	3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	B	1	6	3	3	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	57	Total	O	0	0
			57	57		
4	B	73	Total	O	0	0
			73	73		

4 Data and refinement statistics

Property	Value	Source
Space group	P 43 2 2	Depositor
Cell constants a, b, c, α , β , γ	92.75Å 92.75Å 287.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	31.54 – 2.44 31.52 – 2.44	Depositor EDS
% Data completeness (in resolution range)	94.8 (31.54-2.44) 94.9 (31.52-2.44)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.74 (at 2.45Å)	Xtrriage
Refinement program	REFMAC 5.8.0403	Depositor
R, R_{free}	0.192 , 0.262 0.197 , 0.260	Depositor DCC
R_{free} test set	2189 reflections (4.85%)	wwPDB-VP
Wilson B-factor (Å ²)	32.7	Xtrriage
Anisotropy	0.362	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 37.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.26$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7536	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 52.23 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.0123e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.44	0/3712	0.83	0/5047
1	B	0.44	0/3701	0.81	0/5032
All	All	0.44	0/7413	0.82	0/10079

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	6
All	All	0	8

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (8) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	336	ASN	Peptide
1	A	410	ARG	Sidechain
1	B	162	GLY	Peptide
1	B	174	ARG	Sidechain
1	B	231	ARG	Sidechain
1	B	365	ARG	Sidechain
1	B	410	ARG	Sidechain
1	B	467	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3651	0	3642	53	0
1	B	3641	0	3628	59	0
2	A	48	0	25	0	0
2	B	48	0	25	1	0
3	A	6	0	8	1	0
3	B	12	0	16	5	0
4	A	57	0	0	0	0
4	B	73	0	0	1	0
All	All	7536	0	7344	108	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (108) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:291:PRO:HB2	1:A:406:ILE:HD11	1.24	1.10
1:B:179:HIS:HD2	3:B:502:GOL:O2	1.49	0.95
1:B:179:HIS:CD2	3:B:502:GOL:O2	2.20	0.95
1:B:30:ASN:HB2	1:B:82:THR:HG22	1.60	0.82
1:A:291:PRO:CB	1:A:406:ILE:HD11	2.11	0.77
1:B:179:HIS:HE1	1:B:449:GLU:OE1	1.68	0.75
1:A:81:GLY:HA2	1:A:118:THR:HG22	1.69	0.73
1:B:179:HIS:HD2	3:B:502:GOL:HO2	1.35	0.73
1:B:81:GLY:HA2	1:B:118:THR:HG22	1.70	0.73
1:A:266:GLY:H	1:A:269:HIS:CD2	2.10	0.69
1:A:210:PRO:HG3	1:A:232:TRP:CH2	2.29	0.67
1:B:82:THR:HG23	1:B:118:THR:HA	1.77	0.67
1:A:265:TYR:HE2	1:A:462:GLU:HG3	1.59	0.67
1:B:97:SER:HB3	1:B:98:GLU:HA	1.77	0.66
1:B:43:VAL:HG13	1:B:47:ASP:HB2	1.79	0.64
1:A:266:GLY:H	1:A:269:HIS:HD2	1.45	0.64
1:A:29:ASN:O	1:A:81:GLY:O	2.16	0.62
1:A:47:ASP:O	1:A:51:SER:HB2	1.99	0.61
1:B:57:HIS:O	1:B:398:LEU:CD1	2.49	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:301:GLY:O	1:A:380:LEU:HA	2.01	0.60
1:B:170:ALA:HB2	1:B:195:ILE:HD12	1.84	0.59
1:A:59:ILE:O	1:A:60:SER:HB2	2.03	0.59
1:A:265:TYR:CE2	1:A:462:GLU:HG3	2.38	0.59
1:B:140:ILE:HG21	1:B:457:ILE:HG13	1.86	0.58
1:B:285:THR:OG1	3:B:503:GOL:H12	2.03	0.58
1:A:32:VAL:HG21	1:A:148:TYR:CE1	2.39	0.58
1:B:434:ILE:O	1:B:440:LEU:HD12	2.03	0.58
1:B:455:GLN:OE1	1:B:458:ARG:NH1	2.37	0.58
1:A:328:ARG:HH21	1:A:328:ARG:HB2	1.68	0.58
1:B:324:HIS:CD2	1:B:326:GLY:H	2.23	0.57
1:A:187:THR:OG1	1:B:5:HIS:HE1	1.89	0.56
1:B:392:PHE:O	1:B:396:VAL:HG23	2.06	0.55
1:B:100:THR:HB	1:B:101:ASP:HA	1.88	0.55
1:B:320:ILE:HA	1:B:332:ALA:HB2	1.90	0.54
1:B:421:GLY:HA3	3:B:502:GOL:H32	1.91	0.53
1:B:446:TYR:N	1:B:446:TYR:CD1	2.76	0.53
1:A:15:ALA:O	1:A:18:MET:HB2	2.08	0.53
1:A:184:GLY:HA3	1:A:198:ASN:OD1	2.09	0.52
1:B:448:ASN:HD22	2:B:501:NAP:H72N	1.56	0.52
1:B:438:ARG:HD3	4:B:673:HOH:O	2.08	0.52
1:A:187:THR:OG1	1:B:5:HIS:CE1	2.62	0.52
1:A:179:HIS:HE1	1:A:449:GLU:OE1	1.90	0.52
1:A:455:GLN:OE1	1:A:458:ARG:NH1	2.43	0.51
1:B:324:HIS:HD2	1:B:326:GLY:H	1.56	0.51
1:B:179:HIS:CE1	1:B:449:GLU:OE1	2.58	0.51
1:B:414:VAL:CG1	1:B:419:PHE:CE2	2.95	0.50
1:B:456:VAL:O	1:B:460:VAL:HG23	2.11	0.50
1:A:52:HIS:HE1	1:A:68:SER:OG	1.94	0.50
1:B:300:TYR:OH	1:B:388:GLU:OE1	2.23	0.50
1:A:157:VAL:HA	1:A:205:ILE:O	2.12	0.50
1:A:177:SER:HA	1:B:417:THR:HG21	1.92	0.49
1:A:227:ASP:OD2	1:A:241:HIS:HE1	1.95	0.48
1:B:291:PRO:CB	1:B:406:ILE:HD11	2.44	0.48
1:A:175:ARG:HG2	1:A:175:ARG:HH21	1.78	0.48
1:A:166:LEU:HD12	1:A:204:VAL:HG21	1.95	0.48
1:A:111:ALA:N	1:A:112:PRO:CD	2.77	0.48
1:A:122:PRO:HA	1:A:150:GLY:O	2.14	0.48
1:A:221:ASN:HD22	1:A:246:GLY:HA2	1.79	0.47
1:A:52:HIS:CE1	1:A:68:SER:OG	2.68	0.47
1:B:323:PHE:CD1	1:B:324:HIS:N	2.82	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:414:VAL:HG11	1:B:419:PHE:CZ	2.50	0.47
1:A:93:SER:O	1:A:105:PHE:CZ	2.68	0.47
1:A:281:ALA:O	1:A:282:ALA:HB3	2.15	0.47
1:B:375:VAL:HG22	1:B:443:TYR:CD2	2.49	0.46
1:B:266:GLY:H	1:B:269:HIS:CD2	2.33	0.46
1:B:294:LYS:HB2	1:B:352:LEU:HD11	1.98	0.46
1:A:324:HIS:CD2	1:A:325:LYS:H	2.34	0.45
1:B:376:LEU:O	1:B:441:VAL:HA	2.17	0.45
1:B:414:VAL:CG1	1:B:419:PHE:CZ	2.99	0.45
1:A:36:PHE:CB	1:A:58:ILE:HD13	2.47	0.45
1:A:98:GLU:H	1:A:98:GLU:CD	2.20	0.45
1:B:396:VAL:HA	1:B:400:SER:HB3	1.99	0.44
1:A:36:PHE:HB3	1:A:58:ILE:HD13	1.99	0.44
1:B:301:GLY:O	1:B:380:LEU:HA	2.17	0.44
1:B:59:ILE:HG23	1:B:60:SER:N	2.33	0.44
1:A:35:VAL:HG23	1:A:40:LEU:HD11	1.99	0.43
1:A:19:VAL:N	1:A:20:PRO:HD2	2.34	0.43
1:B:238:LEU:HD11	1:B:252:LEU:HD22	2.00	0.43
1:A:198:ASN:O	1:B:5:HIS:HD2	2.01	0.43
1:A:59:ILE:O	1:A:60:SER:CB	2.66	0.43
1:A:186:ILE:HA	1:A:196:TRP:O	2.19	0.43
1:B:432:ALA:HB1	1:B:443:TYR:CD1	2.54	0.43
1:A:324:HIS:CD2	1:A:325:LYS:N	2.87	0.42
1:A:158:VAL:O	1:A:206:TYR:HA	2.18	0.42
1:B:59:ILE:HD11	1:B:92:TYR:CD1	2.54	0.42
1:B:100:THR:CB	1:B:101:ASP:HA	2.49	0.42
1:A:53:ARG:NH1	1:A:56:ARG:HH11	2.16	0.42
1:B:57:HIS:O	1:B:398:LEU:HD13	2.20	0.42
1:B:82:THR:HG23	1:B:118:THR:CA	2.48	0.42
1:A:265:TYR:HE2	1:A:462:GLU:CG	2.28	0.42
1:A:267:ILE:HD11	1:A:291:PRO:HA	2.01	0.42
1:A:308:GLU:OE2	1:A:362:ASN:HB2	2.20	0.42
1:A:412:PRO:O	1:A:413:GLU:HB2	2.19	0.42
1:B:97:SER:CB	1:B:98:GLU:HA	2.41	0.41
1:A:79:ASN:O	1:A:115:GLY:N	2.53	0.41
1:B:111:ALA:N	1:B:112:PRO:CD	2.83	0.41
1:B:291:PRO:HB2	1:B:406:ILE:HD11	2.03	0.41
1:B:471:TYR:HA	1:B:472:PRO:HA	1.86	0.41
1:B:138:ALA:O	1:B:142:VAL:HG23	2.21	0.41
1:A:127:LEU:O	1:A:156:ILE:HA	2.21	0.41
1:B:400:SER:O	1:B:403:ARG:HB2	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:446:TYR:N	1:B:446:TYR:HD1	2.18	0.41
1:B:32:VAL:HG21	1:B:148:TYR:CE1	2.55	0.41
1:A:27:ARG:HH11	1:A:27:ARG:HG2	1.86	0.40
1:B:52:HIS:HE1	1:B:68:SER:OG	2.03	0.40
1:B:266:GLY:H	1:B:269:HIS:HD2	1.70	0.40
1:A:30:ASN:HD22	1:A:118:THR:HB	1.86	0.40
1:A:179:HIS:HD2	3:A:502:GOL:O1	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	464/488 (95%)	440 (95%)	22 (5%)	2 (0%)	34	41
1	B	464/488 (95%)	428 (92%)	33 (7%)	3 (1%)	25	29
All	All	928/976 (95%)	868 (94%)	55 (6%)	5 (0%)	29	34

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	327	SER
1	A	114	ILE
1	B	274	ALA
1	B	371	VAL
1	A	371	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	398/417 (95%)	381 (96%)	17 (4%)	29	38
1	B	396/417 (95%)	383 (97%)	13 (3%)	38	49
All	All	794/834 (95%)	764 (96%)	30 (4%)	33	43

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	ASN
1	A	97	SER
1	A	100	THR
1	A	117	GLU
1	A	123	THR
1	A	165	ASP
1	A	231	ARG
1	A	294	LYS
1	A	328	ARG
1	A	341	GLU
1	A	382	THR
1	A	385	ASP
1	A	401	ASP
1	A	423	THR
1	A	446	TYR
1	A	448	ASN
1	A	470	VAL
1	B	43	VAL
1	B	58	ILE
1	B	207	SER
1	B	221	ASN
1	B	295	VAL
1	B	325	LYS
1	B	340	THR
1	B	356	GLU
1	B	382	THR
1	B	417	THR
1	B	436	THR
1	B	446	TYR
1	B	472	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (32) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	4	ASN
1	A	5	HIS
1	A	29	ASN
1	A	30	ASN
1	A	52	HIS
1	A	89	GLN
1	A	179	HIS
1	A	192	ASN
1	A	221	ASN
1	A	241	HIS
1	A	269	HIS
1	A	324	HIS
1	A	381	ASN
1	A	390	ASN
1	A	404	GLN
1	B	4	ASN
1	B	5	HIS
1	B	29	ASN
1	B	52	HIS
1	B	79	ASN
1	B	89	GLN
1	B	179	HIS
1	B	192	ASN
1	B	221	ASN
1	B	228	ASN
1	B	240	GLN
1	B	241	HIS
1	B	269	HIS
1	B	297	ASN
1	B	315	ASN
1	B	324	HIS
1	B	390	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAP	B	501	-	45,52,52	0.71	1 (2%)	56,80,80	1.04	5 (8%)
3	GOL	B	502	-	5,5,5	0.28	0	5,5,5	0.64	0
2	NAP	A	501	-	45,52,52	0.75	0	56,80,80	1.11	7 (12%)
3	GOL	B	503	-	5,5,5	0.19	0	5,5,5	0.48	0
3	GOL	A	502	-	5,5,5	0.12	0	5,5,5	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	B	501	-	-	7/31/67/67	0/5/5/5
3	GOL	B	502	-	-	4/4/4/4	-
2	NAP	A	501	-	-	6/31/67/67	0/5/5/5
3	GOL	B	503	-	-	4/4/4/4	-
3	GOL	A	502	-	-	4/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	NAP	O4D-C1D	2.04	1.43	1.41

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	NAP	O2A-PA-O1A	2.88	126.50	112.24
2	B	501	NAP	C5A-C6A-N6A	2.85	124.68	120.35
2	A	501	NAP	C5A-C6A-N6A	2.63	124.34	120.35
2	A	501	NAP	O2B-C2B-C3B	2.53	120.84	111.68
2	B	501	NAP	O3B-C3B-C2B	2.52	118.33	111.17
2	B	501	NAP	O2A-PA-O1A	2.44	124.30	112.24
2	A	501	NAP	C2N-C3N-C4N	2.40	120.98	118.26
2	B	501	NAP	C6N-N1N-C2N	-2.32	119.86	121.97
2	A	501	NAP	O3B-C3B-C4B	-2.22	104.64	111.05
2	A	501	NAP	PN-O3-PA	-2.18	125.34	132.83
2	A	501	NAP	C3N-C2N-N1N	-2.12	118.35	120.43
2	B	501	NAP	PN-O3-PA	-2.01	125.93	132.83

There are no chirality outliers.

All (25) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	501	NAP	O4D-C1D-N1N-C2N
2	A	501	NAP	O4D-C1D-N1N-C6N
2	A	501	NAP	C2D-C1D-N1N-C2N
2	A	501	NAP	C2D-C1D-N1N-C6N
2	B	501	NAP	O4D-C1D-N1N-C2N
2	B	501	NAP	O4D-C1D-N1N-C6N
2	B	501	NAP	C2D-C1D-N1N-C6N
3	A	502	GOL	C1-C2-C3-O3
3	B	502	GOL	O1-C1-C2-C3
3	B	502	GOL	C1-C2-C3-O3
3	B	503	GOL	O1-C1-C2-C3
3	B	503	GOL	C1-C2-C3-O3
2	A	501	NAP	C3B-C2B-O2B-P2B
2	B	501	NAP	C3B-C2B-O2B-P2B
3	A	502	GOL	O1-C1-C2-C3
2	B	501	NAP	C1B-C2B-O2B-P2B
3	B	502	GOL	O1-C1-C2-O2
3	B	503	GOL	O1-C1-C2-O2
3	B	503	GOL	O2-C2-C3-O3
3	A	502	GOL	O2-C2-C3-O3
3	B	502	GOL	O2-C2-C3-O3
2	A	501	NAP	O4B-C4B-C5B-O5B
2	B	501	NAP	C2D-C1D-N1N-C2N
3	A	502	GOL	O1-C1-C2-O2
2	B	501	NAP	O4B-C4B-C5B-O5B


There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	501	NAP	1	0
3	B	502	GOL	4	0
3	B	503	GOL	1	0
3	A	502	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	468/488 (95%)	-0.18	29 (6%) 20 17	15, 35, 100, 125	0
1	B	468/488 (95%)	-0.25	19 (4%) 37 34	15, 37, 94, 139	0
All	All	936/976 (95%)	-0.21	48 (5%) 28 25	15, 36, 98, 139	0

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	60	SER	7.4
1	B	61	LYS	4.7
1	A	471	TYR	4.7
1	A	60	SER	4.4
1	A	100	THR	4.2
1	A	77	ASP	4.1
1	B	108	ASP	3.8
1	A	59	ILE	3.6
1	A	66	GLU	3.6
1	A	114	ILE	3.5
1	A	104	ALA	3.5
1	A	90	LEU	3.5
1	A	3	HIS	3.5
1	B	56	ARG	3.4
1	A	58	ILE	3.3
1	A	55	ALA	3.3
1	A	79	ASN	3.3
1	B	100	THR	3.2
1	A	118	THR	3.2
1	B	97	SER	3.1
1	A	53	ARG	3.1
1	A	54	TYR	3.0
1	A	101	ASP	2.9
1	B	92	TYR	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	104	ALA	2.9
1	A	98	GLU	2.8
1	B	114	ILE	2.8
1	B	98	GLU	2.8
1	A	95	GLU	2.8
1	B	356	GLU	2.8
1	A	56	ARG	2.8
1	B	58	ILE	2.7
1	B	118	THR	2.7
1	B	115	GLY	2.7
1	B	93	SER	2.7
1	A	92	TYR	2.6
1	A	4	ASN	2.5
1	A	6	LYS	2.4
1	A	91	ALA	2.4
1	B	4	ASN	2.4
1	A	111	ALA	2.3
1	A	97	SER	2.3
1	A	89	GLN	2.3
1	A	70	ASP	2.2
1	B	77	ASP	2.2
1	B	59	ILE	2.2
1	B	99	SER	2.1
1	A	106	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GOL	B	502	6/6	0.86	0.18	32,41,46,46	0
3	GOL	B	503	6/6	0.92	0.25	52,64,67,69	0
3	GOL	A	502	6/6	0.94	0.13	38,40,43,46	0
2	NAP	B	501	48/48	0.98	0.10	17,26,33,42	0
2	NAP	A	501	48/48	0.98	0.10	14,19,24,27	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.