



wwPDB X-ray Structure Validation Summary Report ⓘ

May 13, 2020 – 12:46 pm BST

PDB ID : 1JE1
Title : 5'-DEOXY-5'-METHYLTHIOADENOSINE PHOSPHORYLASE COM-
PLEX WITH GUANOSINE AND SULFATE
Authors : Appleby, T.C.; Mathews, I.I.; Porcelli, M.; Cacciapuoti, G.; Ealick, S.E.
Deposited on : 2001-06-15
Resolution : 1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	236	
1	B	236	
1	C	236	
1	D	236	
1	E	236	
1	F	236	

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 11287 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 5'-METHYLTHIOADENOSINE PHOSPHORYLASE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	232	Total 1790	C 1149	N 295	O 341	S 5	0	0	0
1	B	231	Total 1776	C 1136	N 294	O 341	S 5	0	0	0
1	C	235	Total 1806	C 1157	N 299	O 345	S 5	0	0	0
1	D	231	Total 1782	C 1143	N 294	O 340	S 5	0	0	0
1	E	230	Total 1768	C 1130	N 293	O 340	S 5	0	0	0
1	F	234	Total 1798	C 1153	N 297	O 343	S 5	0	0	0

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	C	1	Total O S 5 4 1	0	0
2	D	1	Total O S 5 4 1	0	0
2	E	1	Total O S 5 4 1	0	0
2	F	1	Total O S 5 4 1	0	0

- Molecule 3 is GUANOSINE (three-letter code: GMP) (formula: C₁₀H₁₃N₅O₅).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O 20 10 5 5	0	0
3	B	1	Total C N O 20 10 5 5	0	0
3	C	1	Total C N O 20 10 5 5	0	0
3	D	1	Total C N O 20 10 5 5	0	0
3	E	1	Total C N O 20 10 5 5	0	0
3	F	1	Total C N O 20 10 5 5	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	75	Total 75	O 75	0	0
4	B	65	Total 65	O 65	0	0
4	C	72	Total 72	O 72	0	0
4	D	75	Total 75	O 75	0	0
4	E	64	Total 64	O 64	0	0
4	F	66	Total 66	O 66	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: 5'-METHYLTHIOADENOSINE PHOSPHORYLASE

Chain A:  82% 14%





- Molecule 1: 5'-METHYLTHIOADENOSINE PHOSPHORYLASE

Chain B:  74% 23%




- Molecule 1: 5'-METHYLTHIOADENOSINE PHOSPHORYLASE

Chain C:  82% 17%




- Molecule 1: 5'-METHYLTHIOADENOSINE PHOSPHORYLASE

Chain D:  86% 11%




S236

- Molecule 1: 5'-METHYLTHIOADENOSINE PHOSPHORYLASE

Chain E:  82% 14% ..

- Molecule 1: 5'-METHYLTHIOADENOSINE PHOSPHORYLASE

Chain F:  81% 17% ..

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	101.80Å 87.60Å 101.80Å 90.00° 119.99° 90.00°	Depositor
Resolution (Å)	20.00 – 1.80	Depositor
% Data completeness (in resolution range)	(Not available) (20.00-1.80)	Depositor
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS	Depositor
R, R_{free}	0.217 , 0.237	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	11287	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GMP, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/1822	0.61	0/2467
1	B	0.33	0/1806	0.59	0/2445
1	C	0.32	0/1839	0.59	0/2492
1	D	0.35	0/1814	0.60	0/2456
1	E	0.34	0/1798	0.59	0/2434
1	F	0.32	0/1831	0.59	0/2480
All	All	0.33	0/10910	0.60	0/14774

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1790	0	1811	27	0
1	B	1776	0	1795	43	0
1	C	1806	0	1823	33	0
1	D	1782	0	1800	24	0
1	E	1768	0	1784	23	0
1	F	1798	0	1818	36	0
2	A	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	5	0	0	0	0
2	C	5	0	0	0	0
2	D	5	0	0	0	0
2	E	5	0	0	0	0
2	F	5	0	0	0	0
3	A	20	0	13	1	0
3	B	20	0	13	1	0
3	C	20	0	13	1	0
3	D	20	0	13	1	0
3	E	20	0	13	1	0
3	F	20	0	13	1	0
4	A	75	0	0	1	0
4	B	65	0	0	1	0
4	C	72	0	0	0	0
4	D	75	0	0	0	0
4	E	64	0	0	0	0
4	F	66	0	0	1	0
All	All	11287	0	10909	185	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 185 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:164:ASP:O	1:B:167:PHE:HB3	1.76	0.85
1:E:164:ASP:O	1:E:167:PHE:HB3	1.81	0.79
1:C:25:ARG:HG3	1:C:25:ARG:HH11	1.47	0.79
1:F:25:ARG:HG3	1:F:25:ARG:HH11	1.47	0.78
1:D:215:THR:HG23	1:D:216:LYS:H	1.50	0.76

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	228/236 (97%)	220 (96%)	5 (2%)	3 (1%)	12	3
1	B	227/236 (96%)	224 (99%)	3 (1%)	0	100	100
1	C	233/236 (99%)	226 (97%)	7 (3%)	0	100	100
1	D	227/236 (96%)	219 (96%)	6 (3%)	2 (1%)	17	6
1	E	226/236 (96%)	220 (97%)	6 (3%)	0	100	100
1	F	232/236 (98%)	227 (98%)	5 (2%)	0	100	100
All	All	1373/1416 (97%)	1336 (97%)	32 (2%)	5 (0%)	34	21

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	215	THR
1	D	215	THR
1	A	114	GLY
1	A	216	LYS
1	D	114	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	193/195 (99%)	191 (99%)	2 (1%)	76	71
1	B	192/195 (98%)	187 (97%)	5 (3%)	46	32
1	C	194/195 (100%)	193 (100%)	1 (0%)	88	87
1	D	192/195 (98%)	190 (99%)	2 (1%)	76	71
1	E	191/195 (98%)	188 (98%)	3 (2%)	62	54
1	F	193/195 (99%)	191 (99%)	2 (1%)	76	71
All	All	1155/1170 (99%)	1140 (99%)	15 (1%)	69	62

5 of 15 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	219	LEU
1	C	16	ARG
1	E	216	LYS
1	B	180	GLU
1	E	180	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	123	ASN
1	B	153	ASN
1	E	35	ASN
1	B	112	GLN
1	D	123	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	C	3250	-	4,4,4	0.27	0	6,6,6	0.08	0
2	SO4	A	1250	-	4,4,4	0.28	0	6,6,6	0.07	0
3	GMP	C	3260	-	18,22,22	2.22	6 (33%)	20,33,33	2.25	6 (30%)
2	SO4	B	2250	-	4,4,4	0.31	0	6,6,6	0.07	0
3	GMP	E	5260	-	18,22,22	2.31	6 (33%)	20,33,33	2.23	6 (30%)
3	GMP	A	1260	-	18,22,22	2.30	6 (33%)	20,33,33	2.24	6 (30%)
2	SO4	D	4250	-	4,4,4	0.29	0	6,6,6	0.06	0
3	GMP	F	6260	-	18,22,22	2.28	6 (33%)	20,33,33	2.25	6 (30%)
2	SO4	E	5250	-	4,4,4	0.29	0	6,6,6	0.06	0
3	GMP	D	4260	-	18,22,22	2.30	6 (33%)	20,33,33	2.24	6 (30%)
2	SO4	F	6250	-	4,4,4	0.29	0	6,6,6	0.07	0
3	GMP	B	2260	-	18,22,22	2.36	6 (33%)	20,33,33	2.25	6 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GMP	C	3260	-	-	1/2/22/22	0/3/3/3
3	GMP	E	5260	-	-	0/2/22/22	0/3/3/3
3	GMP	A	1260	-	-	0/2/22/22	0/3/3/3
3	GMP	F	6260	-	-	2/2/22/22	0/3/3/3
3	GMP	D	4260	-	-	1/2/22/22	0/3/3/3
3	GMP	B	2260	-	-	0/2/22/22	0/3/3/3

The worst 5 of 36 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	2260	GMP	C2-N2	5.32	1.44	1.33
3	E	5260	GMP	C2-N2	5.32	1.44	1.33
3	D	4260	GMP	C2-N2	5.24	1.44	1.33
3	F	6260	GMP	C2-N2	5.14	1.44	1.33
3	C	3260	GMP	C2-N2	5.06	1.44	1.33

The worst 5 of 36 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	6260	GMP	C6-C5-C4	-5.84	115.23	120.80
3	C	3260	GMP	C6-C5-C4	-5.76	115.30	120.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	4260	GMP	C6-C5-C4	-5.75	115.31	120.80
3	B	2260	GMP	C6-C5-C4	-5.74	115.32	120.80
3	E	5260	GMP	C6-C5-C4	-5.73	115.33	120.80

There are no chirality outliers.

All (4) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	F	6260	GMP	O4'-C4'-C5'-O5'
3	F	6260	GMP	C3'-C4'-C5'-O5'
3	C	3260	GMP	O4'-C4'-C5'-O5'
3	D	4260	GMP	O4'-C4'-C5'-O5'



There are no ring outliers.



6 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	3260	GMP	1	0
3	E	5260	GMP	1	0
3	A	1260	GMP	1	0
3	F	6260	GMP	1	0
3	D	4260	GMP	1	0
3	B	2260	GMP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.