



Full wwPDB X-ray Structure Validation Report ⓘ

May 12, 2020 – 11:51 pm BST

PDB ID : 4JEP
Title : Crystal structure of *Toxoplasma gondii* nucleoside triphosphate diphosphohydrolase 1 (NTPDase1)
Authors : Krug, U.; Totzauer, R.; Strater, N.
Deposited on : 2013-02-27
Resolution : 3.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1094 (3.10-3.10)
Clashscore	141614	1184 (3.10-3.10)
Ramachandran outliers	138981	1141 (3.10-3.10)
Sidechain outliers	138945	1141 (3.10-3.10)
RSRZ outliers	127900	1067 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	611	 5% 86% 8% 6%
1	B	611	 10% 83% 7% 10%

2 Entry composition i

There is only 1 type of molecule in this entry. The entry contains 8772 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nucleoside-triphosphatase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	574	4471	2805	789	853	24	0	0	0
1	B	552	4301	2703	756	816	26	0	0	0


There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	25	MET	-	EXPRESSION TAG	UNP Q27895
A	629	GLU	-	EXPRESSION TAG	UNP Q27895
A	630	HIS	-	EXPRESSION TAG	UNP Q27895
A	631	HIS	-	EXPRESSION TAG	UNP Q27895
A	632	HIS	-	EXPRESSION TAG	UNP Q27895
A	633	HIS	-	EXPRESSION TAG	UNP Q27895
A	634	HIS	-	EXPRESSION TAG	UNP Q27895
A	635	HIS	-	EXPRESSION TAG	UNP Q27895
B	25	MET	-	EXPRESSION TAG	UNP Q27895
B	629	GLU	-	EXPRESSION TAG	UNP Q27895
B	630	HIS	-	EXPRESSION TAG	UNP Q27895
B	631	HIS	-	EXPRESSION TAG	UNP Q27895
B	632	HIS	-	EXPRESSION TAG	UNP Q27895
B	633	HIS	-	EXPRESSION TAG	UNP Q27895
B	634	HIS	-	EXPRESSION TAG	UNP Q27895
B	635	HIS	-	EXPRESSION TAG	UNP Q27895


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nucleoside-triphosphatase 2



- Molecule 1: Nucleoside-triphosphatase 2



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	72.74Å 161.52Å 236.94Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.31 – 3.10 39.49 – 3.10	Depositor EDS
% Data completeness (in resolution range)	99.4 (36.31-3.10) 99.7 (39.49-3.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.20 (at 3.12Å)	Xtrriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.270 , 0.305 0.301 , 0.342	Depositor DCC
R_{free} test set	1306 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	65.1	Xtrriage
Anisotropy	0.228	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 39.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.81	EDS
Total number of atoms	8772	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.36	0/4554	0.55	0/6161
1	B	0.37	0/4381	0.54	0/5922
All	All	0.37	0/8935	0.55	0/12083

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4471	0	4420	19	0
1	B	4301	0	4249	16	0
All	All	8772	0	8669	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:338:LYS:HB2	1:A:389:LEU:HD11	1.86	0.56
1:B:616:GLY:HA2	1:B:620:TYR:HD1	1.73	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:185:LEU:HB3	1:B:228:THR:HG23	1.89	0.54
1:A:69:ILE:HG12	1:A:78:THR:HG22	1.89	0.53
1:B:289:LEU:HD11	1:B:293:THR:HG21	1.90	0.53
1:A:185:LEU:HB3	1:A:228:THR:HG23	1.91	0.52
1:A:368:GLY:HA2	1:A:387:ASN:HA	1.92	0.52
1:B:322:VAL:HG11	1:B:471:PHE:HB2	1.91	0.51
1:A:124:ALA:HB2	1:A:138:LEU:HD21	1.93	0.50
1:A:407:VAL:HG22	1:B:476:GLN:HE21	1.76	0.49
1:B:244:LEU:HD22	1:B:275:VAL:HB	1.96	0.47
1:B:415:GLU:HG2	1:B:458:LEU:HD13	1.96	0.47
1:A:188:THR:HG22	1:A:231:ILE:HD11	1.95	0.46
1:B:124:ALA:HB2	1:B:138:LEU:HD21	1.97	0.46
1:B:269:ARG:HH21	1:B:318:ASP:HB2	1.81	0.46
1:B:309:LEU:HA	1:B:607:HIS:HD2	1.81	0.45
1:B:39:GLU:HA	1:B:42:ILE:HD12	1.98	0.45
1:A:244:LEU:HD22	1:A:275:VAL:HB	1.98	0.44
1:A:112:LEU:HD22	1:A:202:LEU:HD21	1.98	0.44
1:A:464:ILE:HG12	1:A:470:GLY:HA2	2.00	0.44
1:A:376:GLY:HA3	1:A:532:ILE:HG12	2.00	0.43
1:B:602:VAL:HA	1:B:605:ILE:HD12	1.99	0.43
1:A:181:ILE:HD12	1:A:219:TYR:HB3	2.01	0.43
1:B:322:VAL:HG13	1:B:469:SER:HB2	2.00	0.43
1:A:322:VAL:HG11	1:A:471:PHE:HB2	2.00	0.43
1:A:385:ARG:HA	1:A:390:LYS:HD2	2.00	0.43
1:B:88:CYS:HB2	1:B:91:LYS:HB2	2.00	0.43
1:A:617:ARG:HA	1:A:621:GLU:HB2	2.00	0.42
1:A:298:SER:HB3	1:B:463:ASN:HD22	1.85	0.41
1:A:304:LEU:HD23	1:A:309:LEU:HD13	2.02	0.41
1:A:515:VAL:HG22	1:A:568:LEU:HB3	2.03	0.41
1:B:186:CYS:HB3	1:B:231:ILE:HG12	2.03	0.40
1:A:65:ALA:HB3	1:A:181:ILE:HG12	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	566/611 (93%)	545 (96%)	20 (4%)	1 (0%)	47	79
1	B	538/611 (88%)	516 (96%)	22 (4%)	0	100	100
All	All	1104/1222 (90%)	1061 (96%)	42 (4%)	1 (0%)	51	83

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	329	MET

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	491/527 (93%)	481 (98%)	10 (2%)	55	80
1	B	475/527 (90%)	461 (97%)	14 (3%)	42	72
All	All	966/1054 (92%)	942 (98%)	24 (2%)	47	75

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	251	LEU
1	A	408	THR
1	A	507	LEU
1	A	531	ILE
1	A	567	LEU
1	A	586	SER
1	A	591	ASP
1	A	607	HIS
1	A	613	GLU
1	A	627	ASN
1	B	82	LEU
1	B	110	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	133	VAL
1	B	200	ASP
1	B	251	LEU
1	B	340	LEU
1	B	347	LEU
1	B	403	ILE
1	B	415	GLU
1	B	435	ILE
1	B	472	GLU
1	B	553	LEU
1	B	591	ASP
1	B	593	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	101	GLN
1	B	463	ASN
1	B	476	GLN
1	B	607	HIS
1	B	608	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	574/611 (93%)	0.46	31 (5%) 25 12	32, 61, 99, 128	0
1	B	552/611 (90%)	0.62	60 (10%) 5 2	38, 78, 112, 173	0
All	All	1126/1222 (92%)	0.54	91 (8%) 12 5	32, 69, 107, 173	0

All (91) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	39	GLU	5.3
1	A	50	GLN	4.7
1	B	341	CYS	4.4
1	B	357	LEU	4.3
1	A	37	ASP	4.0
1	B	344	ASP	3.9
1	A	376	GLY	3.9
1	A	530	VAL	3.9
1	A	48	HIS	3.7
1	B	252	GLY	3.5
1	B	525	PHE	3.5
1	B	297	SER	3.5
1	B	289	LEU	3.2
1	A	421	THR	3.2
1	B	268	CYS	3.2
1	B	437	GLU	3.2
1	A	532	ILE	3.1
1	B	557	LEU	3.1
1	A	536	GLY	3.1
1	B	255	PRO	3.0
1	A	110	THR	3.0
1	B	138	LEU	3.0
1	B	139	PHE	3.0
1	A	528	SER	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	535	ASP	2.9
1	A	33	GLY	2.9
1	B	290	GLN	2.9
1	B	337	LEU	2.7
1	B	411	MET	2.7
1	B	375	ASP	2.7
1	B	352	CYS	2.7
1	A	423	PRO	2.7
1	B	359	LYS	2.7
1	A	290	GLN	2.7
1	A	372	VAL	2.7
1	B	358	PHE	2.7
1	B	267	HIS	2.6
1	B	485	ALA	2.6
1	B	361	PHE	2.6
1	B	399	HIS	2.6
1	B	197	TRP	2.6
1	B	556	ASP	2.5
1	B	343	ASN	2.5
1	A	35	ASP	2.5
1	A	551	ASN	2.5
1	B	562	ALA	2.5
1	A	133	VAL	2.5
1	B	136	ARG	2.5
1	B	405	PHE	2.5
1	B	552	SER	2.5
1	B	414	ARG	2.4
1	B	573	ALA	2.4
1	A	537	PRO	2.4
1	B	501	LEU	2.4
1	A	425	ALA	2.4
1	B	439	THR	2.4
1	A	111	GLY	2.4
1	B	245	ASN	2.4
1	B	120	LEU	2.4
1	A	245	ASN	2.4
1	B	347	LEU	2.4
1	A	47	THR	2.3
1	A	538	VAL	2.3
1	B	394	THR	2.3
1	B	516	GLU	2.3
1	B	622	ALA	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	127	ASP	2.3
1	B	350	GLY	2.2
1	B	365	CYS	2.2
1	B	383	ASP	2.2
1	B	342	SER	2.2
1	B	400	ASN	2.2
1	A	531	ILE	2.2
1	B	345	GLU	2.2
1	B	565	VAL	2.2
1	A	590	PRO	2.1
1	B	43	ASN	2.1
1	B	57	THR	2.1
1	A	548	GLN	2.1
1	A	554	ASN	2.1
1	B	257	ARG	2.1
1	B	351	ILE	2.1
1	B	508	ARG	2.1
1	A	384	VAL	2.1
1	A	297	SER	2.1
1	B	279	GLY	2.1
1	B	441	ASN	2.1
1	B	551	ASN	2.0
1	B	97	PRO	2.0
1	B	447	SER	2.0
1	B	435	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.