



# Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 05:15 am BST

PDB ID : 5JEJ  
Title : Phosphorylated STING in complex with IRF-3 CTD  
Authors : Li, P.; Shu, C.  
Deposited on : 2016-04-18  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	39	
1	D	39	
1	E	39	
2	A	242	
2	B	242	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4338 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Stimulator of interferon genes protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	P				S
1	C	21	Total 175	C 109	N 27	O 37	P 1	S 1	0	0	0
1	D	10	Total 80	C 48	N 11	O 19	P 1	S 1	0	0	0
1	E	4	Total 36	C 22	N 5	O 9			0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	341	SER	-	expression tag	UNP Q86WV6
C	378	TRP	PHE	conflict	UNP Q86WV6
D	341	SER	-	expression tag	UNP Q86WV6
D	378	TRP	PHE	conflict	UNP Q86WV6
E	341	SER	-	expression tag	UNP Q86WV6
E	378	TRP	PHE	conflict	UNP Q86WV6

- Molecule 2 is a protein called Interferon regulatory factor 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	A	234	Total 1838	C 1173	N 317	O 338	S 10	0	0	0
2	B	233	Total 1829	C 1168	N 315	O 336	S 10	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	186	SER	-	expression tag	UNP Q14653
A	187	GLU	-	expression tag	UNP Q14653
A	188	PHE	-	expression tag	UNP Q14653
B	186	SER	-	expression tag	UNP Q14653

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	187	GLU	-	expression tag	UNP Q14653
B	188	PHE	-	expression tag	UNP Q14653



- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	C	11	Total O 11 11	0	0
3	D	3	Total O 3 3	0	0
3	E	2	Total O 2 2	0	0
3	A	190	Total O 190 190	0	0
3	B	174	Total O 174 174	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Stimulator of interferon genes protein





- Molecule 1: Stimulator of interferon genes protein




- Molecule 1: Stimulator of interferon genes protein



- Molecule 2: Interferon regulatory factor 3



- Molecule 2: Interferon regulatory factor 3



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	64.08Å 56.40Å 75.55Å 90.00° 104.64° 90.00°	Depositor
Resolution (Å)	44.65 – 2.00 44.65 – 1.99	Depositor EDS
% Data completeness (in resolution range)	98.0 (44.65-2.00) 88.9 (44.65-1.99)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.82 (at 2.00Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, $R_{free}$	0.176 , 0.237 0.180 , 0.238	Depositor DCC
$R_{free}$ test set	2000 reflections (5.70%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	21.8	Xtrriage
Anisotropy	0.893	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 57.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4338	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.64% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	C	0.31	0/168	0.57	0/225
1	D	0.28	0/69	0.54	0/90
1	E	0.33	0/37	0.56	0/49
2	A	0.38	0/1891	0.50	0/2577
2	B	0.37	0/1882	0.49	0/2565
All	All	0.37	0/4047	0.50	0/5506

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	175	0	169	4	1
1	D	80	0	74	0	0
1	E	36	0	25	2	0
2	A	1838	0	1786	14	0
2	B	1829	0	1778	12	0
3	A	190	0	0	3	1
3	B	174	0	0	2	1
3	C	11	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	3	0	0	0	0
3	E	2	0	0	0	0
All	All	4338	0	3832	30	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:233:VAL:HB	2:A:234:GLY:HA2	1.63	0.80
1:E:376:THR:HG21	2:B:256:GLY:HA3	1.80	0.63
2:A:262:ARG:NH2	3:A:503:HOH:O	2.32	0.61
1:C:377:ASP:OD1	1:C:379:SER:OG	2.22	0.56
2:A:213:ARG:NH1	3:A:502:HOH:O	2.30	0.54
2:A:345:TRP:CZ2	2:A:366:LYS:HD3	2.41	0.54
2:A:301:PRO:HB3	2:A:345:TRP:CD2	2.42	0.53
2:B:303:SER:OG	3:B:501:HOH:O	2.17	0.53
1:C:372:LEU:HA	1:C:375:ARG:HE	1.74	0.52
2:B:378:MET:SD	2:B:391:VAL:HG22	2.51	0.51
2:B:399:HIS:HB3	2:B:400:PRO:HD2	1.93	0.49
2:B:301:PRO:HB3	2:B:345:TRP:CD2	2.47	0.48
2:A:194:ARG:NH2	2:A:200:GLU:OE1	2.46	0.47
2:A:219:THR:HG23	3:A:647:HOH:O	2.14	0.46
2:B:357:PRO:HG2	2:B:360:LYS:HG3	1.98	0.46
2:B:378:MET:HB2	2:B:391:VAL:HG13	1.99	0.45
1:C:370:LYS:H	1:C:370:LYS:HG3	1.53	0.45
2:A:194:ARG:HA	2:A:197:VAL:HG23	1.99	0.45
1:E:376:THR:CG2	2:B:256:GLY:HA3	2.47	0.45
2:A:233:VAL:CB	2:A:234:GLY:HA2	2.42	0.43
2:A:281:LEU:HB2	2:A:320:PHE:HB3	1.99	0.43
2:B:272:LEU:HD21	2:B:346:PHE:CD2	2.55	0.42
2:A:254:ASP:OD1	2:A:254:ASP:N	2.52	0.42
2:B:345:TRP:CZ2	2:B:366:LYS:HD3	2.56	0.41
2:B:281:LEU:HB2	2:B:320:PHE:HB3	2.02	0.41
2:A:293:TRP:HZ3	2:A:295:VAL:HG23	1.85	0.41
1:C:372:LEU:HB3	1:C:373:PRO:HD3	2.02	0.41
2:A:235:ASP:OD1	2:A:236:ARG:N	2.54	0.40
2:A:390:THR:HA	2:A:401:LEU:O	2.22	0.40
2:B:380:ARG:HD3	3:B:533:HOH:O	2.20	0.40



All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:360:GLU:OE2	1:C:370:LYS:NZ[2_11516]	2.04	0.16
3:A:575:HOH:O	3:B:668:HOH:O[1_655]	2.14	0.06

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	18/39 (46%)	16 (89%)	1 (6%)	1 (6%)	2	0
1	D	7/39 (18%)	7 (100%)	0	0	100	100
1	E	2/39 (5%)	2 (100%)	0	0	100	100
2	A	232/242 (96%)	227 (98%)	4 (2%)	1 (0%)	34	30
2	B	231/242 (96%)	220 (95%)	11 (5%)	0	100	100
All	All	490/601 (82%)	472 (96%)	16 (3%)	2 (0%)	34	30

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	369	GLU
2	A	233	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	19/35 (54%)	17 (90%)	2 (10%)	7	4
1	D	8/35 (23%)	8 (100%)	0	100	100
1	E	4/35 (11%)	3 (75%)	1 (25%)	0	0
2	A	199/205 (97%)	194 (98%)	5 (2%)	47	49
2	B	198/205 (97%)	194 (98%)	4 (2%)	55	58
All	All	428/515 (83%)	416 (97%)	12 (3%)	43	44

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	360	GLU
1	C	374	LEU
1	E	376	THR
2	A	232	GLU
2	A	254	ASP
2	A	255	ARG
2	A	416	VAL
2	A	422	GLN
2	B	197	VAL
2	B	237	THR
2	B	255	ARG
2	B	391	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	SEP	D	366	1	8,9,10	1.51	1 (12%)	8,12,14	1.16	0
1	SEP	C	366	1	8,9,10	1.56	1 (12%)	8,12,14	1.71	2 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	D	366	1	-	0/5/8/10	-
1	SEP	C	366	1	-	0/5/8/10	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	366	SEP	P-O1P	3.34	1.61	1.50
1	D	366	SEP	P-O1P	3.23	1.61	1.50

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	366	SEP	OG-CB-CA	3.20	111.26	108.14
1	C	366	SEP	P-OG-CB	-3.09	109.78	118.30

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	C	20/39 (51%)	0.28	2 (10%) 7 6	37, 59, 88, 90	0
1	D	9/39 (23%)	1.00	2 (22%) 0 0	31, 43, 80, 89	0
1	E	4/39 (10%)	0.83	0 100 100	54, 61, 64, 76	0
2	A	234/242 (96%)	-0.22	9 (3%) 40 39	18, 30, 57, 99	0
2	B	233/242 (96%)	-0.23	4 (1%) 70 68	18, 30, 54, 104	0
All	All	500/601 (83%)	-0.18	17 (3%) 45 44	18, 31, 68, 104	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	234	GLY	9.8
2	A	234	GLY	8.1
2	A	233	VAL	6.9
2	B	235	ASP	6.8
2	B	233	VAL	4.6
1	D	368	MET	4.3
2	B	236	ARG	4.1
2	A	235	ASP	3.5
2	A	236	ARG	3.2
2	A	336	SER	3.1
2	A	232	GLU	3.1
1	C	372	LEU	2.9
2	A	255	ARG	2.8
2	A	305	HIS	2.7
1	D	367	GLY	2.4
2	A	422	GLN	2.2
1	C	369	GLU	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	SEP	D	366	10/11	0.96	0.10	42,53,58,59	0
1	SEP	C	366	10/11	0.97	0.09	39,47,71,71	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.