



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 9, 2020 – 06:02 AM BST

PDB ID : 6JIV
Title : SspE crystal structure
Authors : Bing, Y.Z.; Yang, H.G.
Deposited on : 2019-02-23
Resolution : 3.31 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric | Whole archive (#Entries) | Similar resolution (#Entries, resolution range(Å)) |
|-----------------------|-----------------------------|---|
| R_{free} | 130704 | 1089 (3.36-3.28) |
| Clashscore | 141614 | 1137 (3.36-3.28) |
| Ramachandran outliers | 138981 | 1115 (3.36-3.28) |
| Sidechain outliers | 138945 | 1114 (3.36-3.28) |
| RSRZ outliers | 127900 | 1059 (3.36-3.28) |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 1 | A | 771 | |
| 1 | B | 771 | |
| 1 | C | 771 | |
| 1 | D | 771 | |

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 15028 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SspE protein.

| Mol | Chain | Residues | Atoms | | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|----|---------|---------|-------|
| | | | Total | C | N | O | S | Se | | | |
| 1 | D | 560 | 3711 | 2310 | 674 | 713 | 3 | 11 | 0 | 0 | 0 |
| 1 | A | 560 | 3711 | 2310 | 674 | 713 | 3 | 11 | 0 | 0 | 0 |
| 1 | B | 560 | 3711 | 2310 | 674 | 713 | 3 | 11 | 0 | 0 | 0 |
| 1 | C | 557 | 3895 | 2439 | 697 | 743 | 4 | 12 | 0 | 0 | 0 |

4 Data and refinement statistics

| Property | Value | Source |
|---|---|------------------|
| Space group | P 21 21 21 | Depositor |
| Cell constants a, b, c, α , β , γ | 111.07Å 138.16Å 293.68Å 90.00° 90.00° 90.00° | Depositor |
| Resolution (Å) | 125.33 – 3.31 125.02 – 3.31 | Depositor EDS |
| % Data completeness (in resolution range) | 99.5 (125.33-3.31) 99.5 (125.02-3.31) | Depositor EDS |
| R_{merge} | (Not available) | Depositor |
| R_{sym} | (Not available) | Depositor |
| $\langle I/\sigma(I) \rangle$ ¹ | 1.77 (at 3.33Å) | Xtrriage |
| Refinement program | REFMAC 5.8.0258 | Depositor |
| R, R_{free} | 0.406 , 0.435 0.407 , 0.439 | Depositor DCC |
| R_{free} test set | 3357 reflections (4.96%) | wwPDB-VP |
| Wilson B-factor (Å ²) | 122.6 | Xtrriage |
| Anisotropy | 0.299 | Xtrriage |
| Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²) | 0.29 , 110.6 | EDS |
| L-test for twinning ² | $\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$ | Xtrriage |
| Estimated twinning fraction | No twinning to report. | Xtrriage |
| F_o, F_c correlation | 0.80 | EDS |
| Total number of atoms | 15028 | wwPDB-VP |
| Average B, all atoms (Å ²) | 109.0 | wwPDB-VP |

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|----------------|-------------|-----------------|
| | | RMSZ | # $ Z > 5$ | RMSZ | # $ Z > 5$ |
| 1 | A | 0.60 | 0/3747 | 0.84 | 17/5100 (0.3%) |
| 1 | B | 0.52 | 0/3747 | 0.75 | 16/5100 (0.3%) |
| 1 | C | 0.57 | 0/3943 | 0.82 | 15/5351 (0.3%) |
| 1 | D | 0.73 | 5/3747 (0.1%) | 0.90 | 16/5100 (0.3%) |
| All | All | 0.61 | 5/15184 (0.0%) | 0.83 | 64/20651 (0.3%) |

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

| Mol | Chain | #Chirality outliers | #Planarity outliers |
|-----|-------|---------------------|---------------------|
| 1 | A | 0 | 1 |
| 1 | C | 0 | 2 |
| All | All | 0 | 3 |

All (5) bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|--------|--------|-------------|----------|
| 1 | D | 146 | CYS | CB-SG | -10.19 | 1.65 | 1.82 |
| 1 | D | 110 | VAL | CB-CG1 | -5.86 | 1.40 | 1.52 |
| 1 | D | 133 | TYR | CE1-CZ | -5.46 | 1.31 | 1.38 |
| 1 | D | 260 | SER | CA-CB | -5.23 | 1.45 | 1.52 |
| 1 | D | 68 | VAL | CB-CG1 | -5.01 | 1.42 | 1.52 |

The worst 5 of 64 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|-----------|-------|-------------|----------|
| 1 | C | 484 | LEU | CB-CG-CD1 | -9.98 | 94.03 | 111.00 |
| 1 | D | 160 | PRO | N-CA-CB | 9.37 | 114.54 | 103.30 |
| 1 | A | 160 | PRO | N-CA-CB | 8.07 | 112.98 | 103.30 |
| 1 | A | 329 | PRO | N-CA-CB | 7.82 | 112.69 | 103.30 |
| 1 | B | 329 | PRO | N-CA-CB | 7.54 | 112.35 | 103.30 |

There are no chirality outliers.

All (3) planarity outliers are listed below:

| Mol | Chain | Res | Type | Group |
|-----|-------|-----|------|---------|
| 1 | A | 127 | GLU | Peptide |
| 1 | C | 477 | ARG | Peptide |
| 1 | C | 584 | VAL | Peptide |

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | A | 3711 | 0 | 3036 | 217 | 0 |
| 1 | B | 3711 | 0 | 3038 | 219 | 0 |
| 1 | C | 3895 | 0 | 3376 | 323 | 0 |
| 1 | D | 3711 | 0 | 3038 | 246 | 0 |
| All | All | 15028 | 0 | 12488 | 970 | 0 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 35.

The worst 5 of 970 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|----------------|------------------|--------------------------|-------------------|
| 1:C:515:PHE:CZ | 1:C:578:ARG:HD3 | 1.32 | 1.63 |
| 1:D:356:LEU:CA | 1:D:427:LEU:HD11 | 1.17 | 1.59 |
| 1:A:356:LEU:CA | 1:A:427:LEU:HD11 | 1.16 | 1.59 |
| 1:D:356:LEU:HA | 1:D:427:LEU:CD1 | 1.30 | 1.55 |
| 1:D:395:LYS:CE | 1:D:526:ARG:HD3 | 1.37 | 1.55 |

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|-----------------|------------|-----------|----------|-------------|---|
| 1 | A | 552/771 (72%) | 429 (78%) | 73 (13%) | 50 (9%) | 1 | 5 |
| 1 | B | 552/771 (72%) | 437 (79%) | 70 (13%) | 45 (8%) | 1 | 6 |
| 1 | C | 549/771 (71%) | 423 (77%) | 80 (15%) | 46 (8%) | 1 | 6 |
| 1 | D | 552/771 (72%) | 427 (77%) | 75 (14%) | 50 (9%) | 1 | 5 |
| All | All | 2205/3084 (72%) | 1716 (78%) | 298 (14%) | 191 (9%) | 1 | 5 |

5 of 191 Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | D | 2 | GLU |
| 1 | D | 4 | LYS |
| 1 | D | 6 | ILE |
| 1 | D | 61 | SER |
| 1 | D | 129 | SER |

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|-----------------|------------|-----------|-------------|----|
| 1 | A | 268/637 (42%) | 241 (90%) | 27 (10%) | 7 | 28 |
| 1 | B | 268/637 (42%) | 245 (91%) | 23 (9%) | 10 | 36 |
| 1 | C | 322/637 (50%) | 293 (91%) | 29 (9%) | 9 | 33 |
| 1 | D | 268/637 (42%) | 240 (90%) | 28 (10%) | 7 | 26 |
| All | All | 1126/2548 (44%) | 1019 (90%) | 107 (10%) | 8 | 30 |

5 of 107 residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | A | 303 | VAL |
| 1 | B | 67 | THR |
| 1 | C | 515 | PHE |
| 1 | A | 358 | PHE |
| 1 | A | 525 | TYR |

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | A | 391 | GLN |
| 1 | B | 156 | HIS |
| 1 | B | 429 | GLN |
| 1 | A | 156 | HIS |
| 1 | B | 391 | GLN |

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

| Mol | Chain | Analysed | <RSRZ> | #RSRZ>2 | OWAB(Å ²) | Q<0.9 |
|-----|-------|-----------------|--------|---------------|-----------------------|-------|
| 1 | A | 548/771 (71%) | 0.30 | 41 (7%) 14 14 | 53, 109, 176, 219 | 0 |
| 1 | B | 548/771 (71%) | 0.87 | 92 (16%) 1 1 | 62, 120, 274, 329 | 0 |
| 1 | C | 545/771 (70%) | 0.37 | 28 (5%) 28 27 | 30, 104, 136, 171 | 0 |
| 1 | D | 548/771 (71%) | 0.36 | 36 (6%) 18 19 | 37, 91, 169, 223 | 0 |
| All | All | 2189/3084 (70%) | 0.47 | 197 (8%) 9 10 | 30, 106, 216, 329 | 0 |

The worst 5 of 197 RSRZ outliers are listed below:

| Mol | Chain | Res | Type | RSRZ |
|-----|-------|-----|------|------|
| 1 | B | 446 | SER | 14.8 |
| 1 | B | 448 | ASP | 12.9 |
| 1 | B | 533 | VAL | 10.3 |
| 1 | B | 522 | ASP | 9.0 |
| 1 | B | 532 | VAL | 8.3 |

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.