

Full wwPDB X-ray Structure Validation Report (i)

Aug 28, 2023 - 04:52 AM EDT

hy-
ן

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolDrobity		4.02b-467
5		
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 1.93 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\# \textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
Clashscore	141614	1023 (1.94-1.94)
Ramachandran outliers	138981	1007 (1.94-1.94)
Sidechain outliers	138945	1007 (1.94-1.94)
RSRZ outliers	127900	4250 (1.96-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	122	76%	17%	• 6%
1	В	122	80%	16%	·
1	С	122	7%82%	14%	•••



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 2915 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	А	115	Total	С	Ν	Ο	OS 0 2	9	Ο	
	Л	110	883	563	152	162	6	0	2	0
1	В	117	Total	С	Ν	Ο	\mathbf{S}	0	9	0
	D	117	901	573	156	166	6	0	2	0
1	С	118	Total	С	Ν	Ο	S	0	ე	0
	U	110	912	580	159	167	6	U		U

• Molecule 1 is a protein called Macrophage migration inhibitory factor.

Chain	Residue	Modelled	Actual	Comment	Reference
А	115	LEU	-	expression tag	UNP P14174
А	116	GLU	-	expression tag	UNP P14174
А	117	HIS	-	expression tag	UNP P14174
А	118	HIS	-	expression tag	UNP P14174
А	119	HIS	-	expression tag	UNP P14174
А	120	HIS	-	expression tag	UNP P14174
А	121	HIS	-	expression tag	UNP P14174
А	122	HIS	-	expression tag	UNP P14174
В	115	LEU	-	expression tag	UNP P14174
В	116	GLU	-	expression tag	UNP P14174
В	117	HIS	-	expression tag	UNP P14174
В	118	HIS	-	expression tag	UNP P14174
В	119	HIS	-	expression tag	UNP P14174
В	120	HIS	-	expression tag	UNP P14174
В	121	HIS	-	expression tag	UNP P14174
В	122	HIS	-	expression tag	UNP P14174
С	115	LEU	-	expression tag	UNP P14174
С	116	GLU	-	expression tag	UNP P14174
С	117	HIS	-	expression tag	UNP P14174
С	118	HIS	-	expression tag	UNP P14174
С	119	HIS	-	expression tag	UNP P14174
С	120	HIS	-	expression tag	UNP P14174
С	121	HIS	-	expression tag	UNP P14174
				Continued	on nort nado


There are 24 discrepancies between the modelled and reference sequences:



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
С	122	HIS	-	expression tag	UNP P14174


• Molecule 2 is 7-(2-fluorobenzyl)quinolin-8-ol (three-letter code: XV1) (formula: $C_{16}H_{12}FNO$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	Δ	1	Total	С	F	Ν	Ο	0	0
	Л	I	19	16	1	1	1	0	0
2	В	1	Total	С	F	Ν	Ο	0	0
	D	1	19	16	1	1	1	0	0
2	С	1	Total	С	F	Ν	Ο	0	0
	U	1	19	16	1	1	1	0	

• Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O_4S).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	В	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
3	В	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0


• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	45	Total O 45 45	0	0
4	В	65	Total O 65 65	0	0
4	С	42	$\begin{array}{cc} \text{Total} & \text{O} \\ 42 & 42 \end{array}$	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Macrophage migration inhibitory factor



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	67.65\AA 67.62\AA 88.58\AA	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	23.91 - 1.93	Depositor
Resolution (A)	23.91 - 1.93	EDS
% Data completeness	$95.7\ (23.91-1.93)$	Depositor
(in resolution range)	$95.9\ (23.91\text{-}1.93)$	EDS
R _{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.24 (at 1.93 \text{\AA})$	Xtriage
Refinement program	CNS, CNX 2005	Depositor
R, R_{free}	0.187 , 0.218	Depositor
II, II, ree	0.179 , (Not available)	DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor $(Å^2)$	27.3	Xtriage
Anisotropy	0.203	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.41 , 46.9	EDS
L-test for twinning ²	$< L > = 0.48, < L^2 > = 0.31$	Xtriage
Estimated twinning fraction	0.024 for k,h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2915	wwPDB-VP
Average B, all atoms $(Å^2)$	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 6.27% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, $\rm XV1$

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	lengths	Bond angles		
	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.27	0/909	0.49	0/1237	
1	В	0.27	0/928	0.48	0/1263	
1	С	0.25	0/940	0.46	0/1279	
All	All	0.26	0/2777	0.47	0/3779	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	883	0	887	27	0
1	В	901	0	899	21	0
1	С	912	0	908	20	0
2	А	19	0	10	2	0
2	В	19	0	10	1	0
2	С	19	0	10	1	0
3	В	10	0	0	1	0
4	А	45	0	0	0	0
4	В	65	0	0	0	0
4	С	42	0	0	0	0
All	All	2915	0	2724	64	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:33:PRO:HG2	1:B:36:TYR:CD2	2.22	0.74
1:B:4:ILE:HD12	1:B:40:HIS:HB3	1.70	0.72
1:A:3:PHE:CE2	1:A:5[B]:VAL:HG13	2.28	0.68
1:C:5[B]:VAL:CG2	1:C:41:VAL:HG22	2.25	0.65
1:B:3:PHE:CE2	1:B:5[B]:VAL:HG13	2.32	0.64
1:B:19:LEU:HD22	1:B:39[B]:VAL:HG23	1.79	0.64
1:A:2:MET:HE3	1:C:58:LEU:HD13	1.79	0.64
1:C:33:PRO:HG2	1:C:36:TYR:CE2	2.32	0.64
1:C:27:ALA:HA	1:C:37[B]:ILE:HD11	1.80	0.64
1:B:33:PRO:HG2	1:B:36:TYR:CE2	2.34	0.63
1:A:1:PRO:HG3	1:A:32:LYS:HG3	1.79	0.62
1:B:11:ARG:HD3	3:B:124:SO4:O1	1.99	0.60
1:B:33:PRO:HG2	1:B:36:TYR:HD2	1.66	0.59
1:C:4:ILE:CD1	1:C:40:HIS:HB3	2.32	0.59
1:C:33:PRO:HG2	1:C:36:TYR:CD2	2.37	0.59
1:C:5[B]:VAL:HG22	1:C:41:VAL:HG22	1.86	0.58
1:C:4:ILE:HD12	1:C:40:HIS:HB3	1.87	0.55
1:A:5[B]:VAL:CG2	1:A:41:VAL:HG22	2.37	0.55
1:B:5[B]:VAL:CG2	1:B:41:VAL:HG22	2.37	0.55
1:B:95:TYR:CD2	2:C:123:XV1:HAN	2.42	0.54
1:B:5[B]:VAL:HG23	1:B:41:VAL:HA	1.90	0.54
1:C:102:ASN:HD22	1:C:104:ALA:H	1.54	0.54
1:C:27:ALA:CA	1:C:37[B]:ILE:HD11	2.39	0.53
1:C:102:ASN:ND2	1:C:104:ALA:H	2.07	0.52
1:A:5[B]:VAL:HG23	1:A:41:VAL:HA	1.91	0.51
1:B:3:PHE:CE2	1:B:5[A]:VAL:HG22	2.45	0.51
1:A:3:PHE:CE2	1:A:5[A]:VAL:CG2	2.94	0.51
1:A:27:ALA:HA	1:A:37[B]:ILE:HD11	1.92	0.50
1:B:4:ILE:CD1	1:B:40:HIS:HB3	2.41	0.49
1:A:3:PHE:CE2	1:A:5[A]:VAL:HG22	2.47	0.49
1:A:27:ALA:CA	1:A:37[B]:ILE:HD11	2.43	0.48
1:B:5[B]:VAL:HG12	1:B:59:CYS:SG	2.53	0.48
1:A:3:PHE:CE2	1:A:5[B]:VAL:CG1	2.95	0.48
1:A:4:ILE:HD12	1:A:40:HIS:HB3	1.95	0.48
1:B:3:PHE:CE2	1:B:5[A]:VAL:CG2	2.97	0.47
1:A:1:PRO:HG2	1:A:37[A]:ILE:CD1	2.44	0.47
1:A:105:ASN:HD22	1:C:72:ASN:ND2	2.13	0.47



		Interatomic	Clash
Atom-1	Atom-2	distance (\AA)	overlap (Å)
1:A:95:TYR:CD2	2:B:123:XV1:HAN	2.50	0.46
1:A:1:PRO:HD2	1:A:37[A]:ILE:HD13	1.98	0.46
1:A:1:PRO:CG	1:A:32:LYS:HG3	2.45	0.46
1:A:109:ASN:O	1:A:110:ASN:HB2	2.16	0.46
1:A:33:PRO:HA	1:A:34:PRO:HD3	1.88	0.45
1:C:5[B]:VAL:HG21	1:C:41:VAL:HG22	1.98	0.45
1:A:115:LEU:HD13	1:C:73:ARG:HD3	1.99	0.45
1:B:109:ASN:O	1:B:110:ASN:HB2	2.17	0.45
1:C:25:GLN:HE21	1:C:78:LEU:HD11	1.81	0.45
1:A:5[B]:VAL:HG12	1:A:59:CYS:SG	2.57	0.44
1:A:72:ASN:ND2	1:B:105:ASN:HD22	2.15	0.44
1:B:33:PRO:HA	1:B:34:PRO:HD3	1.84	0.44
1:C:5[B]:VAL:CG2	1:C:41:VAL:HA	2.47	0.44
1:A:1:PRO:N	2:A:123:XV1:OAK	2.51	0.44
2:A:123:XV1:HAH	1:C:95:TYR:CE2	2.53	0.44
1:C:4:ILE:HD13	1:C:40:HIS:HB3	2.01	0.42
1:B:19:LEU:HD22	1:B:39[B]:VAL:CG2	2.49	0.42
1:A:4:ILE:CD1	1:A:40:HIS:HB3	2.49	0.42
1:C:109:ASN:O	1:C:110:ASN:HB2	2.19	0.42
1:A:10:PRO:HA	1:A:44:ASP:OD1	2.19	0.42
1:A:5[B]:VAL:CG2	1:A:41:VAL:HA	2.50	0.41
1:B:1:PRO:HA	1:B:62:HIS:O	2.19	0.41
1:A:1:PRO:HA	1:A:62:HIS:O	2.21	0.41
1:B:1:PRO:CG	1:B:32:LYS:HD2	2.50	0.41
1:B:1:PRO:HG3	1:B:32:LYS:HD2	2.02	0.40

Continued from previous page...

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured Allowed		Outliers	Percentiles	
1	А	115/122~(94%)	113~(98%)	2(2%)	0	100 100	



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	В	117/122~(96%)	114 (97%)	3~(3%)	0	100	100
1	С	118/122~(97%)	117 (99%)	1 (1%)	0	100	100
All	All	350/366~(96%)	344 (98%)	6(2%)	0	100	100

Continued from previous page...

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	А	97/102~(95%)	96~(99%)	1 (1%)	76 71		
1	В	99/102~(97%)	99 (100%)	0	100 100		
1	С	100/102~(98%)	99~(99%)	1 (1%)	76 71		
All	All	296/306~(97%)	294 (99%)	2(1%)	84 81		

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	109	ASN
1	С	109	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	45	GLN
1	А	105	ASN
1	А	109	ASN
1	В	25	GLN
1	В	105	ASN
1	В	109	ASN
1	С	25	GLN
1	С	102	ASN
1	С	105	ASN



Continued from previous page...

Mol	Chain	Res	Type
1	С	109	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Type Chain		Dec	Tinle	Bond lengths			Bond angles		
	Mol Type Chain	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	XV1	А	123	1	21,21,21	0.97	0	29,29,29	1.32	3 (10%)
3	SO4	В	125	-	4,4,4	0.25	0	6,6,6	0.29	0
2	XV1	В	123	1	21,21,21	0.97	0	29,29,29	1.27	3 (10%)
3	SO4	В	124	-	4,4,4	0.21	0	6,6,6	0.33	0
2	XV1	С	123	1	21,21,21	0.96	0	29,29,29	1.28	<mark>3 (10%)</mark>

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	XV1	В	123	1	-	0/4/4/4	0/3/3/3
2	XV1	А	123	1	-	0/4/4/4	0/3/3/3
2	XV1	С	123	1	-	0/4/4/4	0/3/3/3

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
2	А	123	XV1	CAM-NAG-CAE	4.02	122.30	117.30
2	В	123	XV1	CAM-NAG-CAE	3.94	122.20	117.30
2	С	123	XV1	CAM-NAG-CAE	3.83	122.07	117.30
2	А	123	XV1	CAI-CAE-NAG	-2.54	119.84	122.62
2	В	123	XV1	CAI-CAE-NAG	-2.42	119.97	122.62
2	С	123	XV1	CAI-CAE-NAG	-2.32	120.08	122.62
2	А	123	XV1	CAQ-CAM-NAG	-2.22	120.54	123.94
2	В	123	XV1	CAQ-CAM-NAG	-2.18	120.59	123.94
2	С	123	XV1	CAQ-CAM-NAG	-2.18	120.60	123.94

There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	А	123	XV1	2	0
2	В	123	XV1	1	0
3	В	124	SO4	1	0
2	С	123	XV1	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	115/122~(94%)	0.04	3 (2%) 56 63	20, 26, 38, 55	0
1	В	117/122~(95%)	-0.12	2 (1%) 70 75	20, 25, 34, 41	0
1	С	118/122~(96%)	0.23	8 (6%) 17 23	21, 29, 42, 53	0
All	All	350/366~(95%)	0.05	13 (3%) 41 49	20, 26, 38, 55	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	С	118	HIS	3.8
1	А	115	LEU	3.6
1	С	12	ALA	3.5
1	С	88	ARG	2.8
1	С	28	GLN	2.8
1	С	73	ARG	2.7
1	А	88	ARG	2.6
1	В	88	ARG	2.6
1	С	24	GLN	2.3
1	С	16	ASP	2.3
1	В	117	HIS	2.1
1	С	11	ARG	2.1
1	А	35	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q<0.9
2	XV1	С	123	19/19	0.89	0.14	$34,\!36,\!38,\!38$	0
2	XV1	В	123	19/19	0.91	0.11	29,31,34,35	0
2	XV1	А	123	19/19	0.92	0.11	35,37,39,40	0
3	SO4	В	125	5/5	0.93	0.33	$67,\!67,\!68,\!68$	0
3	SO4	В	124	5/5	0.94	0.24	61,63,64,64	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.









6.5 Other polymers (i)

There are no such residues in this entry.

