

Full wwPDB NMR Structure Validation Report (i)

May 29, 2020 – 08:19 am BST

PDB ID : 5JTM

Title : The structure of chaperone SecB in complex with unstructured PhoA binding

site a

Authors: Huang, C.; Saio, T.; Rossi, P.; Kalodimos, C.G.

Deposited on : 2016-05-09

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/NMRValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)

NmrClust : Kelley et al. (1996)

MolProbity: 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

RCI : v 1n 11 5 13 A (Berjanski et al., 2005)

PANAV : Wang et al. (2010)

ShiftChecker : 2.11

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment is 13%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric | Whole archive | NMR archive | | |
|-----------------------|-------------------------|-------------------------|--|--|
| Metric | $(\# \mathrm{Entries})$ | $(\# \mathrm{Entries})$ | | |
| Clashscore | 158937 | 12864 | | |
| Ramachandran outliers | 154571 | 11451 | | |
| Sidechain outliers | 154315 | 11428 | | |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

| Mol | Chain | Length | Quality of chain | | |
|-----|-------|--------|------------------|-----|-----|
| 1 | A | 155 | 71% | • | 26% |
| 1 | В | 155 | 77% | 5% | 19% |
| 1 | С | 155 | 75% | ••• | 20% |
| 1 | D | 155 | 77% | · | 21% |
| 2 | Е | 25 | 100% | | |
| 2 | F | 25 | 100% | | |
| 2 | G | 25 | 100% | | |
| 2 | Н | 25 | 100% | | |



2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 20 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: lowest energy.

The following residues are included in the computation of the global validation metrics.

| Well-defined (core) protein residues | | | | | | | |
|---|------------------------|------|----|--|--|--|--|
| Well-defined core Residue range (total) Backbone RMSD (Å) Medoid mode | | | | | | | |
| 1 | A:11-A:86, A:95-A:133, | 0.44 | 20 | | | | |
| | B:9-B:134, C:11-C:134, | | | | | | |
| | D:10-D:131 (487) | | | | | | |

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 2 single-model clusters were found.

| Cluster number | Models |
|-----------------------|---|
| 1 | 1, 2, 5, 8, 9, 10, 11, 14, 16, 17, 18, 20 |
| 2 | 6, 13 |
| 3 | 7, 12 |
| 4 | 3, 4 |
| Single-model clusters | 15; 19 |



3 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 11104 atoms, of which 5488 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called Protein-export protein SecB.

| Mol | Chain | Residues | | Atoms | | | | | Trace |
|-----|-------|----------|-------|-------|------|-----|-----|---|-------|
| 1 | A | 155 | Total | С | Н | N | О | S | 0 |
| 1 | A | 155 | 2367 | 762 | 1155 | 198 | 243 | 9 | |
| 1 | В | 155 | Total | С | Н | N | О | S | 0 |
| 1 | D | 155 | 2367 | 762 | 1155 | 198 | 243 | 9 | 0 |
| 1 | С | 155 | Total | С | Н | N | О | S | 0 |
| 1 | | 155 | 2367 | 762 | 1155 | 198 | 243 | 9 | 0 |
| 1 | D | 155 | Total | С | Н | N | О | S | 0 |
| 1 | ש | 199 | 2367 | 762 | 1155 | 198 | 243 | 9 | 0 |

• Molecule 2 is a protein called Alkaline phosphatase.

| Mol | Chain | Residues | | Atoms | | | | | Trace |
|-----|-------|----------|-------|-------|-----|----|----|---|-------|
| 2 | Е | 25 | Total | С | Н | N | О | S | 0 |
| 2 | 15 | 29 | 409 | 126 | 217 | 31 | 34 | 1 | |
| 2 | F | 25 | Total | С | Н | N | О | S | 0 |
| | 1' | 20 | 409 | 126 | 217 | 31 | 34 | 1 | |
| 2 | G | 25 | Total | С | Н | N | О | S | 0 |
| | G | 20 | 409 | 126 | 217 | 31 | 34 | 1 | |
| 2 | Н | 25 | Total | С | Н | N | О | S | 0 |
| | 11 | 20 | 409 | 126 | 217 | 31 | 34 | 1 | U |




4 Residue-property plots (i)


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: Protein-export protein SecB











21%

E147 G148 T149 E150 E151 H152 Q153 D154

Chain D:

• Molecule 1: Protein-export protein SecB

• Molecule 2: Alkaline phosphatase

Chain E:

• Molecule 2: Alkaline phosphatase

Chain F:

• Molecule 2: Alkaline phosphatase

Chain G:


• Molecule 2: Alkaline phosphatase

Chain H: 100%

4.2.2 Score per residue for model 2


• Molecule 1: Protein-export protein SecB

Chain A: 65% 9% 26%



T149 E150 E151 H152 Q153 D154









4.2.3 Score per residue for model 3


• Molecule 1: Protein-export protein SecB




• Molecule 1: Protein-export protein SecB



• Molecule 1: Protein-export protein SecB



• Molecule 1: Protein-export protein SecB



• Molecule 2: Alkaline phosphatase

Chain E: 100%

• Molecule 2: Alkaline phosphatase

Chain F: 100%



• Molecule 2: Alkaline phosphatase


Chain G:

• Molecule 2: Alkaline phosphatase

Chain H:


4.2.4 Score per residue for model 4

• Molecule 1: Protein-export protein SecB




G148 T149 E150 E151 H152 Q153 D154

• Molecule 1: Protein-export protein SecB



• Molecule 1: Protein-export protein SecB



• Molecule 1: Protein-export protein SecB


Chain D: 73% 5% 21%






| Chain E: | 100% | | | |
|--|--|--|--|----------------------|
| M S S S S S S S S S S S S S S S S S S S | 114 116 116 117 119 120 120 120 123 123 123 | | | |
| • Molecule 2: All | kaline phosphatase | | | |
| Chain F: | 100% | | | |
| M1 K2 K2 C43 C43 C54 C16 C16 C10 C10 C11 C11 C11 C11 C11 C11 C11 C11 | 116 116 116 116 119 119 119 123 123 123 123 | | | |
| • Molecule 2: All | kaline phosphatase | | | |
| Chain G: | 100% | | | |
| M2 C3 | 114 116 117 119 119 119 119 119 119 119 119 119 | | | |
| • Molecule 2: All | kaline phosphatase | | | |
| Chain H: | 100% | | | |
| M1 K2 K2 K2 K2 K4 | 114 116 119 119 119 123 1733 1733 1733 1733 | | | |
| _ | er residue for model 5 otein-export protein SecB | | | |
| Chain A: | 63% | 11% • | 26% | |
| 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 | 74 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 | A73 Q77 Q77 I86 A87 G88 G88 G99 G91 G91 | 093 M94 1106 D134 A136 L136 | F137 M138 N139 |
| 1141 0142 0143 0144 0144 0146 0146 0148 0148 0148 0148 0148 | A165 | | | |
| • Molecule 1: Pro | otein-export protein SecB | | | |
| Chain B: | 77% | 5% | 19% | |
| M | 744 744 7445 7446 7446 7446 7446 7446 74 | E147 (148 T149 E150 E151 (1152 (1153 D154 A155 | | |
| • Molecule 1: Pre | otein-export protein SecB | | | |
| Chain C: | 67% | 12% • | 20% | |
| | | | | |











 \bullet Molecule 2: Alkaline phosphatase




























4.2.11 Score per residue for model 11









• Molecule 2: Alkaline phosphatase

Chain E:












• Molecule 1: Protein-export protein SecB


Chain B: 72% 9% 19%






4.2.15 Score per residue for model 15












| • Molecule 2: Alkaline phospha | atase | | | |
|---|--|--|--|------------------------------|
| Chain F: | 100% | | | |
| 新 16 16 16 16 16 17 17 17 17 18 17 18 17 18 17 18 17 18 18 18 18 18 18 18 18 18 18 18 18 18 | 2 | | | |
| • Molecule 2: Alkaline phospha | atase | | | |
| Chain G: | 100% | | | |
| M 72 | 25 4 4 4 5 5 5 4 5 5 5 5 5 5 5 5 5 5 5 5 | | | |
| • Molecule 2: Alkaline phospha | atase | | | |
| Chain H: | 100% | | | |
| 所 23 24 24 25 26 27 27 27 27 27 27 27 27 27 27 | 2 | | | |
| 4.2.17 Score per residue f | or model 17 | | | |
| - | | | | |
| • Molecule 1: Protein-export p | orotein SecB | | | |
| Chain A: | 5% 9% | 269 | % | |
| M1 | V64 T65 T73 473 186 186 188 189 192 192 192 193 194 | L98 M117 D134 A135 L136 | F137 M138 M139 Y140 L141 Q142 Q143 | Q144 A145 G146 E147 |
| 0148 2150 2150 2151 2151 2153 20153 20154 4155 | | | | |
| • Molecule 1: Protein-export p | orotein SecB | | | |
| Chain B: | 72% | 8% • | 19% | |
| M | E112 1114 1114 1116 1118 1119 1110 1100 | 1141 Q142 Q143 Q144 A145 G146 G148 | T149 E150 E151 H152 Q153 D154 | |
| • Molecule 1: Protein-export p | orotein SecB | | | |
| Chain C: | 70% | 10% | 20% | |
| 표 | M17 M129 M135 M136 M136 M136 M136 M136 M136 M136 M136 | F137 M138 M139 V140 Q142 Q143 Q144 | A145 G146 E147 G148 T149 E150 | H152 Q153 D154 A155 |
| • Molecule 1: Protein-export p | orotein SecB | | | |
| Chain D: | 74% | 5% | 21% | |
| | | | | |



100%

Chain E:



• Molecule 2: Alkaline phosphatase

Chain F: 100%

• Molecule 2: Alkaline phosphatase

Chain G: 100%

• Molecule 2: Alkaline phosphatase

Chain H: 100%

Score per residue for model 18 4.2.18


• Molecule 1: Protein-export protein SecB



• Molecule 1: Protein-export protein SecB


Chain B: 74% 19%







E150 E151 H152 Q153 D154





4.2.20 Score per residue for model 20 (medoid)

• Molecule 1: Protein-export protein SecB

E151 H152 Q153 D154 A155

• Molecule 1: Protein-export protein SecB

Chain B: 75% 6% 19%

• Molecule 1: Protein-export protein SecB

Chain C: 70% 10% • 20%

A155

• Molecule 1: Protein-export protein SecB

Chain D: 72% 6% 21%

• Molecule 2: Alkaline phosphatase

Chain E:

• Molecule 2: Alkaline phosphatase

Chain F:



| α_1 , α | |
|-----------------------|------|
| Chain G: | 100% |
| Chan Ci. | 100% |

• Molecule 2: Alkaline phosphatase

Chain H: 100%



5 Refinement protocol and experimental data overview (i)



The models were refined using the following method: molecular dynamics.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: target function.

The following table shows the software used for structure solution, optimisation and refinement.

| Software name | Classification | Version |
|---------------|-----------------------|---------|
| CNS | refinement | |
| CYANA | structure calculation | |

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 6 of this report.

| Chemical shift file(s) | $input_cs.cif$ |
|--|-----------------|
| Number of chemical shift lists | 8 |
| Total number of shifts | 5090 |
| Number of shifts mapped to atoms | 5090 |
| Number of unparsed shifts | 0 |
| Number of shifts with mapping errors | 0 |
| Number of shifts with mapping warnings | 0 |
| Assignment completeness (well-defined parts) | 13% |

No validations of the models with respect to experimental NMR restraints is performed at this time.

COVALENT-GEOMETRY INFOmissingINFO

5.1Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes |
|-----|-------|-------|----------|----------|---------|
| 1 | A | 904 | 883 | 881 | 5±1 |
| 1 | В | 981 | 955 | 953 | 6±2 |
| 1 | С | 966 | 939 | 937 | 7±2 |
| 1 | D | 946 | 927 | 925 | 5±2 |
| 2 | Е | 0 | 0 | 0 | 0±0 |
| 2 | F | 0 | 0 | 0 | 0±0 |
| 2 | G | 0 | 0 | 0 | 0±0 |



Continued from previous page...

| Mol | Chain | Non-H | H(model) | $\mathbf{H}(\mathbf{added})$ | Clashes |
|-----|-------|-------|----------|------------------------------|---------|
| 2 | Н | 0 | 0 | 0 | 0±0 |
| All | All | 75940 | 74080 | 73920 | 407 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All unique clashes are listed below, sorted by their clash magnitude.

| A., 1 | A 0 | $oxed{ 	ext{Clash(\AA)} \ 	ext{Distance(Å)} }$ | D: 4 (8) | Models | |
|-----------------|------------------|--|-------------|--------|-------|
| Atom-1 | Atom-2 | | Distance(A) | Worst | Total |
| 1:B:25:ALA:HB1 | 1:B:28:ALA:HB2 | 0.80 | 1.53 | 14 | 7 |
| 1:B:13:ILE:HD11 | 1:B:105:ILE:HD13 | 0.68 | 1.66 | 16 | 1 |
| 1:A:31:VAL:HG21 | 1:A:73:ALA:HA | 0.64 | 1.67 | 5 | 16 |
| 1:B:25:ALA:CB | 1:B:28:ALA:HB2 | 0.61 | 2.24 | 14 | 6 |
| 1:D:20:ASP:HB3 | 1:D:79:GLN:HB2 | 0.60 | 1.73 | 19 | 6 |
| 1:C:34:LYS:HG2 | 1:C:68:LEU:HD13 | 0.59 | 1.73 | 20 | 10 |
| 1:C:31:VAL:HG21 | 1:C:73:ALA:HA | 0.58 | 1.75 | 17 | 20 |
| 1:C:11:PHE:HE1 | 1:C:102:CYS:HG | 0.58 | 1.42 | 18 | 1 |
| 1:B:98:LEU:HA | 1:B:102:CYS:SG | 0.58 | 2.39 | 3 | 4 |
| 1:A:44:LEU:HD23 | 1:A:60:LEU:HD21 | 0.58 | 1.74 | 12 | 1 |
| 1:B:42:LEU:HD22 | 1:B:128:LEU:HD11 | 0.57 | 1.75 | 3 | 1 |
| 1:B:60:LEU:HD22 | 1:B:103:PRO:HB3 | 0.57 | 1.75 | 14 | 2 |
| 1:D:43:ASP:HB2 | 1:D:63:THR:HB | 0.57 | 1.75 | 17 | 5 |
| 1:C:64:VAL:HG23 | 1:C:114:ILE:HD13 | 0.57 | 1.75 | 13 | 3 |
| 1:B:21:ILE:HD11 | 1:B:113:CYS:SG | 0.57 | 2.40 | 14 | 6 |
| 1:B:41:LYS:HB2 | 1:B:65:THR:HB | 0.56 | 1.76 | 10 | 1 |
| 1:A:43:ASP:HB2 | 1:A:63:THR:HB | 0.56 | 1.77 | 3 | 4 |
| 1:D:86:ILE:HG12 | 1:D:89:ILE:HD13 | 0.55 | 1.77 | 5 | 2 |
| 1:C:45:ASP:HB3 | 1:C:61:ARG:HD2 | 0.55 | 1.76 | 19 | 2 |
| 1:A:86:ILE:HD11 | 1:A:102:CYS:SG | 0.54 | 2.42 | 15 | 1 |
| 1:A:64:VAL:HG11 | 1:A:126:LEU:HD21 | 0.54 | 1.77 | 13 | 3 |
| 1:B:9:MET:SD | 1:B:10:THR:N | 0.54 | 2.80 | 15 | 4 |
| 1:D:40:VAL:HG13 | 1:D:64:VAL:HG13 | 0.54 | 1.79 | 4 | 5 |
| 1:C:41:LYS:HB2 | 1:C:65:THR:HB | 0.54 | 1.80 | 19 | 2 |
| 1:B:43:ASP:HB2 | 1:B:63:THR:HB | 0.54 | 1.79 | 7 | 2 |
| 1:C:117:MET:HE2 | 1:D:21:ILE:HG21 | 0.53 | 1.80 | 1 | 10 |
| 1:D:47:ALA:HB3 | 1:D:59:VAL:HB | 0.53 | 1.80 | 8 | 7 |
| 1:B:11:PHE:CZ | 1:B:13:ILE:HG13 | 0.53 | 2.38 | 16 | 1 |
| 1:A:23:PHE:HB2 | 1:A:117:MET:SD | 0.53 | 2.43 | 2 | 5 |
| 1:D:90:GLU:HA | 1:D:94:MET:HE3 | 0.53 | 1.80 | 6 | 1 |
| 1:A:111:ARG:NH1 | 1:C:105:ILE:HG23 | 0.53 | 2.18 | 20 | 2 |
| 1:A:20:ASP:HB3 | 1:A:79:GLN:HB2 | 0.52 | 1.81 | 5 | 5 |
| 1:C:60:LEU:HD22 | 1:C:103:PRO:HB2 | 0.52 | 1.80 | 1 | 1 |



Continued from previous page...

| Atom-1 | Atom-2 | $\operatorname{Clash}(ext{\AA})$ | $\operatorname{Distance}(\operatorname{\AA})$ | Models | |
|------------------|------------------|-----------------------------------|---|--------|-------|
| | | Clash(A) | Distance(A) | Worst | Total |
| 1:C:20:ASP:HB3 | 1:C:79:GLN:HB2 | 0.52 | 1.82 | 17 | 5 |
| 1:C:51:LEU:HD11 | 1:C:83:ILE:HD12 | 0.52 | 1.82 | 9 | 14 |
| 1:A:51:LEU:HD11 | 1:A:57:GLU:HB2 | 0.52 | 1.82 | 2 | 4 |
| 1:D:76:CYS:SG | 1:D:114:ILE:HG12 | 0.52 | 2.45 | 15 | 4 |
| 1:B:117:MET:SD | 1:B:120:ARG:HD2 | 0.51 | 2.45 | 20 | 3 |
| 1:C:19:LYS:HE3 | 1:D:28:ALA:HB3 | 0.51 | 1.81 | 3 | 2 |
| 1:B:100:ALA:HB1 | 1:B:134:ASP:HA | 0.51 | 1.83 | 2 | 2 |
| 1:C:97:CYS:SG | 1:C:102:CYS:SG | 0.51 | 3.08 | 17 | 3 |
| 1:B:64:VAL:HG21 | 1:B:114:ILE:HG21 | 0.51 | 1.83 | 8 | 4 |
| 1:A:61:ARG:HB2 | 1:A:79:GLN:HG2 | 0.51 | 1.83 | 14 | 1 |
| 1:D:38:PRO:HD3 | 1:D:124:PRO:HG2 | 0.51 | 1.83 | 15 | 1 |
| 1:A:47:ALA:HB3 | 1:A:59:VAL:HB | 0.51 | 1.83 | 16 | 4 |
| 1:D:31:VAL:HG22 | 1:D:68:LEU:HD12 | 0.50 | 1.83 | 10 | 4 |
| 1:B:12:GLN:HE22 | 1:D:129:ALA:HB2 | 0.50 | 1.67 | 15 | 1 |
| 1:D:25:ALA:HB1 | 1:D:28:ALA:HB2 | 0.50 | 1.83 | 8 | 1 |
| 1:D:11:PHE:HE2 | 1:D:102:CYS:SG | 0.50 | 2.29 | 2 | 2 |
| 1:A:45:ASP:HB3 | 1:A:61:ARG:HB3 | 0.49 | 1.84 | 2 | 5 |
| 1:D:40:VAL:HG21 | 1:D:126:LEU:HG | 0.49 | 1.85 | 1 | 1 |
| 1:B:21:ILE:CD1 | 1:B:113:CYS:SG | 0.49 | 3.00 | 19 | 5 |
| 1:C:19:LYS:HE2 | 1:D:29:PRO:HD3 | 0.49 | 1.84 | 8 | 2 |
| 1:C:100:ALA:HB1 | 1:C:134:ASP:HA | 0.49 | 1.84 | 11 | 1 |
| 1:B:21:ILE:HD12 | 1:B:113:CYS:SG | 0.49 | 2.47 | 19 | 2 |
| 1:D:54:ASP:HB3 | 1:D:86:ILE:O | 0.49 | 2.08 | 14 | 1 |
| 1:C:100:ALA:HB1 | 1:C:134:ASP:HB2 | 0.48 | 1.84 | 13 | 1 |
| 1:A:130:PRO:HG3 | 1:C:11:PHE:HD2 | 0.48 | 1.68 | 16 | 1 |
| 1:B:31:VAL:HG22 | 1:B:68:LEU:HD12 | 0.48 | 1.84 | 13 | 2 |
| 1:B:76:CYS:SG | 1:B:114:ILE:HA | 0.48 | 2.49 | 2 | 1 |
| 1:B:89:ILE:HG22 | 1:B:94:MET:HB2 | 0.48 | 1.85 | 6 | 1 |
| 1:D:11:PHE:HE2 | 1:D:102:CYS:HG | 0.48 | 1.52 | 1 | 1 |
| 1:A:11:PHE:HE2 | 1:A:84:PHE:HB3 | 0.47 | 1.69 | 2 | 11 |
| 1:B:86:ILE:HD13 | 1:B:98:LEU:HD21 | 0.47 | 1.86 | 2 | 1 |
| 1:B:9:MET:HB3 | 1:B:89:ILE:HG12 | 0.47 | 1.85 | 14 | 2 |
| 1:B:11:PHE:CE1 | 1:B:13:ILE:HG13 | 0.47 | 2.44 | 16 | 1 |
| 1:C:80:GLN:HB2 | 1:C:110:ALA:HB2 | 0.47 | 1.86 | 3 | 2 |
| 1:C:30:HIS:O | 1:C:33:GLN:HG2 | 0.47 | 2.09 | 11 | 2 |
| 1:B:125:GLN:HB2 | 1:C:124:PRO:HA | 0.46 | 1.85 | 7 | 3 |
| 1:D:86:ILE:HG13 | 1:D:97:CYS:SG | 0.46 | 2.51 | 20 | 1 |
| 1:A:111:ARG:HH12 | 1:C:105:ILE:HD12 | 0.46 | 1.71 | 3 | 1 |
| 1:B:9:MET:HB2 | 1:B:89:ILE:HB | 0.46 | 1.87 | 9 | 1 |
| 1:A:97:CYS:HA | 1:A:101:TYR:HB3 | 0.46 | 1.88 | 16 | 4 |
| 1:D:31:VAL:HG21 | 1:D:73:ALA:HA | 0.46 | 1.88 | 16 | 3 |
| | _ | _ | 01:1 | L | |



Continued from previous page...

| Atom-1 | Atom-2 | $oxed{ { m Clash}({ m \AA}) \ \ { m Distance}({ m \AA}) }$ | Models | | |
|-----------------|------------------|--|-------------|-------|-------|
| | | | Distance(A) | Worst | Total |
| 1:D:39:GLU:O | 1:D:66:ALA:HA | 0.46 | 2.11 | 16 | 1 |
| 1:B:40:VAL:HG13 | 1:B:64:VAL:HG13 | 0.46 | 1.88 | 12 | 1 |
| 1:A:13:ILE:HG12 | 1:A:84:PHE:HD2 | 0.45 | 1.71 | 14 | 7 |
| 1:B:56:TYR:HB2 | 1:B:86:ILE:HD11 | 0.45 | 1.89 | 7 | 2 |
| 1:B:22:SER:HB2 | 1:B:77:GLU:HB3 | 0.45 | 1.88 | 6 | 2 |
| 1:A:119:SER:HB2 | 1:A:125:GLN:HG3 | 0.45 | 1.89 | 20 | 1 |
| 1:B:10:THR:O | 1:B:86:ILE:HA | 0.45 | 2.12 | 1 | 1 |
| 1:C:94:MET:SD | 1:C:98:LEU:HD12 | 0.45 | 2.51 | 1 | 1 |
| 1:C:11:PHE:HE1 | 1:C:102:CYS:SG | 0.45 | 2.34 | 8 | 2 |
| 1:D:45:ASP:HB3 | 1:D:61:ARG:HB3 | 0.45 | 1.88 | 1 | 4 |
| 1:A:13:ILE:CD1 | 1:A:105:ILE:HG21 | 0.45 | 2.42 | 6 | 2 |
| 1:B:18:THR:HG21 | 1:B:21:ILE:HD11 | 0.45 | 1.88 | 11 | 2 |
| 1:B:130:PRO:HG3 | 1:D:16:ILE:HD13 | 0.45 | 1.87 | 17 | 1 |
| 1:A:111:ARG:NH1 | 1:C:105:ILE:HD12 | 0.45 | 2.27 | 1 | 5 |
| 1:B:9:MET:SD | 1:B:89:ILE:HB | 0.45 | 2.51 | 10 | 1 |
| 1:D:60:LEU:HD22 | 1:D:103:PRO:HB3 | 0.45 | 1.87 | 5 | 2 |
| 1:B:47:ALA:HB3 | 1:B:59:VAL:HB | 0.45 | 1.88 | 6 | 2 |
| 1:B:113:CYS:O | 1:B:117:MET:HG2 | 0.44 | 2.13 | 4 | 4 |
| 1:A:20:ASP:HA | 1:B:23:PHE:O | 0.44 | 2.12 | 17 | 3 |
| 1:D:76:CYS:SG | 1:D:114:ILE:HA | 0.44 | 2.52 | 16 | 1 |
| 1:A:28:ALA:HB3 | 1:A:29:PRO:HD3 | 0.44 | 1.88 | 3 | 2 |
| 1:B:61:ARG:HG3 | 1:B:79:GLN:HG2 | 0.44 | 1.90 | 19 | 1 |
| 1:D:90:GLU:HA | 1:D:94:MET:SD | 0.44 | 2.53 | 8 | 1 |
| 1:C:43:ASP:HB3 | 1:C:63:THR:HB | 0.44 | 1.90 | 10 | 2 |
| 1:B:9:MET:HB3 | 1:B:89:ILE:HB | 0.44 | 1.89 | 16 | 2 |
| 1:B:45:ASP:HB3 | 1:B:61:ARG:HB3 | 0.44 | 1.87 | 2 | 2 |
| 1:B:11:PHE:HE1 | 1:B:102:CYS:SG | 0.44 | 2.36 | 3 | 2 |
| 1:B:9:MET:SD | 1:B:89:ILE:HG22 | 0.44 | 2.53 | 9 | 1 |
| 1:A:96:HIS:O | 1:A:100:ALA:HB3 | 0.44 | 2.13 | 1 | 1 |
| 1:D:41:LYS:HB2 | 1:D:65:THR:HB | 0.44 | 1.90 | 19 | 1 |
| 1:C:96:HIS:O | 1:C:100:ALA:HB3 | 0.43 | 2.12 | 5 | 2 |
| 1:C:19:LYS:HE2 | 1:D:28:ALA:HB3 | 0.43 | 1.88 | 9 | 2 |
| 1:D:13:ILE:HG21 | 1:D:16:ILE:HD11 | 0.43 | 1.89 | 10 | 1 |
| 1:D:16:ILE:N | 1:D:16:ILE:HD12 | 0.43 | 2.27 | 10 | 1 |
| 1:C:11:PHE:HE2 | 1:C:84:PHE:HB3 | 0.43 | 1.73 | 1 | 3 |
| 1:C:90:GLU:HA | 1:C:94:MET:HE3 | 0.43 | 1.90 | 13 | 1 |
| 1:B:9:MET:SD | 1:B:11:PHE:HB2 | 0.43 | 2.54 | 4 | 1 |
| 1:C:51:LEU:HD11 | 1:C:57:GLU:HB2 | 0.43 | 1.90 | 8 | 1 |
| 1:A:120:ARG:HD3 | 1:B:113:CYS:SG | 0.43 | 2.53 | 11 | 1 |
| 1:C:21:ILE:HG12 | 1:C:113:CYS:SG | 0.43 | 2.53 | 11 | 1 |
| 1:C:64:VAL:HG21 | 1:C:114:ILE:HG21 | 0.43 | 1.89 | 15 | 1 |



Continued from previous page...

| | Atom 2 | $oxed{ 	ext{Clash(\AA)} \ 	ext{Distance(Å)} }$ | Distance (Å) | Models | |
|------------------|------------------|--|--------------|--------|-------|
| Atom-1 | Atom-2 | Clash(A) | | | Total |
| 1:A:43:ASP:HB3 | 1:A:63:THR:HB | 0.43 | 1.90 | 5 | 1 |
| 1:B:11:PHE:HE1 | 1:B:102:CYS:HG | 0.43 | 1.55 | 3 | 1 |
| 1:A:130:PRO:HD3 | 1:C:13:ILE:HG13 | 0.43 | 1.90 | 20 | 1 |
| 1:B:31:VAL:HG21 | 1:B:73:ALA:HA | 0.42 | 1.91 | 13 | 1 |
| 1:A:98:LEU:HD12 | 1:A:99:GLY:N | 0.42 | 2.29 | 1 | 1 |
| 1:C:76:CYS:SG | 1:C:114:ILE:HG23 | 0.42 | 2.54 | 11 | 4 |
| 1:B:98:LEU:HD23 | 1:B:98:LEU:H | 0.42 | 1.73 | 13 | 1 |
| 1:A:21:ILE:CG2 | 1:A:117:MET:SD | 0.42 | 3.07 | 2 | 1 |
| 1:C:11:PHE:CE1 | 1:C:101:TYR:CE2 | 0.42 | 3.07 | 17 | 3 |
| 1:A:104:ASN:HD21 | 1:A:133:PHE:HB2 | 0.42 | 1.73 | 9 | 1 |
| 1:A:13:ILE:HG21 | 1:C:130:PRO:HG2 | 0.42 | 1.91 | 14 | 1 |
| 1:D:57:GLU:HB2 | 1:D:83:ILE:HG12 | 0.42 | 1.90 | 14 | 1 |
| 1:B:51:LEU:HD11 | 1:B:57:GLU:HB2 | 0.42 | 1.90 | 15 | 1 |
| 1:B:94:MET:SD | 1:B:98:LEU:HD21 | 0.42 | 2.53 | 16 | 1 |
| 1:D:86:ILE:HB | 1:D:89:ILE:HD12 | 0.42 | 1.92 | 18 | 1 |
| 1:B:61:ARG:HB2 | 1:B:79:GLN:HG2 | 0.42 | 1.91 | 9 | 1 |
| 1:C:59:VAL:HG12 | 1:C:79:GLN:HG2 | 0.42 | 1.91 | 9 | 1 |
| 1:D:28:ALA:O | 1:D:31:VAL:HG12 | 0.42 | 2.15 | 5 | 3 |
| 1:A:40:VAL:HA | 1:A:65:THR:O | 0.42 | 2.15 | 17 | 1 |
| 1:D:86:ILE:HD13 | 1:D:98:LEU:HD21 | 0.42 | 1.91 | 2 | 1 |
| 1:B:96:HIS:O | 1:B:100:ALA:HB3 | 0.42 | 2.14 | 3 | 1 |
| 1:D:64:VAL:CG2 | 1:D:114:ILE:HD13 | 0.42 | 2.45 | 3 | 1 |
| 1:D:51:LEU:HD11 | 1:D:57:GLU:HB2 | 0.42 | 1.92 | 3 | 1 |
| 1:C:45:ASP:HB3 | 1:C:61:ARG:HB3 | 0.42 | 1.92 | 17 | 3 |
| 1:A:52:ALA:HB3 | 1:A:55:VAL:HB | 0.42 | 1.92 | 9 | 1 |
| 1:C:17:TYR:HB3 | 1:D:122:THR:HG23 | 0.42 | 1.92 | 19 | 1 |
| 1:B:56:TYR:HB3 | 1:B:86:ILE:HD13 | 0.41 | 1.90 | 12 | 1 |
| 1:A:116:SER:O | 1:A:120:ARG:HB2 | 0.41 | 2.15 | 19 | 2 |
| 1:C:44:LEU:HA | 1:C:61:ARG:O | 0.41 | 2.14 | 13 | 1 |
| 1:C:20:ASP:HA | 1:D:23:PHE:O | 0.41 | 2.15 | 14 | 2 |
| 1:A:21:ILE:HD13 | 1:B:117:MET:SD | 0.41 | 2.56 | 16 | 1 |
| 1:C:18:THR:HG22 | 1:C:80:GLN:HE22 | 0.41 | 1.74 | 5 | 1 |
| 1:C:47:ALA:HB3 | 1:C:59:VAL:HB | 0.41 | 1.93 | 17 | 1 |
| 1:D:89:ILE:HG22 | 1:D:90:GLU:N | 0.41 | 2.31 | 11 | 1 |
| 1:D:14:GLN:HE21 | 1:D:85:SER:HB2 | 0.41 | 1.76 | 10 | 1 |
| 1:D:56:TYR:HB3 | 1:D:86:ILE:HD13 | 0.41 | 1.91 | 11 | 2 |
| 1:B:42:LEU:HD13 | 1:B:128:LEU:HD21 | 0.41 | 1.91 | 15 | 1 |
| 1:D:41:LYS:HB3 | 1:D:65:THR:HB | 0.41 | 1.92 | 12 | 2 |
| 1:A:23:PHE:O | 1:B:20:ASP:HA | 0.41 | 2.15 | 20 | 2 |
| 1:C:44:LEU:HD23 | 1:C:60:LEU:HD11 | 0.41 | 1.91 | 13 | 1 |
| 1:D:11:PHE:CE2 | 1:D:102:CYS:SG | 0.41 | 3.13 | 2 | 1 |



Continued from previous page...

| Atom-1 | Atom-2 | $\operatorname{Clash}(\mathring{\mathrm{A}})$ | $\mathbf{Distance}(\mathbf{\mathring{A}})$ | Models | |
|------------------|------------------|---|--|--------|-------|
| Atom-1 | Atom-2 | Clash(A) | Distance(A) | Worst | Total |
| 1:A:34:LYS:HD3 | 1:A:35:ASP:N | 0.40 | 2.31 | 5 | 1 |
| 1:B:76:CYS:SG | 1:B:114:ILE:HG23 | 0.40 | 2.56 | 8 | 1 |
| 1:B:28:ALA:HB3 | 1:B:29:PRO:HD3 | 0.40 | 1.92 | 11 | 1 |
| 1:C:104:ASN:HD21 | 1:C:133:PHE:HB3 | 0.40 | 1.75 | 11 | 1 |
| 1:B:60:LEU:HD23 | 1:B:106:LEU:HD12 | 0.40 | 1.93 | 13 | 1 |
| 1:C:19:LYS:HE2 | 1:C:59:VAL:HG21 | 0.40 | 1.93 | 14 | 1 |
| 1:D:13:ILE:HD13 | 1:D:105:ILE:HG21 | 0.40 | 1.94 | 2 | 1 |
| 1:B:114:ILE:O | 1:B:118:VAL:HG23 | 0.40 | 2.16 | 17 | 1 |
| 1:C:14:GLN:HB2 | 1:C:83:ILE:HB | 0.40 | 1.93 | 1 | 2 |
| 1:C:61:ARG:HB2 | 1:C:79:GLN:HG2 | 0.40 | 1.94 | 2 | 1 |
| 1:C:44:LEU:HD23 | 1:C:45:ASP:N | 0.40 | 2.31 | 6 | 1 |
| 1:B:34:LYS:HB2 | 1:B:68:LEU:HD13 | 0.40 | 1.93 | 10 | 1 |
| 1:C:98:LEU:HA | 1:C:102:CYS:SG | 0.40 | 2.57 | 16 | 1 |
| 1:C:94:MET:O | 1:C:98:LEU:HG | 0.40 | 2.16 | 16 | 1 |
| 1:D:14:GLN:HB2 | 1:D:83:ILE:HB | 0.40 | 1.92 | 19 | 1 |
| 1:D:114:ILE:O | 1:D:118:VAL:HG23 | 0.40 | 2.16 | 3 | 1 |
| 1:B:125:GLN:HA | 1:B:125:GLN:HE21 | 0.40 | 1.77 | 9 | 1 |
| 1:C:17:TYR:CE2 | 1:C:19:LYS:HD3 | 0.40 | 2.51 | 15 | 1 |

5.2 Torsion angles (i)

5.2.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percei | ntiles |
|-----|-------|---------------------|--------------------------|-----------------------|------------|--------|--------|
| 1 | A | 115/155~(74%) | $109\pm2~(95\pm1\%)$ | $6\pm 2 \ (5\pm 1\%)$ | 0±0 (0±0%) | 100 | 100 |
| 1 | В | 126/155~(81%) | 119±2 (94±2%) | $7\pm2~(6\pm2\%)$ | 0±0 (0±0%) | 50 | 82 |
| 1 | С | $124/155 \; (80\%)$ | $115\pm 2 \ (93\pm 2\%)$ | 9±2 (7±2%) | 0±0 (0±0%) | 54 | 85 |
| 1 | D | 122/155~(79%) | $116\pm 2 \ (95\pm 1\%)$ | 5±2 (4±1%) | 0±0 (0±0%) | 50 | 82 |
| 2 | Е | 0 | - | - | - | - | |
| 2 | F | 0 | - | - | - | - | |
| 2 | G | 0 | - | - | - | - | |
| 2 | Н | 0 | - | - | - | - | |
| All | All | 9740/14400 (68%) | 9171 (94%) | 556 (6%) | 13 (0%) | 54 | 85 |



All 4 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

| Mol | Chain | Res | Type | Models (Total) |
|-----|-------|-----|------|----------------|
| 1 | D | 28 | ALA | 6 |
| 1 | В | 9 | MET | 3 |
| 1 | С | 130 | PRO | 2 |
| 1 | В | 130 | PRO | 2 |

5.2.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles |
|-----|-------|-----------------------|----------------------|-----------------------|-------------|
| 1 | A | 100/132~(76%) | 98±1 (98±1%) | 2±1 (2±1%) | 59 93 |
| 1 | В | $108/132 \; (82\%)$ | $105\pm2~(97\pm2\%)$ | $3\pm 2 \ (3\pm 2\%)$ | 43 88 |
| 1 | С | $106/132 \; (80\%)$ | 102±1 (96±1%) | 4±1 (4±1%) | 36 84 |
| 1 | D | $104/132 \ (79\%)$ | 102±1 (98±1%) | 2±1 (2±1%) | 57 93 |
| 2 | Е | 0 | - | - | - |
| 2 | F | 0 | - | - | - |
| 2 | G | 0 | - | - | - |
| 2 | Н | 0 | - | - | - |
| All | All | $8360/12320 \ (68\%)$ | 8130 (97%) | 230 (3%) | 46 90 |

All 70 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

| Mol | Chain | Res | Type | Models (Total) |
|-----|-------|-----|------|----------------|
| 1 | С | 34 | LYS | 20 |
| 1 | С | 117 | MET | 20 |
| 1 | В | 12 | GLN | 15 |
| 1 | D | 70 | GLU | 13 |
| 1 | A | 86 | ILE | 7 |
| 1 | D | 90 | GLU | 7 |
| 1 | В | 125 | GLN | 7 |
| 1 | С | 112 | GLU | 6 |
| 1 | D | 93 | GLN | 6 |
| 1 | В | 94 | MET | 6 |
| 1 | В | 11 | PHE | 6 |
| 1 | A | 60 | LEU | 5 |

Continued on next page...



Continued from previous page...

| Mol | nued fron Chain | \mathbf{Res} | $\overline{	ext{Type}}$ | Models (Total) |
|-----|--------------------|----------------|-------------------------|----------------|
| 1 | A | 42 | LEU | 5 |
| 1 | В | 9 | MET | 4 |
| 1 | В | 42 | LEU | 4 |
| 1 | A | 14 | GLN | 4 |
| 1 | С | 126 | LEU | 4 |
| 1 | С | 128 | LEU | 4 |
| 1 | С | 11 | PHE | 4 |
| 1 | В | 86 | ILE | 3 |
| 1 | С | 44 | LEU | 3 |
| 1 | С | 86 | ILE | 3 |
| 1 | С | 125 | GLN | 3 |
| 1 | В | 70 | GLU | 3 |
| 1 | A | 56 | TYR | 3 |
| 1 | A | 27 | ASN | 3 |
| 1 | В | 89 | ILE | 3 |
| 1 | D | 86 | ILE | 3 |
| 1 | D | 42 | LEU | 3 |
| 1 | С | 104 | ASN | 2 |
| 1 | В | 126 | LEU | 2 |
| 1 | A | 70 | GLU | 2 |
| 1 | С | 61 | ARG | 2 |
| 1 | С | 107 | PHE | 2 |
| 1 | В | 98 | LEU | 2 |
| 1 | С | 39 | GLU | 2 |
| 1 | С | 98 | LEU | 2 |
| 1 | С | 42 | LEU | 2 |
| 1 | A | 122 | THR | 2 |
| 1 | В | 112 | GLU | 2 |
| 1 | В | 132 | ASN | 2 |
| 1 | D | 98 | LEU | 1 |
| 1 | В | 107 | PHE | 1 |
| 1 | D | 127 | ASN | 1 |
| 1 | В | 60 | LEU | 1 |
| 1 | В | 39 | GLU | 1 |
| 1 | В | 93 | GLN | 1 |
| 1 | A | 125 | GLN | 1 |
| 1 | D | 39 | GLU | 1 |
| 1 | С | 14 | GLN | 1 |
| 1 | A | 98 | LEU | 1 |
| 1 | D | 104 | ASN | 1 |
| 1 | D | 41 | LYS | 1 |
| 1 | В | 128 | LEU | 1 |

Continued on next page...



Continued from previous page...

| Mol | Chain | Res | Type | Models (Total) |
|-----|-------|-----|------|----------------|
| 1 | A | 54 | ASP | 1 |
| 1 | D | 56 | TYR | 1 |
| 1 | В | 56 | TYR | 1 |
| 1 | D | 60 | LEU | 1 |
| 1 | С | 70 | GLU | 1 |
| 1 | A | 31 | VAL | 1 |
| 1 | D | 94 | MET | 1 |
| 1 | В | 35 | ASP | 1 |
| 1 | В | 104 | ASN | 1 |
| 1 | В | 90 | GLU | 1 |
| 1 | D | 126 | LEU | 1 |
| 1 | С | 93 | GLN | 1 |
| 1 | A | 101 | TYR | 1 |
| 1 | A | 128 | LEU | 1 |
| 1 | С | 37 | GLN | 1 |
| 1 | A | 41 | LYS | 1 |

5.2.3 RNA (i)

There are no RNA molecules in this entry.

5.3 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.4 Carbohydrates (i)

There are no carbohydrates in this entry.

5.5 Ligand geometry (i)

There are no ligands in this entry.

5.6 Other polymers (i)

There are no such molecules in this entry.



5.7 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 13% for the well-defined parts and 13% for the entire structure.

6.1 Chemical shift list 1

File name: input cs.cif

Chemical shift list name: assigned_chemical_shift_5

6.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

| Total number of shifts | 226 |
|---|-----|
| Number of shifts mapped to atoms | 226 |
| Number of unparsed shifts | 0 |
| Number of shifts with mapping errors | 0 |
| Number of shifts with mapping warnings | 0 |
| Number of shift outliers (ShiftChecker) | 0 |

6.1.2 Chemical shift referencing (i)

No chemical shift referencing corrections were calculated (not enough data).

6.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 0%, i.e. 0 atoms were assigned a chemical shift out of a possible 5840. 0 out of 80 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

| | Total | $^{1}\mathrm{H}$ | $^{13}\mathbf{C}$ | $^{15}{ m N}$ |
|-----------|------------------|------------------|-------------------|-----------------|
| Backbone | $0/2379 \ (0\%)$ | 0/946~(0%) | 0/974 (0%) | $0/459 \ (0\%)$ |
| Sidechain | 0/2942~(0%) | 0/1716~(0%) | 0/1116 (0%) | 0/110 (0%) |
| Aromatic | 0/519 (0%) | 0/275~(0%) | 0/224~(0%) | 0/20~(0%) |
| Overall | 0/5840 (0%) | 0/2937~(0%) | 0/2314~(0%) | 0/589 (0%) |

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 2%, i.e. 192 atoms were assigned a chemical shift out of a possible 8632. 0 out of 112 assigned methyl groups (LEU and VAL) were assigned stereospecifically.



| | Total | $^{1}\mathrm{H}$ | $^{13}\mathbf{C}$ | $^{15}{ m N}$ |
|-----------|---------------|------------------|-------------------|---------------|
| Backbone | 77/3520~(2%) | 37/1400 (3%) | 19/1440 (1%) | 21/680 (3%) |
| Sidechain | 106/4448~(2%) | 53/2596 (2%) | 53/1684 (3%) | 0/168 (0%) |
| Aromatic | 9/664~(1%) | 5/352 (1%) | 4/284 (1%) | 0/28 (0%) |
| Overall | 192/8632 (2%) | 95/4348 (2%) | 76/3408 (2%) | 21/876 (2%) |


6.1.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.


6.1.5 Random Coil Index (RCI) plots (i)

The images below report random coil index values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:



Random coil index (RCI) for chain E:





6.2 Chemical shift list 2

File name: input_cs.cif

Chemical shift list name: assigned_chemical_shift_6

6.2.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

| Total number of shifts | 226 |
|---|-----|
| Number of shifts mapped to atoms | 226 |
| Number of unparsed shifts | 0 |
| Number of shifts with mapping errors | 0 |
| Number of shifts with mapping warnings | 0 |
| Number of shift outliers (ShiftChecker) | 0 |

6.2.2 Chemical shift referencing (i)

No chemical shift referencing corrections were calculated (not enough data).

6.2.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 0%, i.e. 0 atoms were assigned a chemical shift out of a possible 5840. 0 out of 80 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

| | Total | $^{1}\mathbf{H}$ | $^{13}\mathbf{C}$ | $^{15}{ m N}$ |
|-----------|------------------|------------------|-------------------|---------------|
| Backbone | $0/2379 \ (0\%)$ | 0/946~(0%) | 0/974 (0%) | 0/459~(0%) |
| Sidechain | 0/2942~(0%) | $0/1716 \ (0\%)$ | 0/1116 (0%) | 0/110 (0%) |
| Aromatic | 0/519 (0%) | 0/275~(0%) | 0/224~(0%) | 0/20~(0%) |
| Overall | 0/5840 (0%) | 0/2937~(0%) | 0/2314~(0%) | 0/589 (0%) |

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 2%, i.e. 192 atoms were assigned a chemical shift out of a possible 8632. 0 out of 112 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

| | Total | $^{1}\mathrm{H}$ | $^{13}\mathbf{C}$ | $^{15}{ m N}$ |
|-----------|---------------|------------------|-------------------|---------------|
| Backbone | 77/3520 (2%) | 37/1400 (3%) | 19/1440 (1%) | 21/680 (3%) |
| Sidechain | 106/4448~(2%) | 53/2596~(2%) | 53/1684 (3%) | 0/168 (0%) |
| Aromatic | 9/664 (1%) | 5/352 (1%) | 4/284 (1%) | 0/28 (0%) |
| Overall | 192/8632 (2%) | 95/4348 (2%) | 76/3408 (2%) | 21/876 (2%) |




6.2.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.


6.2.5 Random Coil Index (RCI) plots (1)

The images below report random coil index values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:



Random coil index (RCI) for chain E:



6.3 Chemical shift list 3

File name: input cs.cif

Chemical shift list name: assigned chemical shift 7



6.3.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

| Total number of shifts | 226 |
|---|-----|
| Number of shifts mapped to atoms | 226 |
| Number of unparsed shifts | 0 |
| Number of shifts with mapping errors | 0 |
| Number of shifts with mapping warnings | 0 |
| Number of shift outliers (ShiftChecker) | 0 |

6.3.2 Chemical shift referencing (i)

No chemical shift referencing corrections were calculated (not enough data).

6.3.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 0%, i.e. 0 atoms were assigned a chemical shift out of a possible 5840. 0 out of 80 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

| | Total | $^{1}{ m H}$ | $^{13}\mathbf{C}$ | $^{15}{ m N}$ |
|-----------|-------------|--------------|-------------------|---------------|
| Backbone | 0/2379 (0%) | 0/946~(0%) | 0/974 (0%) | 0/459~(0%) |
| Sidechain | 0/2942 (0%) | 0/1716~(0%) | 0/1116 (0%) | 0/110 (0%) |
| Aromatic | 0/519 (0%) | 0/275~(0%) | 0/224~(0%) | 0/20~(0%) |
| Overall | 0/5840 (0%) | 0/2937~(0%) | 0/2314 (0%) | 0/589~(0%) |

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 2%, i.e. 192 atoms were assigned a chemical shift out of a possible 8632. 0 out of 112 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

| | Total | $^{1}\mathrm{H}$ | $^{13}\mathbf{C}$ | $^{15}{ m N}$ |
|-----------|--------------------|------------------|-------------------|---------------|
| Backbone | 77/3520~(2%) | 37/1400 (3%) | $19/1440 \ (1\%)$ | 21/680 (3%) |
| Sidechain | $106/4448 \ (2\%)$ | 53/2596 (2%) | 53/1684 (3%) | 0/168 (0%) |
| Aromatic | 9/664 (1%) | 5/352 (1%) | 4/284 (1%) | 0/28 (0%) |
| Overall | 192/8632~(2%) | 95/4348 (2%) | 76/3408~(2%) | 21/876 (2%) |

6.3.4 Statistically unusual chemical shifts (i)


There are no statistically unusual chemical shifts.




6.3.5 Random Coil Index (RCI) plots (i)

The images below report random coil index values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:



Random coil index (RCI) for chain E:



6.4 Chemical shift list 4

File name: input cs.cif

Chemical shift list name: assigned chemical shift 8

6.4.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

| Total number of shifts | |
|----------------------------------|-----|
| Number of shifts mapped to atoms | 226 |



| Number of unparsed shifts | | |
|---|---|--|
| Number of shifts with mapping errors | 0 | |
| Number of shifts with mapping warnings | 0 | |
| Number of shift outliers (ShiftChecker) | 0 | |

6.4.2 Chemical shift referencing (i)

No chemical shift referencing corrections were calculated (not enough data).

6.4.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 0%, i.e. 0 atoms were assigned a chemical shift out of a possible 5840. 0 out of 80 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

| | Total | $^{1}{ m H}$ | $^{13}\mathbf{C}$ | $^{15}{ m N}$ |
|-----------|------------------|--------------|-------------------|---------------|
| Backbone | $0/2379 \ (0\%)$ | 0/946~(0%) | 0/974 (0%) | 0/459~(0%) |
| Sidechain | $0/2942 \ (0\%)$ | 0/1716~(0%) | 0/1116 (0%) | 0/110 (0%) |
| Aromatic | 0/519 (0%) | 0/275~(0%) | 0/224~(0%) | 0/20 (0%) |
| Overall | 0/5840 (0%) | 0/2937~(0%) | 0/2314~(0%) | 0/589 (0%) |

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 2%, i.e. 192 atoms were assigned a chemical shift out of a possible 8632. 0 out of 112 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

| | Total | $^{1}\mathrm{H}$ | $^{13}\mathbf{C}$ | $^{15}{ m N}$ |
|-----------|---------------|------------------|-------------------|---------------|
| Backbone | 77/3520~(2%) | 37/1400 (3%) | 19/1440 (1%) | 21/680 (3%) |
| Sidechain | 106/4448~(2%) | 53/2596~(2%) | 53/1684 (3%) | 0/168 (0%) |
| Aromatic | 9/664 (1%) | 5/352 (1%) | 4/284 (1%) | 0/28 (0%) |
| Overall | 192/8632~(2%) | 95/4348 (2%) | 76/3408 (2%) | 21/876 (2%) |

6.4.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.


6.4.5 Random Coil Index (RCI) plots (i)

The images below report random coil index values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication




of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:



Random coil index (RCI) for chain E:



6.5 Chemical shift list 5

File name: input cs.cif

Chemical shift list name: assigned chemical shift list 1

6.5.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

| Total number of shifts | 1049 |
|---|------|
| Number of shifts mapped to atoms | 1049 |
| Number of unparsed shifts | 0 |
| Number of shifts with mapping errors | 0 |
| Number of shifts with mapping warnings | 0 |
| Number of shift outliers (ShiftChecker) | 0 |



6.5.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

| Nucleus | # values | ${\bf Correction}\pm{\bf precision},ppm$ | Suggested action |
|----------------------------|----------|--|-------------------------------------|
| $^{13}\mathrm{C}_{\alpha}$ | 139 | 0.26 ± 0.12 | None needed ($< 0.5 \text{ ppm}$) |
| $^{13}C_{\beta}$ | 128 | 0.85 ± 0.16 | Should be applied |
| ¹³ C′ | 137 | 0.30 ± 0.16 | None needed ($< 0.5 \text{ ppm}$) |
| ^{15}N | 133 | -1.15 ± 0.27 | Should be applied |

6.5.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 13%, i.e. 740 atoms were assigned a chemical shift out of a possible 5840. 0 out of 80 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

| | Total | $^{1}\mathrm{H}$ | $^{13}\mathbf{C}$ | $^{15}{ m N}$ |
|-----------|----------------|--------------------|---------------------|---------------|
| Backbone | 422/2379 (18%) | 103/946 (11%) | 216/974~(22%) | 103/459 (22%) |
| Sidechain | 205/2942~(7%) | $57/1716 \ (3\%)$ | $148/1116 \ (13\%)$ | 0/110 (0%) |
| Aromatic | 113/519 (22%) | 57/275 (21%) | 55/224~(25%) | 1/20 (5%) |
| Overall | 740/5840 (13%) | $217/2937 \ (7\%)$ | 419/2314 (18%) | 104/589 (18%) |

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 11%, i.e. 930 atoms were assigned a chemical shift out of a possible 8632. 0 out of 112 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

| | Total | $^{1}\mathrm{H}$ | $^{13}\mathbf{C}$ | $^{15}{ m N}$ |
|-----------|----------------|---------------------|-------------------|------------------|
| Backbone | 542/3520 (15%) | $133/1400 \ (10\%)$ | 276/1440 (19%) | 133/680 (20%) |
| Sidechain | 255/4448~(6%) | 70/2596~(3%) | 185/1684 (11%) | $0/168 \; (0\%)$ |
| Aromatic | 133/664 (20%) | 67/352 (19%) | 65/284~(23%) | $1/28 \ (4\%)$ |
| Overall | 930/8632 (11%) | 270/4348~(6%) | 526/3408 (15%) | 134/876 (15%) |

6.5.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.


6.5.5 Random Coil Index (RCI) plots (i)

The image below reports random coil index values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-



defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:



6.6 Chemical shift list 6

File name: input cs.cif

Chemical shift list name: assigned chemical shift list 2

6.6.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

| Total number of shifts | 1044 |
|---|------|
| Number of shifts mapped to atoms | 1044 |
| Number of unparsed shifts | 0 |
| Number of shifts with mapping errors | 0 |
| Number of shifts with mapping warnings | 0 |
| Number of shift outliers (ShiftChecker) | 0 |

6.6.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

| Nucleus | # values | ${\bf Correction}\pm{\bf precision},ppm$ | Suggested action |
|----------------------------|----------|--|-------------------------------------|
| $^{13}\mathrm{C}_{\alpha}$ | 140 | 0.28 ± 0.07 | None needed ($< 0.5 \text{ ppm}$) |
| $^{13}C_{\beta}$ | 126 | 0.79 ± 0.12 | Should be applied |
| ¹³ C′ | 135 | 0.26 ± 0.10 | None needed ($< 0.5 \text{ ppm}$) |
| ^{15}N | 132 | -1.10 ± 0.16 | Should be applied |



6.6.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 13%, i.e. 735 atoms were assigned a chemical shift out of a possible 5840. 0 out of 80 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

| | Total | $^{1}\mathrm{H}$ | $^{13}\mathbf{C}$ | $^{15}{ m N}$ |
|-----------|----------------|--------------------|-------------------|--------------------|
| Backbone | 419/2379 (18%) | 102/946 (11%) | 215/974~(22%) | $102/459 \ (22\%)$ |
| Sidechain | 203/2942~(7%) | $57/1716 \ (3\%)$ | 146/1116 (13%) | 0/110 (0%) |
| Aromatic | 113/519 (22%) | 57/275~(21%) | 55/224~(25%) | 1/20~(5%) |
| Overall | 735/5840 (13%) | $216/2937 \ (7\%)$ | 416/2314 (18%) | 103/589 (17%) |

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 11%, i.e. 925 atoms were assigned a chemical shift out of a possible 8632. 0 out of 112 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

| | Total | $^{1}\mathrm{H}$ | $^{13}\mathbf{C}$ | $^{15}{ m N}$ |
|-----------|----------------|--------------------|---------------------|---------------|
| Backbone | 539/3520 (15%) | $132/1400 \ (9\%)$ | $275/1440 \ (19\%)$ | 132/680 (19%) |
| Sidechain | 253/4448~(6%) | 70/2596~(3%) | 183/1684 (11%) | 0/168 (0%) |
| Aromatic | 133/664 (20%) | $67/352 \ (19\%)$ | 65/284~(23%) | 1/28 (4%) |
| Overall | 925/8632 (11%) | 269/4348~(6%) | 523/3408 (15%) | 133/876 (15%) |

6.6.4 Statistically unusual chemical shifts (i)


There are no statistically unusual chemical shifts.

6.6.5 Random Coil Index (RCI) plots (i)

The image below reports random coil index values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:





6.7 Chemical shift list 7

File name: input_cs.cif

Chemical shift list name: $assigned_chemical_shift_list_3$

6.7.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

| Total number of shifts | 1044 |
|---|------|
| Number of shifts mapped to atoms | 1044 |
| Number of unparsed shifts | 0 |
| Number of shifts with mapping errors | 0 |
| Number of shifts with mapping warnings | 0 |
| Number of shift outliers (ShiftChecker) | 0 |

6.7.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

| Nucleus | # values | ${\bf Correction}\pm{\bf precision},ppm$ | Suggested action |
|----------------------------|----------|--|-------------------------------------|
| $^{13}\mathrm{C}_{\alpha}$ | 140 | 0.27 ± 0.10 | None needed ($< 0.5 \text{ ppm}$) |
| $^{13}C_{\beta}$ | 126 | 0.78 ± 0.17 | Should be applied |
| ¹³ C′ | 135 | 0.26 ± 0.10 | None needed ($< 0.5 \text{ ppm}$) |
| ^{15}N | 132 | -1.09 ± 0.21 | Should be applied |

6.7.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 13%, i.e. 735 atoms were assigned a chemical shift out of a possible 5840. 0 out of 80 assigned methyl groups (LEU and VAL) were assigned



stereospecifically.

| | Total | $^{1}\mathrm{H}$ | $^{13}{ m C}$ | $^{15}{ m N}$ |
|-----------|---------------------|--------------------|---------------------|---------------|
| Backbone | $419/2379 \ (18\%)$ | 102/946~(11%) | 215/974~(22%) | 102/459~(22%) |
| Sidechain | 203/2942 (7%) | $57/1716 \ (3\%)$ | 146/1116 (13%) | 0/110 (0%) |
| Aromatic | $113/519 \ (22\%)$ | 57/275~(21%) | 55/224~(25%) | 1/20~(5%) |
| Overall | 735/5840 (13%) | $216/2937 \ (7\%)$ | $416/2314 \ (18\%)$ | 103/589 (17%) |

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 11%, i.e. 925 atoms were assigned a chemical shift out of a possible 8632. 0 out of 112 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

| | Total | $^{1}{ m H}$ | $^{13}\mathbf{C}$ | $^{15}{ m N}$ |
|-----------|----------------|--------------------|---------------------|---------------|
| Backbone | 539/3520 (15%) | $132/1400 \ (9\%)$ | $275/1440 \ (19\%)$ | 132/680 (19%) |
| Sidechain | 253/4448 (6%) | 70/2596~(3%) | 183/1684 (11%) | 0/168 (0%) |
| Aromatic | 133/664 (20%) | $67/352 \ (19\%)$ | 65/284~(23%) | 1/28 (4%) |
| Overall | 925/8632 (11%) | 269/4348~(6%) | 523/3408 (15%) | 133/876 (15%) |


6.7.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

6.7.5 Random Coil Index (RCI) plots (i)

The image below reports random coil index values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:





6.8 Chemical shift list 8

File name: input_cs.cif

Chemical shift list name: assigned_chemical_shift_list_4

6.8.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

| Total number of shifts | 1049 |
|---|------|
| Number of shifts mapped to atoms | 1049 |
| Number of unparsed shifts | 0 |
| Number of shifts with mapping errors | 0 |
| Number of shifts with mapping warnings | 0 |
| Number of shift outliers (ShiftChecker) | 0 |

6.8.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

| Nucleus | # values | ${\bf Correction} \pm {\bf precision}, ppm$ | Suggested action |
|----------------------------|----------|--|-------------------------------------|
| $^{13}\mathrm{C}_{\alpha}$ | 139 | 0.26 ± 0.09 | None needed ($< 0.5 \text{ ppm}$) |
| $^{13}C_{\beta}$ | 128 | 0.84 ± 0.10 | Should be applied |
| ¹³ C′ | 137 | 0.29 ± 0.12 | None needed ($< 0.5 \text{ ppm}$) |
| ^{15}N | 133 | -1.15 ± 0.26 | Should be applied |

6.8.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 13%, i.e. 740 atoms were assigned a chemical shift out of a possible 5840. 0 out of 80 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

| | Total | $^{1}\mathrm{H}$ | $^{13}\mathbf{C}$ | $^{15}{ m N}$ |
|-----------|---------------------|--------------------|---------------------|--------------------|
| Backbone | $422/2379 \ (18\%)$ | 103/946 (11%) | 216/974~(22%) | $103/459 \ (22\%)$ |
| Sidechain | $205/2942 \ (7\%)$ | 57/1716 (3%) | 148/1116 (13%) | 0/110 (0%) |
| Aromatic | $113/519 \ (22\%)$ | 57/275~(21%) | 55/224~(25%) | 1/20~(5%) |
| Overall | $740/5840 \ (13\%)$ | $217/2937 \ (7\%)$ | $419/2314 \ (18\%)$ | $104/589 \ (18\%)$ |

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 11%, i.e. 930 atoms were assigned a chemical shift out of a possible 8632. 0 out of 112 assigned methyl groups (LEU and VAL) were assigned stereospecifically.



| | Total | $^{1}\mathrm{H}$ | $^{13}\mathbf{C}$ | $^{15}{ m N}$ |
|-----------|----------------|---------------------|---------------------|--------------------|
| Backbone | 542/3520 (15%) | $133/1400 \ (10\%)$ | $276/1440 \ (19\%)$ | $133/680 \ (20\%)$ |
| Sidechain | 255/4448~(6%) | $70/2596 \ (3\%)$ | 185/1684 (11%) | 0/168 (0%) |
| Aromatic | 133/664 (20%) | 67/352 (19%) | 65/284~(23%) | $1/28 \ (4\%)$ |
| Overall | 930/8632 (11%) | 270/4348~(6%) | 526/3408 (15%) | 134/876 (15%) |

6.8.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

6.8.5 Random Coil Index (RCI) plots (i)

The image below reports random coil index values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

