

# wwPDB X-ray Structure Validation Summary Report (i)

May 13, 2020 – 02:55 am BST

PDB ID : 5JX0

> Title : Temperature sensitive D4 mutant L110F

Authors Schormann, N.; Chattopadhyay, D.

2016-05-12 Deposited on

2.40 Å(reported) Resolution

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity 4.02b-467

> 1.8.5 (274361), CSD as541be (2020) Mogul

Xtriage (Phenix) 1.13 EDS 2.11

Percentile statistics 20191225.v01 (using entries in the PDB archive December 25th 2019)

> Refmac 5.8.0158

CCP4 7.0.044 (Gargrove) Engh & Huber (2001)

Ideal geometry (proteins) Ideal geometry (DNA, RNA) Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) 2.11

### 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$ 

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$egin{aligned}  ext{Similar resolution} \ (\# ext{Entries},  ext{resolution range}( ext{Å})) \end{aligned}$
$R_{free}$	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	238	87%		8%
1	В	238	86%	5%	9%
1	С	238	87%		9%
1	D	238	84%	6	11%



## 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 7207 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Uracil-DNA glycosylase.

Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf	Trace
1	Λ	218	Total	С	N	О	S	0	0	0
1	A	210	1764	1144	291	323	6	U	U	
1	В	216	Total	С	N	О	S	0	0	0
1	Б	210	1754	1138	289	321	6	0	U	U
1	С	216	Total	С	N	О	S	0	0	0
1		210	1754	1138	289	321	6	0	U	
1	D	213	Total	С	N	О	S	0	0	0
1	ש	213	1732	1123	285	318	6	0	U	U

There are 92 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP P04303
A	-18	GLY	-	expression tag	UNP P04303
A	-17	SER	-	expression tag	UNP P04303
A	-16	SER	_	expression tag	UNP P04303
A	-15	HIS	-	expression tag	UNP P04303
A	-14	HIS	_	expression tag	UNP P04303
A	-13	HIS	_	expression tag	UNP P04303
A	-12	HIS	_	expression tag	UNP P04303
A	-11	HIS	_	expression tag	UNP P04303
A	-10	HIS	_	expression tag	UNP P04303
A	-9	SER	_	expression tag	UNP P04303
A	-8	SER	_	expression tag	UNP P04303
A	-7	GLY	_	expression tag	UNP P04303
A	-6	LEU	_	expression tag	UNP P04303
A	-5	VAL	-	expression tag	UNP P04303
A	-4	PRO	_	expression tag	UNP P04303
A	-3	ARG		expression tag	UNP P04303
A	-2	GLY	-	expression tag	UNP P04303
A	-1	SER	-	expression tag	UNP P04303
A	0	HIS	-	expression tag	UNP P04303
A	17	ASN	ASP	see remark 999	UNP P04303

Continued on next page...



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	110	PHE	LEU	engineered mutation	UNP P04303
A	163	РНЕ	TYR	see remark 999	UNP P04303
В	-19	MET	-	initiating methionine	UNP P04303
В	-18	GLY	-	expression tag	UNP P04303
В	-17	SER	-	expression tag	UNP P04303
В	-16	SER	_	expression tag	UNP P04303
В	-15	HIS	_	expression tag	UNP P04303
В	-14	HIS	_	expression tag	UNP P04303
В	-13	HIS	_	expression tag	UNP P04303
В	-12	HIS	-	expression tag	UNP P04303
В	-11	HIS	_	expression tag	UNP P04303
В	-10	HIS	_	expression tag	UNP P04303
В	-9	SER	_	expression tag	UNP P04303
В	-8	SER	_	expression tag	UNP P04303
В	-7	GLY	_	expression tag	UNP P04303
В	-6	LEU	_	expression tag	UNP P04303
В	-5	VAL	-	expression tag	UNP P04303
В	-4	PRO	_	expression tag	UNP P04303
В	-3	ARG	_	expression tag	UNP P04303
В	-2	GLY	-	expression tag	UNP P04303
В	-1	SER	-	expression tag	UNP P04303
В	0	HIS	-	expression tag	UNP P04303
В	17	ASN	ASP	see remark 999	UNP P04303
В	110	PHE	LEU	engineered mutation	UNP P04303
В	163	PHE	TYR	see remark 999	UNP P04303
С	-19	MET	-	initiating methionine	UNP P04303
С	-18	GLY	_	expression tag	UNP P04303
С	-17	SER	-	expression tag	UNP P04303
С	-16	SER	_	expression tag	UNP P04303
С	-15	HIS	_	expression tag	UNP P04303
С	-14	HIS	-	expression tag	UNP P04303
С	-13	HIS	_	expression tag	UNP P04303
С	-12	HIS	-	expression tag	UNP P04303
С	-11	HIS	-	expression tag	UNP P04303
С	-10	HIS	-	expression tag	UNP P04303
С	-9	SER	=	expression tag	UNP P04303
С	-8	SER		expression tag	UNP P04303
С	-7	GLY		expression tag	UNP P04303
С	-6	LEU		expression tag	UNP P04303
С	-5	VAL	-	expression tag	UNP P04303
С	-4	PRO	-	expression tag	UNP P04303
С	-3	ARG	_	expression tag	UNP P04303

Continued on next page...



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled  Modelled	Actual	Comment	Reference
С	-2	GLY	-	expression tag	UNP P04303
С	-1	SER	-	expression tag	UNP P04303
С	0	HIS	-	expression tag	UNP P04303
С	17	ASN	ASP	see remark 999	UNP P04303
С	110	PHE	LEU	engineered mutation	UNP P04303
С	163	PHE	TYR	see remark 999	UNP P04303
D	-19	MET	-	initiating methionine	UNP P04303
D	-18	GLY	-	expression tag	UNP P04303
D	-17	SER	-	expression tag	UNP P04303
D	-16	SER	-	expression tag	UNP P04303
D	-15	HIS	-	expression tag	UNP P04303
D	-14	HIS	-	expression tag	UNP P04303
D	-13	HIS	-	expression tag	UNP P04303
D	-12	HIS	-	expression tag	UNP P04303
D	-11	HIS	-	expression tag	UNP P04303
D	-10	HIS	-	expression tag	UNP P04303
D	-9	SER	-	expression tag	UNP P04303
D	-8	SER	-	expression tag	UNP P04303
D	-7	GLY	-	expression tag	UNP P04303
D	-6	LEU	-	expression tag	UNP P04303
D	-5	VAL	-	expression tag	UNP P04303
D	-4	PRO	_	expression tag	UNP P04303
D	-3	ARG	-	expression tag	UNP P04303
D	-2	GLY	-	expression tag	UNP P04303
D	-1	SER	-	expression tag	UNP P04303
D	0	HIS		expression tag	UNP P04303
D	17	ASN	ASP	see remark 999	UNP P04303
D	110	PHE	LEU	engineered mutation	UNP P04303
D	163	PHE	TYR	see remark 999	UNP P04303

• Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0
2	В	1	Total C O 6 3 3	0	0
2	С	1	Total C O 6 3 3	0	0
2	D	1	Total C O 6 3 3	0	0

• Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	В	1	Total Cl 1 1	0	0
3	A	1	Total Cl 1 1	0	0
3	D	1	Total Cl 1 1	0	0
3	С	1	Total Cl 1 1	0	0

• Molecule 4 is water.

$\mathbf{Mol}$	Chain	Residues	${f Atoms}$	ZeroOcc	AltConf
4	A	38	Total O 38 38	0	0

Continued on next page...



 $Continued\ from\ previous\ page...$ 


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	В	60	Total O 60 60	0	0
4	С	49	Total O 49 49	0	0
4	D	28	Total O 28 28	0	0



### 3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Uracil-DNA glycosylase





## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	92.35Å 102.92Å 116.59Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.08 - 2.40	Depositor
resolution (A)	47.08 - 2.40	EDS
% Data completeness	99.9 (47.08-2.40)	Depositor
(in resolution range)	99.9 (47.08-2.40)	EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.48 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
P. P.	0.228 , $0.253$	Depositor
$R, R_{free}$	0.232 , $0.256$	DCC
$R_{free}$ test set	2223 reflections $(5.04\%)$	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	44.4	Xtriage
Anisotropy	0.039	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.34, 27.4	EDS
L-test for twinning <sup>2</sup>	$ < L > = 0.49, < L^2> = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	7207	wwPDB-VP
Average B, all atoms $(Å^2)$	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.72% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $< L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Boı	nd lengths	Во	ond angles
MIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	A	0.58	1/1814 (0.1%)	0.74	$4/2468 \; (0.2\%)$
1	В	0.58	0/1803	0.72	0/2451
1	С	0.55	1/1803 (0.1%)	0.71	$2/2451 \ (0.1\%)$
1	D	0.56	0/1781	0.75	$4/2422 \ (0.2\%)$
All	All	0.57	$2/7201 \ (0.0\%)$	0.73	$10/9792 \ (0.1\%)$

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	${f Observed(\AA)}$	$\operatorname{Ideal}( ext{\AA})$
1	A	36	TRP	CB-CG	-6.14	1.39	1.50
1	С	36	TRP	CB-CG	5.28	1.59	1.50

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
1	A	193	ARG	NE-CZ-NH1	7.53	124.06	120.30
1	D	37	LEU	CA-CB-CG	7.19	131.84	115.30
1	D	39	ARG	NE-CZ-NH1	6.00	123.30	120.30
1	С	58	ARG	NE-CZ-NH1	5.99	123.30	120.30
1	D	187	ARG	NE-CZ-NH1	5.87	123.23	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within



.i	: 1	1.	C	$\alpha_1$	1: 4		1 4 1	1 1.
the asymmetric	unit.	wnereas	5vmm-	Ciasnes	IISUS S	vmmetrv	related	ciasnes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1764	0	1746	4	0
1	В	1754	0	1741	6	0
1	С	1754	0	1741	5	0
1	D	1732	0	1712	9	0
2	A	6	0	8	0	0
2	В	6	0	8	0	0
2	С	6	0	8	0	0
2	D	6	0	8	0	0
3	A	1	0	0	0	0
3	В	1	0	0	0	0
3	С	1	0	0	0	0
3	D	1	0	0	0	0
4	A	38	0	0	0	0
4	В	60	0	0	0	0
4	С	49	0	0	1	0
4	D	28	0	0	1	0
All	All	7207	0	6972	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 20 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{aligned}  ext{Interatomic} \  ext{distance} & ( ext{Å}) \end{aligned}$	$egin{aligned}  ext{Clash} \  ext{overlap } ( ext{Å}) \end{aligned}$
1:C:204:LEU:HD21	1:D:197:ILE:HD13	1.66	0.75
1:B:6:VAL:HG21	1:B:30:TYR:HD2	1.65	0.61
1:C:204:LEU:CD2	1:D:197:ILE:HD13	2.31	0.59
1:C:204:LEU:HD21	1:D:197:ILE:HG21	1.91	0.53
1:D:22:VAL:CG2	1:D:143:LEU:HG	2.41	0.51

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Analysed Favoured Allowed		Outliers	Perce	ntiles
1	A	216/238 (91%)	210 (97%)	5 (2%)	1 (0%)	29	41
1	В	$212/238 \ (89\%)$	206 (97%)	5 (2%)	1 (0%)	29	41
1	С	212/238 (89%)	206 (97%)	5 (2%)	1 (0%)	29	41
1	D	209/238~(88%)	203 (97%)	5 (2%)	1 (0%)	29	41
All	All	849/952 (89%)	825 (97%)	20 (2%)	4 (0%)	29	41

#### All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	138	ASP
1	С	138	ASP
1	D	138	ASP
1	В	138	ASP

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	Percentiles		
1	A	197/216 (91%)	197 (100%)	0	100	100		
1	В	197/216 (91%)	197 (100%)	0	100	100		
1	С	197/216 (91%)	197 (100%)	0	100	100		
1	D	$195/216 \ (90\%)$	195 (100%)	0	100	100		
All	All	786/864 (91%)	786 (100%)	0	100	100		

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

$\mathbf{Mol}$	Chain	${f Res}$	Type
1	A	206	ASN



#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry (i)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Res	Link	Bond lengths			Bond angles			
10101	Type	Chain	nes	Lilik	TIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GOL	В	301	_	5,5,5	0.48	0	5,5,5	0.52	0	
2	GOL	A	301	_	5,5,5	0.46	0	5,5,5	0.67	0	
2	GOL	D	301	_	5,5,5	0.29	0	5,5,5	0.31	0	
2	GOL	С	301	_	5,5,5	0.63	0	5,5,5	0.39	0	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	В	301	-	-	3/4/4/4	_
2	GOL	A	301	_	-	0/4/4/4	-
2	GOL	D	301	_	-	4/4/4/4	_
2	GOL	С	301	_	-	2/4/4/4	-

There are no bond length outliers.



There are no bond angle outliers.

There are no chirality outliers.

5 of 9 torsion outliers are listed below:

Mol	Chain	${f Res}$	Type	Atoms
2	В	301	GOL	C1-C2-C3-O3
2	С	301	GOL	C1-C2-C3-O3
2	D	301	GOL	O1-C1-C2-O2
2	В	301	GOL	O2-C2-C3-O3
2	D	301	GOL	O1-C1-C2-C3

There are no ring outliers.

No monomer is involved in short contacts.

#### 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



### 6 Fit of model and data (i)

#### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$	$OWAB(Å^2)$	Q < 0.9
1	A	218/238 (91%)	0.02	3 (1%) 75 73	28, 47, 77, 100	0
1	В	216/238 (90%)	0.02	2 (0%) 84 82	28, 45, 73, 108	0
1	С	216/238 (90%)	0.03	6 (2%) 53 51	29, 47, 72, 98	0
1	D	213/238 (89%)	0.12	2 (0%) 84 82	29, 50, 75, 100	0
All	All	863/952 (90%)	0.05	13 (1%) 73 72	28, 48, 75, 108	0

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	168	ALA	3.7
1	В	165	ASN	3.4
1	С	1	MET	3.2
1	A	103	ILE	2.7
1	A	36	TRP	2.5

#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

#### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathring{\mathbf{A}}^2)$	Q < 0.9
2	GOL	D	301	6/6	0.75	0.38	57,77,88,91	0
2	GOL	С	301	6/6	0.77	0.24	56,63,68,71	0
2	GOL	В	301	6/6	0.91	0.18	44,51,57,60	0
2	GOL	A	301	6/6	0.92	0.19	44,50,52,52	0
3	CL	A	302	1/1	0.95	0.10	52,52,52,52	0
3	CL	С	302	1/1	0.96	0.09	51,51,51,51	0
3	CL	D	302	1/1	0.96	0.14	57,57,57,57	0
3	CL	В	302	1/1	0.98	0.15	43,43,43,43	0

## 6.5 Other polymers (i)

There are no such residues in this entry.

