

wwPDB X-ray Structure Validation Summary Report (i)

Oct 9, 2023 – 02:20 PM EDT

PDB ID : 7JXR

Title : Crystal Structure Human Immunodeficiency Virus-1 Matrix protein Mutant

Q63R Crystal Form 1

Authors: Green, T.J.; Eastep, G.N.; Ghanam, R.H.; Saad, J.S.

Deposited on : 2020-08-27

Resolution : 2.04 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35.1 buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\ DIFFRACTION$

The reported resolution of this entry is 2.04 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	Similar resolution $(\# \text{Entries, resolution range}(\text{\AA}))$
R_{free}	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain			
1	A	137	% 6 6%	10%		24%
1	В	137	76%		5%	19%
1	С	137	67%	10%		23%
1	D	137	67%	9%		24%
1	Е	137	% 6 9%	7%	•	23%



Continued from previous page...

Mol	Chain	Length				
			% ■			
1	F	137	66%	9%	٠	23%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SO4	A	203	-	-	X	-
3	SO4	В	203	-	-	X	-
3	SO4	С	203	-	-	X	-
3	SO4	D	205	-	X	-	-
3	SO4	F	204	-	-	X	-



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 5507 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Matrix protein.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	A	104	Total	С	N	О	S	0	0	0
1	A	104	842	529	153	158	2	0	U	U
1	В	111	Total	С	N	О	S	0	1	0
1	Ъ	111	904	568	166	168	2	0	1	0
1	С	106	Total	С	N	О	S	0	0	0
1		100	857	538	156	161	2	0	U	
1	D	104	Total	С	N	О	S	0	0	0
1	D	104	842	529	153	158	2	0		
1	Е	105	Total	С	N	О	S	0	0	0
1	l L	105	848	532	154	160	2	0		0
1	F	105	Total	С	N	О	S	0	0	0
1	Г	100	848	532	154	160	2	U	U	U

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	63	ARG	GLN	engineered mutation	UNP Q6E183
A	133	HIS	-	expression tag	UNP Q6E183
A	134	HIS	-	expression tag	UNP Q6E183
A	135	HIS	-	expression tag	UNP Q6E183
A	136	HIS	-	expression tag	UNP Q6E183
A	137	HIS	-	expression tag	UNP Q6E183
A	138	HIS	-	expression tag	UNP Q6E183
В	63	ARG	GLN	engineered mutation	UNP Q6E183
В	133	HIS	-	expression tag	UNP Q6E183
В	134	HIS	-	expression tag	UNP Q6E183
В	135	HIS	-	expression tag	UNP Q6E183
В	136	HIS	-	expression tag	UNP Q6E183
В	137	HIS	-	expression tag	UNP Q6E183
В	138	HIS	-	expression tag	UNP Q6E183
С	63	ARG	GLN	engineered mutation	UNP Q6E183
С	133	HIS	-	expression tag	UNP Q6E183
С	134	HIS	-	expression tag	UNP Q6E183



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
С	135	HIS	-	expression tag	UNP Q6E183
С	136	HIS	-	expression tag	UNP Q6E183
С	137	HIS	-	expression tag	UNP Q6E183
С	138	HIS	-	expression tag	UNP Q6E183
D	63	ARG	GLN	engineered mutation	UNP Q6E183
D	133	HIS	-	expression tag	UNP Q6E183
D	134	HIS	-	expression tag	UNP Q6E183
D	135	HIS	-	expression tag	UNP Q6E183
D	136	HIS	-	expression tag	UNP Q6E183
D	137	HIS	-	expression tag	UNP Q6E183
D	138	HIS	-	expression tag	UNP Q6E183
Е	63	ARG	GLN	engineered mutation	UNP Q6E183
Е	133	HIS	-	expression tag	UNP Q6E183
Е	134	HIS	-	expression tag	UNP Q6E183
Е	135	HIS	-	expression tag	UNP Q6E183
Е	136	HIS	-	expression tag	UNP Q6E183
Е	137	HIS	-	expression tag	UNP Q6E183
Е	138	HIS	-	expression tag	UNP Q6E183
F	63	ARG	GLN	engineered mutation	UNP Q6E183
F	133	HIS	-	expression tag	UNP Q6E183
F	134	HIS	-	expression tag	UNP Q6E183
F	135	HIS	-	expression tag	UNP Q6E183
F	136	HIS	-	expression tag	UNP Q6E183
F	137	HIS	-	expression tag	UNP Q6E183
F	138	HIS	-	expression tag	UNP Q6E183


 \bullet Molecule 2 is HEXAETHYLENE GLYCOL (three-letter code: P6G) (formula: $\mathrm{C_{12}H_{26}O_{7}}).$





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 12 8 4	0	0
2	В	1	Total C O 16 10 6	0	0
2	С	1	Total C O 12 8 4	0	0
2	D	1	Total C O 16 10 6	0	0
2	Е	1	Total C O 12 8 4	0	0
2	F	1	Total C O 11 7 4	0	0

 \bullet Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: $\mathrm{O_4S}).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S	0	0
	11	1	5 4 1 Total O S		0
3	В	1	5 4 1	0	0
3	В	1	Total O S 5 4 1	0	0
3	С	1	Total O S	0	0
			5 4 1 Total O S		
3	С	1	$\begin{bmatrix} 10001 & 0 & 0 \\ 5 & 4 & 1 \end{bmatrix}$	0	0



 $Continued\ from\ previous\ page...$

Mol	Chain	Residues	Ato	oms		ZeroOcc	AltConf
3	D	1	Total	О	S	0	0
3	D	1	5	4	1	0	U
3	D	1	Total	О	S	0	0
J	D	1	5	4	1	U	U
3	D	1	Total	О	S	0	0
3	D	1	5	4	1	0	U
3	D	1	Total	О	S	0	0
J	D	1	5	4	1	U	U
3	E	1	Total	Ο	\mathbf{S}	0	0
	Ľ	1	5	4	1	U	U
3	F	1	Total	Ο	S	0	0
	I.	1	5	4	1	U	U
3	F	1	Total	О	S	0	0
	I'	1	5	4	1		U
3	F	1	Total	О	S	0	0
3	I'	1	5	4	1		U

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	30	Total O 30 30	0	0
4	В	33	Total O 33 33	0	0
4	С	40	Total O 40 40	0	0
4	D	41	Total O 41 41	0	0
4	Е	30	Total O 30 30	0	0
4	F	43	Total O 43 43	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Matrix protein 24% HIS HIS HIS HIS • Molecule 1: Matrix protein Chain B: 76% 5% 19% • Molecule 1: Matrix protein Chain C: 67% 10% 23% • Molecule 1: Matrix protein Chain D: 67% 9% 24% • Molecule 1: Matrix protein Chain E: 69% 23%







4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	63.14Å 97.92Å 84.82Å	Depositor
a, b, c, α , β , γ	90.00° 99.94° 90.00°	Depositor
Resolution (Å)	46.21 - 2.04	Depositor
rtesolution (A)	46.21 - 2.04	EDS
% Data completeness	94.7 (46.21-2.04)	Depositor
(in resolution range)	92.2 (46.21-2.04)	EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.78 (at 2.03Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874	Depositor
P. P.	0.203 , 0.234	Depositor
R, R_{free}	0.203 , 0.234	DCC
R_{free} test set	2000 reflections (3.24%)	wwPDB-VP
Wilson B-factor (Å ²)	41.4	Xtriage
Anisotropy	0.116	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.36, 54.1	EDS
L-test for twinning ²	$ < L > = 0.49, < L^2> = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5507	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 10.71% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: P6G, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.36	0/854	0.57	0/1146	
1	В	0.38	0/919	0.52	0/1229	
1	С	0.40	0/869	0.57	0/1165	
1	D	0.45	0/854	0.55	0/1146	
1	Е	0.37	0/860	0.52	0/1154	
1	F	0.40	0/860	0.57	0/1154	
All	All	0.40	0/5216	0.55	0/6994	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	842	0	864	11	0
1	В	904	0	942	7	0
1	С	857	0	882	15	0
1	D	842	0	864	9	0
1	Е	848	0	869	9	0
1	F	848	0	869	12	0
2	A	12	0	15	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	В	16	0	21	1	0
2	С	12	0	12	0	0
2	D	16	0	21	0	0
2	Е	12	0	15	1	0
2	F	11	0	13	0	0
3	A	10	0	0	3	0
3	В	10	0	0	2	0
3	С	10	0	0	6	0
3	D	20	0	0	2	0
3	Е	5	0	0	0	0
3	F	15	0	0	3	0
4	A	30	0	0	2	0
4	В	33	0	0	1	0
4	С	40	0	0	1	0
4	D	41	0	0	6	0
4	Е	30	0	0	2	0
4	F	43	0	0	2	0
All	All	5507	0	5387	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 62 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{aligned} & ext{Interatomic} \ & ext{distance} \ & ext{(Å)} \end{aligned}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:E:58:ARG:HH22	1:E:104:ILE:HG23	1.49	0.76
3:D:205:SO4:S	4:D:328:HOH:O	2.45	0.75
1:D:7:VAL:N	4:D:302:HOH:O	2.22	0.71
1:C:40:GLU:OE1	1:C:43:ARG:NH2	2.22	0.71
1:D:58:ARG:NH1	4:D:304:HOH:O	2.24	0.71

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the r	number of residu	ues for which	the backbone	conformation	was
analysed, and the total number of	residues.				

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	102/137~(74%)	102 (100%)	0	0	100	100
1	В	110/137~(80%)	110 (100%)	0	0	100	100
1	\mathbf{C}	$104/137\ (76\%)$	104 (100%)	0	0	100	100
1	D	102/137~(74%)	102 (100%)	0	0	100	100
1	${ m E}$	103/137~(75%)	103 (100%)	0	0	100	100
1	F	103/137~(75%)	103 (100%)	0	0	100	100
All	All	624/822 (76%)	624 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	A	92/117 (79%)	90 (98%)	2 (2%)	52	46	
1	В	99/117 (85%)	99 (100%)	0	100	100	
1	С	94/117 (80%)	94 (100%)	0	100	100	
1	D	92/117 (79%)	91 (99%)	1 (1%)	73	73	
1	E	93/117 (80%)	92 (99%)	1 (1%)	73	73	
1	F	93/117 (80%)	90 (97%)	3 (3%)	39	32	
All	All	563/702 (80%)	556 (99%)	7 (1%)	71	70	

5 of 7 residues with a non-rotameric sidechain are listed below:

	Mol	Chain	Res	Type
Ī	1	Ε	30	LYS
	1	F	43	ARG
	1	F	110	LYS
	1	F	65	GLN
	1	D	43	ARG



Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

20 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

N / - 1	Т	Clasia	Das	T : 1-	Во	ond leng	ths	В	ond ang	gles
Mol	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	D	205	-	4,4,4	0.08	0	6,6,6	5.31	6 (100%)
3	SO4	Е	202	-	4,4,4	0.12	0	6,6,6	0.16	0
3	SO4	С	203	-	4,4,4	0.09	0	6,6,6	0.16	0
3	SO4	D	204	-	4,4,4	0.14	0	6,6,6	0.09	0
3	SO4	F	203	-	4,4,4	0.16	0	6,6,6	0.07	0
3	SO4	В	203	-	4,4,4	0.14	0	6,6,6	0.13	0
2	P6G	A	201	-	11,11,18	0.55	0	10,10,17	0.36	0
2	P6G	С	202	-	11,11,18	0.48	0	10,10,17	0.39	0
2	P6G	В	201	-	15,15,18	0.50	0	14,14,17	0.34	0
3	SO4	D	203	-	4,4,4	0.18	0	6,6,6	0.23	0
2	P6G	F	201	-	10,10,18	0.51	0	9,9,17	0.48	0
3	SO4	D	202	-	4,4,4	0.17	0	6,6,6	0.28	0
2	P6G	D	201	-	15,15,18	0.53	0	14,14,17	0.52	0
3	SO4	A	203	-	4,4,4	0.16	0	6,6,6	0.39	0



Mol	Mol Type		Res	Link	Во	Bond lengths			Bond angles		
MIOI	туре	Chain	nes	nes	es Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	P6G	Е	201	-	11,11,18	0.54	0	10,10,17	0.26	0	
3	SO4	F	204	-	4,4,4	0.22	0	6,6,6	0.35	0	
3	SO4	A	202	-	4,4,4	0.15	0	6,6,6	0.25	0	
3	SO4	С	201	-	4,4,4	0.17	0	6,6,6	0.47	0	
3	SO4	F	202	-	4,4,4	0.15	0	6,6,6	0.17	0	
3	SO4	В	202	-	4,4,4	0.16	0	6,6,6	0.36	0	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	P6G	Е	201	-	-	4/9/9/16	-
2	P6G	В	201	-	-	9/13/13/16	-
2	P6G	F	201	-	-	6/8/8/16	-
2	P6G	D	201	-	-	7/13/13/16	-
2	P6G	A	201	-	-	6/9/9/16	-
2	P6G	С	202	-	-	7/9/9/16	-

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$Ideal(^{o})$
3	D	205	SO4	O4-S-O3	-7.33	77.78	109.06
3	D	205	SO4	O3-S-O1	-6.76	74.04	109.31
3	D	205	SO4	O4-S-O1	5.57	138.36	109.31
3	D	205	SO4	O4-S-O2	-4.33	86.72	109.31
3	D	205	SO4	O2-S-O1	-3.32	84.90	109.43

There are no chirality outliers.

5 of 39 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	201	P6G	C15-C14-O13-C12
2	A	201	P6G	O13-C14-C15-O16
2	F	201	P6G	O7-C8-C9-O10
2	С	202	P6G	O10-C11-C12-O13
2	С	202	P6G	O7-C8-C9-O10




There are no ring outliers.


9 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	205	SO4	1	0
3	С	203	SO4	6	0
3	F	203	SO4	1	0
3	В	203	SO4	2	0
2	В	201	P6G	1	0
3	D	203	SO4	1	0
3	A	203	SO4	3	0
2	Е	201	P6G	1	0
3	F	204	SO4	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$	$\mathbf{OWAB}(\mathbf{\mathring{A}}^2)$	Q<0.9
1	A	104/137~(75%)	0.29	2 (1%) 66 71	36, 51, 70, 108	0
1	В	111/137 (81%)	0.39	4 (3%) 42 46	36, 50, 93, 119	0
1	С	106/137 (77%)	0.33	2 (1%) 66 71	33, 49, 72, 107	0
1	D	104/137 (75%)	0.24	0 100 100	31, 46, 59, 85	0
1	E	105/137 (76%)	0.28	2 (1%) 66 71	38, 53, 87, 108	0
1	F	105/137 (76%)	0.30	1 (0%) 82 84	33, 45, 64, 99	0
All	All	635/822 (77%)	0.31	11 (1%) 70 73	31, 48, 75, 119	0

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	Е	111	SER	3.8
1	В	114	LYS	3.3
1	В	91	ARG	2.7
1	С	26	LYS	2.6
1	В	43	ARG	2.6

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}({ m \AA}^2)$	Q < 0.9
3	SO4	F	203	5/5	0.79	0.26	59,60,64,67	5
3	SO4	D	204	5/5	0.80	0.26	63,63,68,73	5
2	P6G	A	201	12/19	0.81	0.16	51,56,62,68	0
2	P6G	В	201	16/19	0.82	0.17	52,60,66,67	0
2	P6G	D	201	16/19	0.83	0.14	47,54,60,60	0
3	SO4	С	203	5/5	0.84	0.26	80,85,86,98	0
2	P6G	С	202	12/19	0.86	0.16	48,51,57,61	0
3	SO4	F	204	5/5	0.86	0.34	49,55,63,77	5
3	SO4	D	203	5/5	0.88	0.20	54,56,63,64	5
2	P6G	F	201	11/19	0.89	0.15	39,52,57,57	0
2	P6G	Е	201	12/19	0.89	0.20	53,58,59,63	0
3	SO4	В	203	5/5	0.91	0.15	54,58,64,70	5
3	SO4	D	205	5/5	0.94	0.81	63,64,69,72	0
3	SO4	В	202	5/5	0.95	0.14	60,61,64,69	0
3	SO4	A	203	5/5	0.96	0.47	52,56,64,65	0
3	SO4	F	202	5/5	0.97	0.12	60,64,65,66	0
3	SO4	С	201	5/5	0.98	0.18	64,64,66,69	0
3	SO4	A	202	5/5	0.98	0.16	51,52,55,55	0
3	SO4	D	202	5/5	0.98	0.18	53,55,58,63	0
3	SO4	E	202	5/5	0.99	0.13	61,63,67,67	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.














Electron density around P6G D 201:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around P6G C 202:


 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)













6.5 Other polymers (i)

There are no such residues in this entry.

