



wwPDB EM Validation Summary Report ⓘ

Oct 22, 2024 – 12:26 AM JST

PDB ID : 8JXO
EMDB ID : EMD-36707
Title : Cryo-EM structure of BICHR2 class one
Authors : Zhang, M.F.
Deposited on : 2023-07-01
Resolution : 2.70 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : **FAILED**
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4.02b-467
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : **FAILED**
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	391	42% 19% 6% • 33%
1	D	391	43% 18% 5% • 33%
1	E	391	42% 19% 5% • 33%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	RET	A	401	-	-	X	-
2	RET	D	401	-	-	X	-
2	RET	E	401	-	-	X	-

2 Entry composition i


There are 4 unique types of molecules in this entry. The entry contains 6294 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BICHR2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	262	Total 2069	C 1372	N 318	O 365	S 14	0	0
1	D	262	Total 2069	C 1372	N 318	O 365	S 14	0	0
1	E	262	Total 2069	C 1372	N 318	O 365	S 14	0	0

- Molecule 2 is RETINAL (three-letter code: RET) (formula: $C_{20}H_{28}O$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms		AltConf
2	A	1	Total 20	C 20	0
2	D	1	Total 20	C 20	0
2	E	1	Total 20	C 20	0

- Molecule 3 is POTASSIUM ION (three-letter code: K) (formula: K) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	AltConf
3	A	6	Total K 6 6	0
3	D	6	Total K 6 6	0
3	E	6	Total K 6 6	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	AltConf
4	A	3	Total O 3 3	0
4	D	3	Total O 3 3	0
4	E	3	Total O 3 3	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: BICHR2



- Molecule 1: BICHR2



- Molecule 1: BICHR2



ILE	ARG	F189	P110	MET
GLN	ALA	F200	L111	S2
LYS	SER	F201	L112	E3
GLU	SER	P202	V113	C4
VAL	GLY	L203	L114	C5
ASP	PHE	L204	D115	E6
SER	PHE	L207	F116	L7
ARG	GLN	H210	C117	C10
SER	GLN	I215	L122	E13
LYS	GLY	I220	L126	T27
LYS	ILE	Y220	S129	
MET	PRO	A225	S130	
GLN	ASP	I226	S131	
ARG	ASP	L227	I132	
GLU	GLN	I228	A133	M38
THR	THR	L229	L136	K41
ALA	VAL	L230	A137	I42
ARG	SER	C231	I138	A43
PHE	SER	K232	A139	
ALA	VAL	A233	V140	H53
ILE	GLY	F237	S141	
ILE	GLY	L244	T142	E56
GLN	GLY	ARG	F143	E61
ASP	GLY	ALA	I144	
ASP	ARG	ARG	V145	N62
GLU	ARG	ARG	D146	W63
ASP	ARG	ARG	A147	L64
GLY	GLN	ARG	P148	S65
GLY	GLY	HIS	Y149	P66
ARG	MET	E255	R150	G67
GLY	LEU	A256	M153	I68
ALA	TYR	E257	L158	L71
ALA	TYR	K260	F161	C72
GLY	LEU	L261	L168	G73
ALA	ALA	E262	E171	F78
ARG	ALA	G263	I172	I81
ARG	SER	ASP	Q175	Y86
ARG	TYR	LYS	R176	A94
ARG	LEU	GLU	E177	F95
LYS	LEU	ALA	S182	S96
ASN	LYS	ASN	L187	L97
ALA	ASP	ASP	L191	V100
ARG	PHO	THR	L192	W101
ARG	ALA	GLY	L193	Y102
ARG	ALA	LEU	R194	S103
ASN	GLU	LYS	L195	L106
ALA	GLY	ILE	ARG	I107
ARG	LEU	ARG	HIS	T108
ARG	LEU	ARG	TYR	I109
ARG	LEU	ARG	ASN	C109
ARG	LYS	ARG		
GLY	LYS	ASN		

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	47000	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	1200	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	FEI FALCON IV (4k x 4k)	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: RET, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.41	0/2135	0.58	3/2915 (0.1%)
1	D	0.41	0/2135	0.58	3/2915 (0.1%)
1	E	0.41	0/2135	0.58	3/2915 (0.1%)
All	All	0.41	0/6405	0.58	9/8745 (0.1%)

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	D	110	PRO	CA-N-CD	-6.53	102.36	111.50
1	E	110	PRO	CA-N-CD	-6.52	102.37	111.50
1	A	110	PRO	CA-N-CD	-6.52	102.38	111.50
1	A	202	PRO	CA-N-CD	-5.94	103.19	111.50
1	E	202	PRO	CA-N-CD	-5.94	103.19	111.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2069	0	2016	89	0
1	D	2069	0	2016	86	0
1	E	2069	0	2016	89	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	20	0	27	21	0
2	D	20	0	27	21	0
2	E	20	0	27	22	0
3	A	6	0	0	1	0
3	D	6	0	0	1	0
3	E	6	0	0	1	0
4	A	3	0	0	1	0
4	D	3	0	0	1	0
4	E	3	0	0	1	0
All	All	6294	0	6129	221	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 221 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:81:ILE:HG21	1:D:153:MET:HE3	1.35	1.07
1:E:136:LEU:HD21	2:E:401:RET:H10	1.37	1.06
1:A:94:ALA:CB	1:D:144:ILE:HG22	1.87	1.05
1:D:94:ALA:CB	1:E:144:ILE:HG22	1.87	1.05
1:D:136:LEU:HD21	2:D:401:RET:H10	1.37	1.05

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	260/391 (66%)	237 (91%)	18 (7%)	5 (2%)	6 17
1	D	260/391 (66%)	237 (91%)	18 (7%)	5 (2%)	6 17

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	260/391 (66%)	237 (91%)	18 (7%)	5 (2%)	6	17
All	All	780/1173 (66%)	711 (91%)	54 (7%)	15 (2%)	9	17

5 of 15 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	232	LYS
1	D	232	LYS
1	E	232	LYS
1	A	53	HIS
1	A	147	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	213/319 (67%)	161 (76%)	52 (24%)	0	1
1	D	213/319 (67%)	161 (76%)	52 (24%)	0	1
1	E	213/319 (67%)	161 (76%)	52 (24%)	0	1
All	All	639/957 (67%)	483 (76%)	156 (24%)	2	1

5 of 156 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	65	SER
1	E	192	LEU
1	E	100	VAL
1	E	141	SER
1	E	244	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	99	GLN
1	D	99	GLN
1	E	99	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 21 ligands modelled in this entry, 18 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	RET	A	401	1	20,20,21	0.73	0	27,27,28	1.98	8 (29%)
2	RET	E	401	1	20,20,21	0.73	0	27,27,28	1.98	8 (29%)
2	RET	D	401	1	20,20,21	0.73	0	27,27,28	1.98	8 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	RET	A	401	1	-	4/13/30/31	0/1/1/1
2	RET	E	401	1	-	4/13/30/31	0/1/1/1
2	RET	D	401	1	-	4/13/30/31	0/1/1/1

There are no bond length outliers.

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	401	RET	C11-C10-C9	-5.05	120.11	127.31
2	A	401	RET	C11-C10-C9	-5.04	120.11	127.31
2	E	401	RET	C11-C10-C9	-5.04	120.12	127.31
2	D	401	RET	C7-C8-C9	-4.76	119.05	126.23
2	A	401	RET	C7-C8-C9	-4.75	119.05	126.23

There are no chirality outliers.

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	RET	C7-C8-C9-C19
2	D	401	RET	C7-C8-C9-C19
2	E	401	RET	C7-C8-C9-C19
2	A	401	RET	C7-C8-C9-C10
2	D	401	RET	C7-C8-C9-C10


There are no ring outliers.


3 monomers are involved in 64 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	RET	21	0
2	E	401	RET	22	0
2	D	401	RET	21	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.