



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 20, 2023 – 04:44 AM EDT

PDB ID : 5JZJ  
Title : Crystal structure of DCLK1-KD in complex with AMPPN  
Authors : Patel, O.; Lucet, I.  
Deposited on : 2016-05-17  
Resolution : 1.71 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 1.71 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5722 (1.74-1.70)
Clashscore	141614	6152 (1.74-1.70)
Ramachandran outliers	138981	6051 (1.74-1.70)
Sidechain outliers	138945	6051 (1.74-1.70)
RSRZ outliers	127900	5629 (1.74-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	297	 2% 88% 9%
1	B	297	 2% 85% 12%

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4683 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein kinase DCLK1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	271	2111	1356	346	398	11	0	3	0
1	B	261	2018	1302	327	379	10	0	4	0

There are 38 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	353	MET	-	initiating methionine	UNP O15075
A	354	HIS	-	expression tag	UNP O15075
A	355	HIS	-	expression tag	UNP O15075
A	356	HIS	-	expression tag	UNP O15075
A	357	HIS	-	expression tag	UNP O15075
A	358	HIS	-	expression tag	UNP O15075
A	359	HIS	-	expression tag	UNP O15075
A	360	HIS	-	expression tag	UNP O15075
A	361	HIS	-	expression tag	UNP O15075
A	362	GLY	-	expression tag	UNP O15075
A	363	SER	-	expression tag	UNP O15075
A	364	GLU	-	expression tag	UNP O15075
A	365	ASN	-	expression tag	UNP O15075
A	366	LEU	-	expression tag	UNP O15075
A	367	TYR	-	expression tag	UNP O15075
A	368	PHE	-	expression tag	UNP O15075
A	369	GLN	-	expression tag	UNP O15075
A	370	GLY	-	expression tag	UNP O15075
A	371	SER	-	expression tag	UNP O15075
B	353	MET	-	initiating methionine	UNP O15075
B	354	HIS	-	expression tag	UNP O15075
B	355	HIS	-	expression tag	UNP O15075
B	356	HIS	-	expression tag	UNP O15075
B	357	HIS	-	expression tag	UNP O15075
B	358	HIS	-	expression tag	UNP O15075

*Continued on next page...*


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	359	HIS	-	expression tag	UNP O15075
B	360	HIS	-	expression tag	UNP O15075
B	361	HIS	-	expression tag	UNP O15075
B	362	GLY	-	expression tag	UNP O15075
B	363	SER	-	expression tag	UNP O15075
B	364	GLU	-	expression tag	UNP O15075
B	365	ASN	-	expression tag	UNP O15075
B	366	LEU	-	expression tag	UNP O15075
B	367	TYR	-	expression tag	UNP O15075
B	368	PHE	-	expression tag	UNP O15075
B	369	GLN	-	expression tag	UNP O15075
B	370	GLY	-	expression tag	UNP O15075
B	371	SER	-	expression tag	UNP O15075

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Mg 1 1	0	0
2	B	1	Total Mg 1 1	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0

- Molecule 4 is AMP PHOSPHORAMIDATE (three-letter code: AN2) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>6</sub>O<sub>9</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C N O P 27 10 6 9 2	0	0
4	B	1	Total C N O P 27 10 6 9 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	271	Total O 271 271	0	0
5	B	222	Total O 222 222	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Serine/threonine-protein kinase DCLK1

Chain A: 



- Molecule 1: Serine/threonine-protein kinase DCLK1

Chain B: 



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	56.39Å 81.27Å 59.61Å 90.00° 92.10° 90.00°	Depositor
Resolution (Å)	56.35 – 1.71 56.35 – 1.71	Depositor EDS
% Data completeness (in resolution range)	99.6 (56.35-1.71) 99.9 (56.35-1.71)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.88 (at 1.71Å)	Xtrriage
Refinement program	BUSTER-TNT 2.10.0	Depositor
R, $R_{free}$	0.177 , 0.208 0.178 , 0.210	Depositor DCC
$R_{free}$ test set	2932 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.1	Xtrriage
Anisotropy	0.392	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 47.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.027 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4683	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.56% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: AN2, SO4, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.50	0/2162	0.59	0/2939
1	B	0.50	0/2071	0.59	0/2824
All	All	0.50	0/4233	0.59	0/5763

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2111	0	2100	5	0
1	B	2018	0	1998	5	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	5	0	0	0	0
4	A	27	0	14	0	0
4	B	27	0	14	1	0
5	A	271	0	0	0	0
5	B	222	0	0	1	0
All	All	4683	0	4126	10	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including



hydrogen atoms). The all-atom clashscore for this structure is 1.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:396:ILE:HG21	1:A:406:GLU:HG3	1.88	0.54
1:A:476:ALA:O	1:A:480:THR:HG22	2.09	0.51
1:B:457:VAL:HG12	1:B:460:GLU:H	1.79	0.47
1:B:495:LEU:HD13	1:B:517:LEU:HD21	1.97	0.46
1:A:495:LEU:HD13	1:A:517:LEU:HD21	1.98	0.45
1:B:588:ARG:NH2	5:B:1002:HOH:O	2.46	0.45
1:B:460:GLU:HG3	1:B:462:TYR:CE2	2.52	0.44
1:A:437:VAL:CG2	1:A:461:LEU:HD21	2.48	0.42
1:A:480:THR:HG21	1:A:483:TYR:CE1	2.56	0.40
1:B:399:GLY:HA3	4:B:902:AN2:O2B	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	270/297 (91%)	265 (98%)	5 (2%)	0	100	100
1	B	261/297 (88%)	257 (98%)	4 (2%)	0	100	100
All	All	531/594 (89%)	522 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	230/259 (89%)	227 (99%)	3 (1%)	69	55
1	B	218/259 (84%)	214 (98%)	4 (2%)	59	41
All	All	448/518 (86%)	441 (98%)	7 (2%)	73	47

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	398[A]	ASP
1	A	398[B]	ASP
1	A	454	GLU
1	B	398[A]	ASP
1	B	398[B]	ASP
1	B	457	VAL
1	B	480	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	AN2	A	703	2	24,29,29	1.84	6 (25%)	25,45,45	1.55	3 (12%)
3	SO4	A	702	-	4,4,4	0.23	0	6,6,6	0.19	0
4	AN2	B	902	2	24,29,29	2.00	7 (29%)	25,45,45	1.85	3 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	AN2	A	703	2	-	1/9/32/32	0/3/3/3
4	AN2	B	902	2	-	1/9/32/32	0/3/3/3

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	703	AN2	C2-N3	4.68	1.39	1.32
4	B	902	AN2	O4'-C1'	4.35	1.47	1.41
4	A	703	AN2	PB-O3A	4.12	1.64	1.59
4	B	902	AN2	C2-N3	3.93	1.38	1.32
4	B	902	AN2	PB-O2B	-3.54	1.47	1.56
4	B	902	AN2	PB-O3A	3.45	1.63	1.59
4	B	902	AN2	C2-N1	3.24	1.39	1.33
4	A	703	AN2	PB-O2B	-3.19	1.48	1.56
4	A	703	AN2	C2-N1	3.06	1.39	1.33
4	B	902	AN2	C5-C4	-2.31	1.34	1.40
4	B	902	AN2	O3'-C3'	2.06	1.47	1.43
4	A	703	AN2	C5-C4	-2.06	1.35	1.40
4	A	703	AN2	C6-C5	-2.04	1.35	1.43

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	902	AN2	N3-C2-N1	-7.09	117.60	128.68
4	A	703	AN2	N3-C2-N1	-5.93	119.41	128.68
4	B	902	AN2	O2B-PB-O1B	3.93	120.45	110.25
4	A	703	AN2	O2B-PB-O1B	3.20	118.56	110.25

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	703	AN2	C4-C5-N7	-2.23	107.07	109.40
4	B	902	AN2	C4-C5-N7	-2.09	107.22	109.40

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	703	AN2	PB-O3A-PA-O1A
4	B	902	AN2	PB-O3A-PA-O2A


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	902	AN2	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	271/297 (91%)	-0.12	5 (1%) 68 72	10, 21, 44, 63	0
1	B	261/297 (87%)	-0.10	5 (1%) 66 70	12, 23, 43, 52	0
All	All	532/594 (89%)	-0.11	10 (1%) 66 70	10, 23, 44, 63	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	375	VAL	3.4
1	B	401	PHE	3.1
1	B	481	ASN	2.7
1	B	400	ASN	2.7
1	A	427	ARG	2.6
1	A	590	SER	2.4
1	A	426	CYS	2.3
1	B	385	THR	2.2
1	A	396	ILE	2.1
1	B	596	VAL	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	AN2	A	703	27/27	0.98	0.08	15,17,20,26	0
4	AN2	B	902	27/27	0.98	0.07	14,16,21,27	0
2	MG	A	701	1/1	0.99	0.03	16,16,16,16	0
2	MG	B	901	1/1	1.00	0.05	16,16,16,16	0
3	SO4	A	702	5/5	1.00	0.06	13,13,15,16	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.