



Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 03:53 am BST

PDB ID : 4K2U
Title : Crystal structure of PfEBA-175 F1 in complex with R218 antibody Fab fragment
Authors : Tolia, N.H.
Deposited on : 2013-04-09
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1544 (2.48-2.44)
Clashscore	141614	1613 (2.48-2.44)
Ramachandran outliers	138981	1598 (2.48-2.44)
Sidechain outliers	138945	1598 (2.48-2.44)
RSRZ outliers	127900	1523 (2.48-2.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	297	
1	B	297	
2	H	233	
2	I	233	
3	L	234	
3	M	234	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	SO4	I	301	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 20975 atoms, of which 10201 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Erythrocyte binding antigen 175.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	230	Total 3869	C 1236	H 1919	N 345	O 353	S 16	0	0	0
1	B	257	Total 4313	C 1376	H 2144	N 376	O 398	S 19	0	0	0

- Molecule 2 is a protein called Antibody Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
2	H	208	Total 3143	C 1023	H 1544	N 257	O 314	S 5	0	0	0
2	I	214	Total 3235	C 1049	H 1589	N 267	O 324	S 6	0	0	0

- Molecule 3 is a protein called Antibody Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
3	L	208	Total 3146	C 1009	H 1526	N 273	O 332	S 6	0	0	0
3	M	200	Total 3038	C 970	H 1479	N 262	O 321	S 6	0	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0
4	H	1	Total O S 5 4 1	0	0
4	H	1	Total O S 5 4 1	0	0
4	H	1	Total O S 5 4 1	0	0
4	I	1	Total O S 5 4 1	0	0
4	I	1	Total O S 5 4 1	0	0
4	I	1	Total O S 5 4 1	0	0
4	L	1	Total O S 5 4 1	0	0
4	L	1	Total O S 5 4 1	0	0
4	M	1	Total O S 5 4 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	31	Total 31	O 31	0	0
5	B	46	Total 46	O 46	0	0
5	H	23	Total 23	O 23	0	0
5	I	36	Total 36	O 36	0	0
5	L	14	Total 14	O 14	0	0
5	M	21	Total 21	O 21	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Erythrocyte binding antigen 175




- Molecule 1: Erythrocyte binding antigen 175




- Molecule 2: Antibody Heavy Chain





- Molecule 2: Antibody Heavy Chain







- Molecule 3: Antibody Light Chain



- Molecule 3: Antibody Light Chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	101.95Å 53.53Å 156.79Å 90.00° 91.18° 90.00°	Depositor
Resolution (Å)	29.79 – 2.45 29.79 – 2.45	Depositor EDS
% Data completeness (in resolution range)	99.4 (29.79-2.45) 99.5 (29.79-2.45)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.48 (at 2.45Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, R_{free}	0.204 , 0.253 0.208 , 0.255	Depositor DCC
R_{free} test set	1998 reflections (3.19%)	wwPDB-VP
Wilson B-factor (Å ²)	55.6	Xtrriage
Anisotropy	0.572	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 56.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.019 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	20975	wwPDB-VP
Average B, all atoms (Å ²)	89.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.22	0/1989	0.35	0/2654
1	B	0.22	0/2209	0.34	0/2947
2	H	0.23	0/1644	0.42	0/2251
2	I	0.23	0/1692	0.42	0/2315
3	L	0.22	0/1656	0.38	0/2245
3	M	0.23	0/1593	0.41	0/2162
All	All	0.22	0/10783	0.39	0/14574

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1950	1919	1926	11	0
1	B	2169	2144	2150	13	0
2	H	1599	1544	1551	8	0
2	I	1646	1589	1596	9	0
3	L	1620	1526	1533	8	0
3	M	1559	1479	1485	11	0
4	A	10	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	5	0	0	0	0
4	H	15	0	0	1	0
4	I	15	0	0	1	0
4	L	10	0	0	0	0
4	M	5	0	0	0	0
5	A	31	0	0	1	0
5	B	46	0	0	1	0
5	H	23	0	0	0	0
5	I	36	0	0	0	0
5	L	14	0	0	0	0
5	M	21	0	0	0	0
All	All	10774	10201	10241	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:M:108:ARG:NH1	3:M:109:ALA:O	1.98	0.97
1:A:30:ASP:OD2	1:A:33:ASN:N	2.17	0.77
1:A:194:ARG:NH2	5:A:423:HOH:O	2.24	0.70
3:L:155:ARG:NH2	3:L:185:GLU:OE2	2.33	0.62
1:B:245:VAL:HB	1:B:246:PRO:CD	2.34	0.58
3:L:29:ILE:HD11	3:L:71:TYR:CE1	2.39	0.57
3:M:145:ASN:OD1	3:M:146:VAL:N	2.37	0.57
2:H:100(B):TYR:OH	4:H:303:SO4:O4	2.17	0.57
1:A:217:ASP:N	1:A:218:PRO:HD2	2.19	0.57
2:H:38:LYS:HE2	2:H:40:ARG:HD2	1.87	0.57
1:B:88:CYS:SG	1:B:166:CYS:CB	2.93	0.56
3:M:24:ARG:NH1	3:M:70:ASP:OD1	2.38	0.56
2:I:100(B):TYR:OH	4:I:303:SO4:O3	2.07	0.54
1:B:18:ARG:NH2	5:B:405:HOH:O	2.42	0.53
2:H:33:TYR:HB2	2:H:95:HIS:HB2	1.92	0.52
2:I:11:LEU:HB2	2:I:147:PRO:HG3	1.91	0.52
1:B:245:VAL:HB	1:B:246:PRO:HD3	1.93	0.50
1:A:17:LYS:NZ	1:A:43:GLN:O	2.44	0.50
1:B:26:LYS:HD2	1:B:109:PHE:CE1	2.47	0.50
3:M:136:LEU:N	3:M:136:LEU:HD12	2.27	0.49
2:I:33:TYR:HB2	2:I:95:HIS:HB2	1.94	0.49
1:B:88:CYS:SG	1:B:166:CYS:HB2	2.53	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:209:ARG:HB3	3:L:210:TYR:HA	1.94	0.49
1:A:87:PHE:CE2	1:A:91:LEU:HD11	2.48	0.49
3:M:33:LEU:HD22	3:M:71:TYR:CB	2.42	0.48
2:I:177:LEU:C	2:I:177:LEU:HD12	2.34	0.47
1:B:217:ASP:N	1:B:218:PRO:HD2	2.30	0.47
1:B:197:LEU:HB3	1:B:198:PRO:HD3	1.97	0.47
1:A:11:LEU:HD23	1:B:63:HIS:NE2	2.31	0.46
3:L:140:TYR:CG	3:L:141:PRO:HA	2.51	0.46
1:B:26:LYS:HD2	1:B:109:PHE:CZ	2.51	0.46
2:H:119:PRO:HB2	2:H:142:VAL:HG13	1.98	0.45
2:H:177:LEU:HD12	2:H:177:LEU:C	2.37	0.45
1:A:214:GLU:O	1:A:215:CYS:HB3	2.16	0.45
1:A:217:ASP:N	1:A:218:PRO:CD	2.79	0.44
2:I:35:HIS:ND1	2:I:50:ARG:HB3	2.32	0.44
1:A:28:LYS:HG3	1:A:109:PHE:CE1	2.52	0.44
2:I:145:TYR:CE1	2:I:150:VAL:HG23	2.53	0.44
3:M:142:LYS:HG2	3:M:142:LYS:O	2.18	0.44
3:L:204:GLN:N	3:L:204:GLN:OE1	2.51	0.43
1:B:245:VAL:CB	1:B:246:PRO:CD	2.95	0.43
2:I:100(C):TYR:HB3	3:M:34:ASN:ND2	2.33	0.43
3:L:136:LEU:N	3:L:136:LEU:HD12	2.34	0.42
3:M:29:ILE:HD11	3:M:71:TYR:CE1	2.53	0.42
3:M:161:ASN:HB3	3:M:175:MET:HE3	2.02	0.42
2:I:172:SER:O	2:I:173:ASP:HB3	2.20	0.42
2:I:188:TRP:CD1	2:I:189:PRO:HA	2.54	0.42
1:A:230:LYS:NZ	4:A:302:SO4:O4	2.49	0.41
3:M:50:TYR:O	3:M:51:THR:HB	2.20	0.41
2:H:96:TYR:CE2	2:H:98:GLY:HA3	2.55	0.41
1:B:246:PRO:O	1:B:247:LYS:C	2.58	0.41
2:H:35:HIS:ND1	2:H:50:ARG:HB3	2.36	0.41
3:L:50:TYR:O	3:L:51:THR:HB	2.20	0.41
2:H:148:GLU:HB3	2:H:149:PRO:HA	2.02	0.41
1:B:169:ILE:O	1:B:169:ILE:HG23	2.20	0.41
3:M:85:THR:HG22	3:M:103:LYS:HA	2.03	0.41
3:L:199:LYS:HB3	3:L:200:LYS:HA	2.03	0.41
1:A:87:PHE:CZ	1:A:91:LEU:HD11	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	220/297 (74%)	212 (96%)	8 (4%)	0	100	100
1	B	243/297 (82%)	235 (97%)	8 (3%)	0	100	100
2	H	202/233 (87%)	197 (98%)	5 (2%)	0	100	100
2	I	210/233 (90%)	201 (96%)	9 (4%)	0	100	100
3	L	204/234 (87%)	188 (92%)	16 (8%)	0	100	100
3	M	198/234 (85%)	189 (96%)	9 (4%)	0	100	100
All	All	1277/1528 (84%)	1222 (96%)	55 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	217/278 (78%)	217 (100%)	0	100	100
1	B	244/278 (88%)	243 (100%)	1 (0%)	91	94
2	H	181/201 (90%)	180 (99%)	1 (1%)	86	91
2	I	186/201 (92%)	185 (100%)	1 (0%)	88	93
3	L	182/205 (89%)	180 (99%)	2 (1%)	73	82
3	M	177/205 (86%)	177 (100%)	0	100	100
All	All	1187/1368 (87%)	1182 (100%)	5 (0%)	91	94

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	26	LYS
2	H	100(B)	TYR
2	I	100(B)	TYR
3	L	90	GLN
3	L	105	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	A	302	-	4,4,4	0.14	0	6,6,6	0.05	0
4	SO4	I	301	-	4,4,4	0.14	0	6,6,6	0.05	0
4	SO4	H	301	-	4,4,4	0.14	0	6,6,6	0.06	0
4	SO4	M	301	-	4,4,4	0.14	0	6,6,6	0.07	0
4	SO4	I	302	-	4,4,4	0.14	0	6,6,6	0.09	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SO4	L	301	-	4,4,4	0.14	0	6,6,6	0.05	0
4	SO4	H	302	-	4,4,4	0.14	0	6,6,6	0.04	0
4	SO4	H	303	-	4,4,4	0.13	0	6,6,6	0.12	0
4	SO4	L	302	-	4,4,4	0.14	0	6,6,6	0.06	0
4	SO4	I	303	-	4,4,4	0.13	0	6,6,6	0.11	0
4	SO4	A	301	-	4,4,4	0.13	0	6,6,6	0.06	0
4	SO4	B	301	-	4,4,4	0.14	0	6,6,6	0.05	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	302	SO4	1	0
4	H	303	SO4	1	0
4	I	303	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	230/297 (77%)	0.78	29 (12%) 3 2	51, 78, 153, 184	0
1	B	257/297 (86%)	0.83	37 (14%) 2 1	46, 72, 138, 166	0
2	H	208/233 (89%)	0.35	11 (5%) 26 23	48, 72, 112, 138	0
2	I	214/233 (91%)	0.47	15 (7%) 16 13	43, 67, 110, 136	0
3	L	208/234 (88%)	0.68	28 (13%) 3 2	51, 86, 142, 166	0
3	M	200/234 (85%)	0.68	23 (11%) 4 3	44, 78, 140, 159	0
All	All	1317/1528 (86%)	0.64	143 (10%) 5 3	43, 75, 139, 184	0

All (143) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	197	LEU	8.6
1	A	196	LYS	7.2
1	B	245	VAL	6.8
1	B	208	TYR	6.5
1	B	213	LYS	6.5
3	M	196	ALA	6.1
1	B	207	LEU	6.0
1	A	280	TYR	5.9
1	A	215	CYS	5.9
2	I	201	ALA	5.9
1	A	135	LYS	5.8
1	B	211	CYS	5.5
3	L	206	THR	5.5
3	M	149	LYS	5.2
1	B	246	PRO	5.1
1	B	210	ALA	4.9
3	M	197	THR	4.9
3	L	200	LYS	4.9
3	M	191	SER	4.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	216	ILE	4.7
3	L	184	ASP	4.5
2	I	100(D)	PHE	4.5
1	A	98	TYR	4.4
1	A	194	ARG	4.4
1	B	94	SER	4.4
1	A	274	ASP	4.4
1	B	283	CYS	4.3
2	I	200	PRO	4.2
1	A	53	LYS	4.1
3	L	209	ARG	4.1
1	A	195	SER	4.0
3	L	185	GLU	3.9
1	A	136	ILE	3.9
1	A	125	LYS	3.8
3	M	36	TYR	3.8
3	M	195	GLU	3.8
3	L	210	TYR	3.7
1	A	188	LEU	3.7
1	B	93	ASN	3.6
3	L	190	ASN	3.6
1	B	212	GLU	3.6
1	B	247	LYS	3.5
2	I	205	LYS	3.5
3	M	150	ILE	3.5
3	L	183	LYS	3.5
3	M	146	VAL	3.5
1	B	133	GLU	3.4
3	L	88	CYS	3.4
3	L	203	PHE	3.3
1	B	215	CYS	3.3
2	I	213	ARG	3.3
2	H	100(D)	PHE	3.3
1	B	98	TYR	3.3
3	L	192	TYR	3.2
1	A	142	LYS	3.2
3	L	150	ILE	3.2
3	L	197	THR	3.2
1	B	197	LEU	3.2
1	A	279	LYS	3.1
2	H	191	GLU	3.1
3	L	186	TYR	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	196	LYS	3.1
3	M	192	TYR	3.0
1	A	214	GLU	2.9
2	I	203	SER	2.9
3	L	198	HIS	2.9
3	M	152	GLY	2.9
2	I	47	TRP	2.9
3	L	173	TYR	2.9
1	A	265	LYS	2.8
3	L	35	TRP	2.8
1	A	139	PHE	2.8
1	B	255	ILE	2.8
3	M	193	THR	2.8
2	I	43	GLN	2.8
1	B	209	GLU	2.8
1	A	94	SER	2.8
2	I	100(C)	TYR	2.8
1	B	201	LYS	2.8
1	A	213	LYS	2.7
1	B	244	LYS	2.7
3	L	191	SER	2.7
3	L	199	LYS	2.7
1	B	248	GLU	2.7
3	L	205	HIS	2.7
3	M	8	THR	2.7
1	B	131	ILE	2.7
3	L	36	TYR	2.7
3	M	190	ASN	2.6
3	M	199	LYS	2.6
2	H	185	SER	2.6
2	H	173	ASP	2.6
2	H	192	THR	2.6
3	L	155	ARG	2.6
2	I	100(B)	TYR	2.5
1	A	126	GLY	2.5
3	L	91	GLY	2.5
2	I	156	SER	2.5
2	I	35	HIS	2.5
2	H	37	VAL	2.4
3	M	147	LYS	2.4
1	B	90	ASP	2.4
1	B	198	PRO	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	I	36	TRP	2.4
2	H	190	SER	2.4
1	B	28	LYS	2.4
1	A	216	ILE	2.4
2	H	47	TRP	2.4
2	H	194	THR	2.4
3	M	131	SER	2.4
1	A	46	ILE	2.4
1	B	9	GLU	2.4
1	B	217	ASP	2.4
3	M	91	GLY	2.3
3	L	107	LYS	2.3
3	L	98	PHE	2.3
3	M	87	PHE	2.3
1	B	91	LEU	2.3
3	M	109	ALA	2.3
2	H	189	PRO	2.3
1	A	275	ALA	2.3
1	A	97	ASP	2.2
1	B	96	LEU	2.2
1	A	29	ASN	2.2
1	B	263	ASP	2.2
1	B	200	SER	2.2
1	A	278	SER	2.2
1	A	64	PHE	2.2
1	B	256	LYS	2.1
1	B	15	ARG	2.1
1	B	257	ILE	2.1
3	M	155	ARG	2.1
3	M	157	ASN	2.1
2	I	70	THR	2.1
2	I	202	SER	2.1
1	B	203	LYS	2.1
3	L	96	TRP	2.1
2	H	159	LEU	2.1
3	M	151	ASP	2.0
3	L	170	ASP	2.0
3	L	89	GLN	2.0
1	A	10	VAL	2.0
3	M	143	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	SO4	I	301	5/5	0.66	0.74	199,200,201,202	0
4	SO4	L	302	5/5	0.78	0.26	145,151,152,153	0
4	SO4	A	302	5/5	0.90	0.35	117,123,125,127	0
4	SO4	A	301	5/5	0.90	0.15	113,117,123,128	0
4	SO4	H	302	5/5	0.91	0.14	136,137,138,138	0
4	SO4	I	303	5/5	0.95	0.35	76,77,103,108	0
4	SO4	B	301	5/5	0.95	0.11	123,124,128,131	0
4	SO4	I	302	5/5	0.96	0.13	80,81,84,87	0
4	SO4	L	301	5/5	0.96	0.12	76,85,95,95	0
4	SO4	M	301	5/5	0.97	0.11	62,71,82,84	0
4	SO4	H	301	5/5	0.97	0.11	64,64,76,81	0
4	SO4	H	303	5/5	0.97	0.21	76,83,92,102	0

6.5 Other polymers [i](#)

There are no such residues in this entry.