



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 17, 2024 – 02:08 PM JST

PDB ID : 8K2I
Title : Crystal structure of Group 2 Oligosaccharide/Monosaccharide-releasing beta-N-acetylhexosaminidase NgaAt from Arabidopsis thaliana in complex with GlcNAc-thiazoline
Authors : Sumida, T.; Fushinobu, S.
Deposited on : 2023-07-12
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	646	 2% 71% 13% • 13%
1	B	646	 % 73% 12% • 13%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 9217 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Oligosaccharide/Monosaccharide-releasing beta-N-acetylhexosaminidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	559	4463	2863	746	836	18	0	1	0
1	A	560	4470	2867	747	838	18	0	1	0

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	GLY	-	expression tag	UNP Q7Y231
B	-1	PRO	-	expression tag	UNP Q7Y231
B	0	GLY	-	expression tag	UNP Q7Y231
A	-2	GLY	-	expression tag	UNP Q7Y231
A	-1	PRO	-	expression tag	UNP Q7Y231
A	0	GLY	-	expression tag	UNP Q7Y231

- Molecule 2 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			7	4	3		
2	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 3 is 3AR,5R,6S,7R,7AR-5-HYDROXYMETHYL-2-METHYL-5,6,7,7A-TETRAHYDRO-3AH-PYRANO[3,2-D]THIAZOLE-6,7-DIOL (three-letter code: NGT) (formula: C₈H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	N	O	S	0	0
			14	8	1	4	1		
3	A	1	Total	C	N	O	S	0	0
			14	8	1	4	1		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	136	Total	O	0	0
			136	136		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	105	Total 105	O 105	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Oligosaccharide/Monosaccharide-releasing beta-N-acetylhexosaminidase



- Molecule 1: Oligosaccharide/Monosaccharide-releasing beta-N-acetylhexosaminidase





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	69.28Å 133.83Å 150.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.71 – 2.50 47.66 – 2.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (47.71-2.50) 100.0 (47.66-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.92 (at 2.51Å)	Xtrriage
Refinement program	REFMAC 5.8.0352	Depositor
R, R_{free}	0.157 , 0.230 0.165 , 0.226	Depositor DCC
R_{free} test set	2489 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	52.6	Xtrriage
Anisotropy	0.045	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 53.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	9217	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, CL, NGT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.75	10/4604 (0.2%)	1.13	17/6288 (0.3%)
1	B	0.76	6/4597 (0.1%)	1.16	15/6278 (0.2%)
All	All	0.76	16/9201 (0.2%)	1.15	32/12566 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	15
1	B	0	9
All	All	0	24

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	273	GLU	CD-OE1	7.74	1.34	1.25
1	A	152	GLU	CD-OE2	7.21	1.33	1.25
1	B	447	GLU	CD-OE1	6.33	1.32	1.25
1	A	294	GLU	CD-OE1	6.00	1.32	1.25
1	A	574	GLU	CD-OE2	-5.85	1.19	1.25
1	B	598	GLU	CD-OE2	5.69	1.31	1.25
1	A	331	GLU	CD-OE1	5.63	1.31	1.25
1	A	567	GLU	CD-OE2	5.44	1.31	1.25
1	A	384	GLU	CD-OE1	5.28	1.31	1.25
1	A	294	GLU	CD-OE2	5.20	1.31	1.25
1	B	294	GLU	CD-OE2	5.13	1.31	1.25
1	B	474	GLU	CD-OE2	5.06	1.31	1.25
1	B	487	GLU	CD-OE1	5.05	1.31	1.25
1	A	144	SER	CB-OG	5.04	1.48	1.42

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	526	GLU	CD-OE2	5.01	1.31	1.25
1	A	611	GLU	CD-OE2	5.01	1.31	1.25

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	517	ARG	NE-CZ-NH2	-11.49	114.56	120.30
1	B	296	ARG	NE-CZ-NH2	-8.21	116.19	120.30
1	A	413	ARG	NE-CZ-NH1	8.07	124.33	120.30
1	A	321	ARG	NE-CZ-NH2	-8.00	116.30	120.30
1	B	350	ASP	CB-CG-OD2	-7.47	111.58	118.30
1	B	486	THR	CA-CB-OG1	-7.02	94.26	109.00
1	B	552	ARG	NE-CZ-NH2	-6.71	116.94	120.30
1	A	633	ARG	NE-CZ-NH2	-6.67	116.97	120.30
1	A	350	ASP	CB-CG-OD2	-6.66	112.31	118.30
1	A	552	ARG	NE-CZ-NH2	-6.58	117.01	120.30
1	B	413	ARG	NE-CZ-NH1	6.55	123.57	120.30
1	A	633	ARG	NE-CZ-NH1	6.45	123.53	120.30
1	B	633	ARG	NE-CZ-NH2	-6.42	117.09	120.30
1	B	378	TYR	CB-CG-CD1	-6.39	117.17	121.00
1	B	488	ASN	CB-CA-C	-6.35	97.69	110.40
1	A	256	ASP	CB-CA-C	6.04	122.47	110.40
1	B	366	ARG	NE-CZ-NH1	5.92	123.26	120.30
1	A	350	ASP	CB-CG-OD1	5.87	123.58	118.30
1	A	75	ARG	NE-CZ-NH1	5.83	123.22	120.30
1	A	387	ARG	NE-CZ-NH1	5.68	123.14	120.30
1	B	75	ARG	NE-CZ-NH2	-5.61	117.50	120.30
1	A	125	ARG	NE-CZ-NH2	-5.59	117.50	120.30
1	A	366	ARG	NE-CZ-NH1	5.43	123.02	120.30
1	B	425	ASP	CB-CG-OD1	5.40	123.16	118.30
1	A	268	ARG	NE-CZ-NH1	-5.36	117.62	120.30
1	A	552	ARG	NE-CZ-NH1	5.29	122.94	120.30
1	A	152	GLU	OE1-CD-OE2	5.21	129.55	123.30
1	A	517	ARG	NE-CZ-NH2	-5.19	117.70	120.30
1	A	502	ASP	CB-CA-C	5.14	120.68	110.40
1	B	378	TYR	CB-CG-CD2	5.13	124.08	121.00
1	B	522	ARG	NE-CZ-NH2	-5.02	117.79	120.30
1	B	86	ARG	NE-CZ-NH2	-5.01	117.80	120.30

There are no chirality outliers.

All (24) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	10	THR	Peptide
1	A	114	ARG	Sidechain
1	A	28	ALA	Peptide
1	A	296	ARG	Sidechain
1	A	321	ARG	Sidechain
1	A	334	ARG	Sidechain
1	A	357	ARG	Sidechain
1	A	380	ARG	Sidechain
1	A	387	ARG	Sidechain
1	A	473	ARG	Sidechain
1	A	512	ARG	Sidechain
1	A	517	ARG	Sidechain
1	A	522	ARG	Sidechain
1	A	627	ARG	Sidechain
1	A	641	ARG	Sidechain
1	B	296	ARG	Sidechain
1	B	333	MET	Peptide
1	B	334	ARG	Sidechain
1	B	380	ARG	Sidechain
1	B	512	ARG	Sidechain
1	B	553	ARG	Sidechain
1	B	583	ARG	Sidechain
1	B	621	ARG	Sidechain
1	B	94	SER	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4470	0	4363	41	0
1	B	4463	0	4356	38	0
2	A	7	0	10	0	0
2	B	7	0	10	0	0
3	A	14	0	13	2	0
3	B	14	0	13	0	0
4	A	1	0	0	1	0
5	A	105	0	0	0	0
5	B	136	0	0	0	0
All	All	9217	0	8765	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:378:TYR:CE2	1:B:382:GLU:HG3	2.16	0.80
1:B:375:ARG:HG3	1:B:375:ARG:HH11	1.48	0.78
1:B:466:SER:OG	1:B:468[A]:TRP:CD1	2.38	0.76
1:A:387:ARG:HG3	1:A:387:ARG:HH11	1.57	0.68
1:B:103:GLN:HG3	1:B:143:VAL:HG22	1.79	0.64
1:A:26:ASN:O	1:A:26:ASN:OD1	2.16	0.63
1:A:377:SER:O	1:A:381:LYS:HG3	1.99	0.63
1:B:346:HIS:ND1	1:B:347:PRO:HD2	2.15	0.61
1:B:369:ILE:O	1:B:369:ILE:HD12	2.00	0.61
1:A:316:TRP:O	1:A:319:GLN:HG2	2.02	0.60
1:A:69:ILE:HD12	1:A:69:ILE:N	2.17	0.59
1:B:366:ARG:HH11	1:B:366:ARG:CB	2.14	0.59
1:A:379:LEU:O	1:A:383:VAL:HG23	2.03	0.59
1:B:375:ARG:HH11	1:B:375:ARG:CG	2.17	0.58
1:B:380:ARG:HG3	1:B:422:TYR:CD1	2.41	0.55
1:A:73:ALA:O	1:A:271:VAL:HA	2.06	0.55
1:A:26:ASN:O	1:A:26:ASN:CG	2.46	0.54
1:B:365:TYR:OH	1:B:418:GLU:OE1	2.26	0.54
1:A:50:CYS:SG	1:A:69:ILE:HD11	2.48	0.53
1:A:409:PHE:HD1	1:A:456:LEU:HD21	1.75	0.52
1:A:641:ARG:HD2	4:A:703:CL:CL	2.47	0.52
1:B:181:LYS:HD2	1:B:182:THR:N	2.25	0.52
1:B:288:VAL:O	1:B:326:PHE:HA	2.11	0.51
1:B:452:VAL:N	1:B:453:PRO:CD	2.74	0.51
1:A:457:ARG:NH1	1:A:490:GLU:OE2	2.44	0.50
1:A:167:PRO:O	1:A:168:PRO:O	2.30	0.50
1:B:126:VAL:HG22	1:B:136:LEU:HD23	1.93	0.49
1:A:366:ARG:HB3	1:A:375:ARG:HD2	1.95	0.49
1:A:117:VAL:O	1:A:118:GLY:C	2.47	0.48
1:B:366:ARG:NH1	1:B:366:ARG:HB2	2.28	0.48
1:A:532:LEU:C	1:A:532:LEU:HD23	2.33	0.48
1:A:181:LYS:O	1:A:182:THR:C	2.53	0.47
1:A:566:GLY:HA2	1:A:576:VAL:HG23	1.96	0.47
1:A:641:ARG:N	1:A:642:PRO:CD	2.77	0.47
1:B:92:ALA:O	1:B:94:SER:N	2.47	0.47
1:A:613:THR:O	1:A:628:PRO:HB3	2.14	0.47
1:B:172:GLU:OE1	1:B:266:LYS:NZ	2.40	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:378:TYR:CD1	1:B:379:LEU:N	2.83	0.47
1:A:121:LEU:HD12	1:A:160:ILE:HG12	1.96	0.46
1:B:387:ARG:HD2	1:B:423:ALA:HB2	1.97	0.46
1:A:296:ARG:NH2	1:A:547:ALA:O	2.45	0.46
1:A:48:VAL:HG22	1:A:85:MET:HG2	1.98	0.46
1:A:364:PRO:HG2	1:A:378:TYR:OH	2.16	0.46
1:A:401:ASP:OD1	3:A:702:NGT:N2	2.49	0.46
1:A:263:LEU:HD23	1:A:263:LEU:C	2.35	0.45
1:B:417:SER:HB3	1:B:459:TYR:CE1	2.52	0.45
1:B:378:TYR:CD1	1:B:378:TYR:C	2.88	0.45
1:A:497:CYS:HB3	3:A:702:NGT:S1	2.56	0.45
1:B:28:ALA:O	1:B:29:GLU:HB3	2.16	0.45
1:B:429:LEU:HD23	1:B:430:THR:N	2.31	0.45
1:A:167:PRO:C	1:A:168:PRO:O	2.54	0.44
1:A:431:THR:HA	1:A:464:CYS:O	2.17	0.44
1:B:366:ARG:HH11	1:B:366:ARG:HB3	1.79	0.44
1:A:387:ARG:HG3	1:A:387:ARG:NH1	2.31	0.44
1:A:505:PRO:O	1:A:506:ASN:HB2	2.18	0.44
1:B:442:ALA:HB1	1:B:443:PRO:CD	2.48	0.44
1:A:349:SER:O	1:A:353:LEU:HG	2.18	0.43
1:B:259:SER:OG	1:B:261:LEU:HB2	2.17	0.43
1:A:504:HIS:HD2	1:A:522:ARG:HH12	1.67	0.43
1:B:369:ILE:HD12	1:B:369:ILE:C	2.39	0.43
1:A:267:LEU:HD12	1:A:267:LEU:N	2.34	0.43
1:A:309:LYS:HA	1:A:309:LYS:HD3	1.73	0.42
1:A:346:HIS:ND1	1:A:347:PRO:HD2	2.34	0.42
1:B:126:VAL:HG22	1:B:136:LEU:CD2	2.48	0.42
1:A:106:ASP:OD1	1:A:117:VAL:HG23	2.20	0.42
1:B:108:CYS:HA	1:B:113:ASP:O	2.19	0.42
1:B:74:ALA:HB1	1:B:274:PHE:O	2.20	0.42
1:B:452:VAL:N	1:B:453:PRO:HD2	2.33	0.42
1:B:366:ARG:CB	1:B:366:ARG:NH1	2.83	0.42
1:B:408:HIS:O	1:B:412:VAL:HG23	2.20	0.41
1:B:450:VAL:O	1:B:483:GLU:HG3	2.19	0.41
1:A:67:GLU:HB3	1:A:68:PRO:CD	2.51	0.41
1:B:276:ILE:HG23	1:B:592:GLU:HG3	2.03	0.41
1:B:323:SER:HB3	1:B:360:ALA:O	2.20	0.41
1:B:375:ARG:CG	1:B:375:ARG:NH1	2.77	0.40
1:B:181:LYS:HD2	1:B:182:THR:H	1.84	0.40
1:A:125:ARG:HD3	1:A:146:LEU:CD1	2.51	0.40
1:A:125:ARG:HD3	1:A:146:LEU:HD12	2.02	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:277:PRO:O	1:A:321:ARG:NH2	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	557/646 (86%)	536 (96%)	18 (3%)	3 (0%)	29	48
1	B	556/646 (86%)	533 (96%)	19 (3%)	4 (1%)	22	39
All	All	1113/1292 (86%)	1069 (96%)	37 (3%)	7 (1%)	25	43

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	168	PRO
1	B	26	ASN
1	B	29	GLU
1	A	497	CYS
1	B	334	ARG
1	B	497	CYS
1	A	303	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	490/566 (87%)	465 (95%)	25 (5%)	24	45
1	B	489/566 (86%)	471 (96%)	18 (4%)	34	60
All	All	979/1132 (86%)	936 (96%)	43 (4%)	29	52

All (43) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	27	ASP
1	B	30	PRO
1	B	31	LEU
1	B	110	SER
1	B	119	GLN
1	B	181	LYS
1	B	261	LEU
1	B	296	ARG
1	B	328	LYS
1	B	345	ASP
1	B	375	ARG
1	B	378	TYR
1	B	387	ARG
1	B	399	LEU
1	B	478	LYS
1	B	512	ARG
1	B	605	GLU
1	B	621	ARG
1	A	11	VAL
1	A	13	VAL
1	A	26	ASN
1	A	75	ARG
1	A	94	SER
1	A	120	SER
1	A	121	LEU
1	A	144	SER
1	A	170	GLN
1	A	300	GLU
1	A	374	SER
1	A	375	ARG
1	A	377	SER
1	A	378	TYR
1	A	379	LEU
1	A	384	GLU
1	A	417	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	458	PRO
1	A	468[A]	TRP
1	A	468[B]	TRP
1	A	474	GLU
1	A	532	LEU
1	A	601	TYR
1	A	625	GLU
1	A	641	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	504	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PEG	A	701	-	6,6,6	0.35	0	5,5,5	0.37	0
3	NGT	B	702	-	13,15,15	1.68	1 (7%)	12,22,22	1.41	3 (25%)
2	PEG	B	701	-	6,6,6	0.33	0	5,5,5	0.31	0
3	NGT	A	702	-	13,15,15	1.11	1 (7%)	12,22,22	1.03	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PEG	A	701	-	-	2/4/4/4	-
3	NGT	B	702	-	-	0/2/30/30	0/2/2/2
2	PEG	B	701	-	-	2/4/4/4	-
3	NGT	A	702	-	-	0/2/30/30	0/2/2/2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	702	NGT	C7-S1	-5.79	1.72	1.77
3	A	702	NGT	C7-S1	-2.96	1.74	1.77

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	702	NGT	O3-C3-C2	2.84	115.63	109.14
3	B	702	NGT	C3-C4-C5	-2.23	106.27	110.24
3	B	702	NGT	O3-C3-C4	-2.10	105.50	110.35

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	701	PEG	O2-C3-C4-O4
2	A	701	PEG	O1-C1-C2-O2
2	B	701	PEG	C1-C2-O2-C3
2	A	701	PEG	C1-C2-O2-C3

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	702	NGT	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	560/646 (86%)	-0.17	10 (1%) 68 71	36, 57, 94, 151	0
1	B	559/646 (86%)	-0.26	4 (0%) 87 89	32, 49, 83, 157	0
All	All	1119/1292 (86%)	-0.22	14 (1%) 77 79	32, 53, 89, 157	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	93	ALA	3.7
1	B	27	ASP	3.4
1	A	422	TYR	3.2
1	B	28	ALA	3.2
1	A	376	GLU	2.8
1	A	384	GLU	2.5
1	B	181	LYS	2.4
1	A	459	TYR	2.4
1	A	386	LEU	2.4
1	B	256	ASP	2.3
1	A	382	GLU	2.1
1	A	390	PRO	2.0
1	A	27	ASP	2.0
1	A	406	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PEG	A	701	7/7	0.93	0.13	58,73,80,81	0
2	PEG	B	701	7/7	0.95	0.11	59,66,74,80	0
4	CL	A	703	1/1	0.97	0.13	82,82,82,82	0
3	NGT	B	702	14/14	0.98	0.12	37,41,47,50	0
3	NGT	A	702	14/14	0.99	0.10	41,45,51,51	0

6.5 Other polymers [i](#)

There are no such residues in this entry.