



wwPDB X-ray Structure Validation Summary Report

Jun 12, 2024 – 11:07 PM EDT

PDB ID : 3K8Z
Title : Crystal Structure of Gudb1 a decriptified secondary glutamate dehydrogenase from *B. subtilis*
Authors : Gunka, K.; Newman, J.A.; Commichau, F.M.; Herzberg, C.; Rodrigues, C.; Hewitt, L.; Lewis, R.J.; Stulke, J.
Deposited on : 2009-10-15
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.20.1
EDS : 2.36.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	423	 7% 60% 27% 11%
1	B	423	 4% 60% 30% 7%
1	C	423	 % 59% 32% 5%
1	D	423	 6% 59% 28% 11%
1	E	423	 3% 60% 30% 7%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	423	 <p>A horizontal bar chart representing the quality of the chain. The bar is divided into three segments: a green segment on the left labeled '58%', a yellow segment in the middle labeled '33%', and a red segment on the right labeled '5%'. A small black dot is visible at the end of the red segment.</p>

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 18791 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NAD-specific glutamate dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	378	2929	1856	506	549	18	0	0	0
1	B	392	3049	1932	525	574	18	0	0	0
1	C	407	3169	2008	545	598	18	0	0	0
1	D	378	2929	1856	506	549	18	0	0	0
1	E	392	3049	1932	525	574	18	0	0	0
1	F	407	3169	2008	545	598	18	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	GLN	SEE REMARK 999	UNP P50735
A	?	-	VAL	SEE REMARK 999	UNP P50735
A	?	-	LYS	SEE REMARK 999	UNP P50735
A	?	-	ALA	SEE REMARK 999	UNP P50735
B	?	-	GLN	SEE REMARK 999	UNP P50735
B	?	-	VAL	SEE REMARK 999	UNP P50735
B	?	-	LYS	SEE REMARK 999	UNP P50735
B	?	-	ALA	SEE REMARK 999	UNP P50735
C	?	-	GLN	SEE REMARK 999	UNP P50735
C	?	-	VAL	SEE REMARK 999	UNP P50735
C	?	-	LYS	SEE REMARK 999	UNP P50735
C	?	-	ALA	SEE REMARK 999	UNP P50735
D	?	-	GLN	SEE REMARK 999	UNP P50735
D	?	-	VAL	SEE REMARK 999	UNP P50735
D	?	-	LYS	SEE REMARK 999	UNP P50735
D	?	-	ALA	SEE REMARK 999	UNP P50735
E	?	-	GLN	SEE REMARK 999	UNP P50735

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	?	-	VAL	SEE REMARK 999	UNP P50735
E	?	-	LYS	SEE REMARK 999	UNP P50735
E	?	-	ALA	SEE REMARK 999	UNP P50735
F	?	-	GLN	SEE REMARK 999	UNP P50735
F	?	-	VAL	SEE REMARK 999	UNP P50735
F	?	-	LYS	SEE REMARK 999	UNP P50735
F	?	-	ALA	SEE REMARK 999	UNP P50735


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	85	Total O 85 85	0	0
2	B	66	Total O 66 66	0	0
2	C	88	Total O 88 88	0	0
2	D	73	Total O 73 73	0	0
2	E	84	Total O 84 84	0	0
2	F	101	Total O 101 101	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NAD-specific glutamate dehydrogenase




- Molecule 1: NAD-specific glutamate dehydrogenase







- Molecule 1: NAD-specific glutamate dehydrogenase




- Molecule 1: NAD-specific glutamate dehydrogenase




- Molecule 1: NAD-specific glutamate dehydrogenase





- Molecule 1: NAD-specific glutamate dehydrogenase

Chain F: 58% 33% 5%



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	85.22Å 192.49Å 89.39Å 90.00° 118.74° 90.00°	Depositor
Resolution (Å)	19.77 – 2.40 19.77 – 2.40	Depositor EDS
% Data completeness (in resolution range)	97.9 (19.77-2.40) 96.6 (19.77-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.64 (at 2.41Å)	Xtrriage
Refinement program	PHENIX 1.5_2	Depositor
R, R_{free}	0.247 , 0.278 0.243 , 0.273	Depositor DCC
R_{free} test set	4788 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	39.1	Xtrriage
Anisotropy	0.389	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 22.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtrriage
Estimated twinning fraction	0.367 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	18791	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.23	0/2984	0.47	2/4025 (0.0%)
1	B	0.27	0/3105	0.49	2/4191 (0.0%)
1	C	0.24	0/3228	0.47	0/4359
1	D	0.23	0/2984	0.47	2/4025 (0.0%)
1	E	0.28	0/3105	0.63	3/4191 (0.1%)
1	F	0.24	0/3228	0.47	0/4359
All	All	0.25	0/18634	0.50	9/25150 (0.0%)

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	339	ARG	NE-CZ-NH1	-18.95	110.83	120.30
1	E	339	ARG	NE-CZ-NH2	18.04	129.32	120.30
1	E	339	ARG	CD-NE-CZ	8.97	136.16	123.60
1	D	339	ARG	NE-CZ-NH2	-6.34	117.13	120.30
1	A	339	ARG	NE-CZ-NH2	-6.32	117.14	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2929	0	2946	96	0
1	B	3049	0	3064	117	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	3169	0	3183	152	0
1	D	2929	0	2946	97	0
1	E	3049	0	3064	109	0
1	F	3169	0	3183	158	0
2	A	85	0	0	8	0
2	B	66	0	0	4	0
2	C	88	0	0	10	0
2	D	73	0	0	8	0
2	E	84	0	0	3	0
2	F	101	0	0	9	0
All	All	18791	0	18386	691	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 691 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:264:ILE:O	1:B:268:LEU:HB2	1.50	1.11
1:B:270:ARG:HH11	1:B:270:ARG:HG2	0.86	1.03
1:B:270:ARG:HH11	1:B:270:ARG:CG	1.73	0.97
1:B:270:ARG:HG2	1:B:270:ARG:NH1	1.65	0.96
1:F:44:GLU:HG3	1:F:45:PRO:HD2	1.50	0.93

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	374/423 (88%)	353 (94%)	19 (5%)	2 (0%)	29 41
1	B	388/423 (92%)	365 (94%)	17 (4%)	6 (2%)	10 14

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	405/423 (96%)	376 (93%)	25 (6%)	4 (1%)	15	23
1	D	374/423 (88%)	353 (94%)	18 (5%)	3 (1%)	19	29
1	E	388/423 (92%)	366 (94%)	19 (5%)	3 (1%)	19	29
1	F	405/423 (96%)	376 (93%)	25 (6%)	4 (1%)	15	23
All	All	2334/2538 (92%)	2189 (94%)	123 (5%)	22 (1%)	17	25

5 of 22 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	156	VAL
1	B	156	VAL
1	B	266	TYR
1	B	267	LEU
1	B	269	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	312/353 (88%)	300 (96%)	12 (4%)	33	51
1	B	326/353 (92%)	313 (96%)	13 (4%)	31	49
1	C	340/353 (96%)	315 (93%)	25 (7%)	13	22
1	D	312/353 (88%)	298 (96%)	14 (4%)	27	44
1	E	326/353 (92%)	313 (96%)	13 (4%)	31	49
1	F	340/353 (96%)	316 (93%)	24 (7%)	14	23
All	All	1956/2118 (92%)	1855 (95%)	101 (5%)	23	38

5 of 101 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	328	THR
1	E	339	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	F	378	LYS
1	D	372	GLU
1	E	50	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 46 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	363	GLN
1	E	305	ASN
1	D	364	ASN
1	E	193	HIS
1	E	364	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	378/423 (89%)	0.28	31 (8%) 11 10	26, 43, 77, 92	0
1	B	392/423 (92%)	0.22	17 (4%) 35 33	27, 43, 77, 92	0
1	C	407/423 (96%)	-0.18	3 (0%) 87 86	24, 39, 59, 74	0
1	D	378/423 (89%)	0.33	25 (6%) 18 17	27, 43, 77, 93	0
1	E	392/423 (92%)	0.12	12 (3%) 49 47	26, 43, 77, 92	0
1	F	407/423 (96%)	-0.10	0 100 100	23, 39, 59, 74	0
All	All	2354/2538 (92%)	0.11	88 (3%) 41 41	23, 42, 75, 93	0

The worst 5 of 88 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	253	TYR	8.7
1	D	257	TYR	5.7
1	A	253	TYR	5.5
1	A	264	ILE	5.0
1	B	254	GLY	4.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.