



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 22, 2024 – 03:01 PM EDT

PDB ID : 6K8I
Title : Crystal structure of Arabidopsis thaliana CRY2
Authors : Ma, L.; Wang, X.; Guan, Z.; Yin, P.
Deposited on : 2019-06-12
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 2.37.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	612	 3% 58% 16% 25%
1	B	612	 2% 57% 18% 25%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7608 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cryptochrome-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	457	3728	2412	632	668	16	0	1	0
1	B	461	3752	2427	638	671	16	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	78	GLN	LYS	See Sequence details	UNP Q96524
A	95	PRO	ALA	See Sequence details	UNP Q96524
A	366	GLY	ALA	See Sequence details	UNP Q96524
A	534	VAL	ALA	See Sequence details	UNP Q96524
B	78	GLN	LYS	See Sequence details	UNP Q96524
B	95	PRO	ALA	See Sequence details	UNP Q96524
B	366	GLY	ALA	See Sequence details	UNP Q96524
B	534	VAL	ALA	See Sequence details	UNP Q96524

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	A	1	53	27	9	15	2	0	0
2	B	1	53	27	9	15	2	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	9	9	9	0	0
3	B	13	13	13	0	0

M378	GLU	ALA	ALA
D387	LEU	GLY	GLY
W397	ALA	ASN	ASN
Y399	THR	THR	THR
I400	ILE	ILE	ILE
S401	LYS	LYS	LYS
G407	GLU	GLU	GLU
H408	PRO	PRO	PRO
E409	GLY	GLY	GLY
L410	LEU	LEU	LEU
D411	CYS	CYS	CYS
R412	PRO	PRO	PRO
	SER	SER	SER
	VAL	VAL	VAL
Q419	SER	SER	SER
	ALA	ALA	ALA
K422	SER	SER	SER
Y423	ASP	ASP	ASP
	GLN	GLN	GLN
R431	LYS	LYS	LYS
	ASN	ASN	ASN
L437	LEU	LEU	LEU
A438	GLU	GLU	GLU
F439	GLY	GLY	GLY
L440	ILE	ILE	ILE
	ARG	ARG	ARG
I445	GLN	GLN	GLN
H446	ASP	ASP	ASP
H447	SER	SER	SER
	GLY	GLY	GLY
	SER	SER	SER
G460	ASP	ASP	ASP
	GLN	GLN	GLN
	ILE	ILE	ILE
N466	THR	THR	THR
	LYS	LYS	LYS
I485	PRO	PRO	PRO
S486	SER	SER	SER
R487	LEU	LEU	LEU
T488	LYS	LYS	LYS
	ASN	ASN	ASN
	GLY	GLY	GLY
	CYS	CYS	CYS
	LYS	LYS	LYS
	GLU	GLU	GLU
	ARG	ARG	ARG
	ASP	ASP	ASP
	MET	MET	MET
	LYS	LYS	LYS
	LYS	LYS	LYS
	ILE	ILE	ILE
	ILE	ILE	ILE
	GLY	GLY	GLY
	SER	SER	SER
	ARG	ARG	ARG
	ALA	ALA	ALA
	ALA	ALA	ALA
	PRO	PRO	PRO
	ASP	ASP	ASP
	GLU	GLU	GLU
	ARG	ARG	ARG
	GLU	GLU	GLU
	ALA	ALA	ALA
	GLN	GLN	GLN
	ALA	ALA	ALA
	GLU	GLU	GLU
	ASP	ASP	ASP
	ILE	ILE	ILE
	LYS	LYS	LYS
	GLY	GLY	GLY
	THR	THR	THR
	LEU	LEU	LEU
	LEU	LEU	LEU
	LYS	LYS	LYS
	ASN	ASN	ASN
	GLY	GLY	GLY
	CYS	CYS	CYS
	LYS	LYS	LYS

4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	123.12Å 160.37Å 139.92Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.03 – 2.70 49.04 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.8 (49.03-2.70) 99.8 (49.04-2.70)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.26 (at 2.69Å)	Xtrriage
Refinement program	PHENIX (1.15.2_3472: ???)	Depositor
R, R_{free}	0.185 , 0.237 0.189 , 0.240	Depositor DCC
R_{free} test set	1948 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	61.6	Xtrriage
Anisotropy	0.681	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 54.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7608	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.44	0/3839	0.60	1/5219 (0.0%)
1	B	0.47	0/3863	0.63	3/5251 (0.1%)
All	All	0.45	0/7702	0.62	4/10470 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	422	LYS	CD-CE-NZ	-5.52	99.01	111.70
1	A	459	SER	C-N-CA	-5.43	110.89	122.30
1	B	121	GLU	CA-CB-CG	-5.37	101.58	113.40
1	B	422	LYS	CB-CG-CD	-5.12	98.27	111.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3728	0	3657	71	0
1	B	3752	0	3690	77	0
2	A	53	0	30	5	0
2	B	53	0	30	3	0
3	A	9	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	13	0	0	0	0
All	All	7608	0	7407	142	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 142 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:701:FAD:C1B	2:A:701:FAD:O4B	1.63	1.14
1:A:266:ARG:NH1	1:B:172:TRP:HE3	1.46	1.13
1:A:266:ARG:NH1	1:B:172:TRP:CE3	2.21	1.09
1:A:266:ARG:CZ	1:B:172:TRP:HE3	1.75	0.99
1:A:189:ILE:HD12	1:A:190:GLU:H	1.32	0.91

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	450/612 (74%)	432 (96%)	18 (4%)	0	100	100
1	B	453/612 (74%)	427 (94%)	26 (6%)	0	100	100
All	All	903/1224 (74%)	859 (95%)	44 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	399/531 (75%)	389 (98%)	10 (2%)	47	76
1	B	402/531 (76%)	392 (98%)	10 (2%)	47	76
All	All	801/1062 (75%)	781 (98%)	20 (2%)	49	76

5 of 20 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	267	MET
1	B	322	ASP
1	B	447	HIS
1	B	378	MET
1	A	440	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	415	ASN
1	B	408	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FAD	B	701	-	54,58,58	3.67	23 (42%)	71,89,89	2.29	18 (25%)
2	FAD	A	701	-	54,58,58	3.84	22 (40%)	71,89,89	2.14	15 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	B	701	-	-	4/30/50/50	0/6/6/6
2	FAD	A	701	-	-	4/30/50/50	0/6/6/6

The worst 5 of 45 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	701	FAD	O4B-C1B	17.37	1.63	1.40
2	B	701	FAD	O4B-C1B	15.96	1.61	1.40
2	A	701	FAD	C4X-N5	7.12	1.46	1.30
2	B	701	FAD	P-O3P	7.09	1.67	1.59
2	A	701	FAD	C10-N1	7.06	1.47	1.33

The worst 5 of 33 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	701	FAD	C5A-C6A-N6A	8.80	133.72	120.31
2	B	701	FAD	C5A-C6A-N6A	8.32	132.98	120.31
2	B	701	FAD	C7M-C7-C6	-7.16	106.96	119.57
2	B	701	FAD	N3A-C2A-N1A	-6.36	120.04	128.67
2	A	701	FAD	N3A-C2A-N1A	-6.02	120.50	128.67

There are no chirality outliers.

5 of 8 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	701	FAD	C5B-O5B-PA-O1A
2	B	701	FAD	C5B-O5B-PA-O2A
2	B	701	FAD	C5B-O5B-PA-O3P
2	B	701	FAD	C4'-C5'-O5'-P
2	A	701	FAD	C4'-C5'-O5'-P


There are no ring outliers.

2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	701	FAD	3	0
2	A	701	FAD	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	457/612 (74%)	0.17	21 (4%) 32 31	41, 69, 121, 165	0
1	B	461/612 (75%)	0.02	11 (2%) 59 60	42, 62, 115, 147	0
All	All	918/1224 (75%)	0.09	32 (3%) 44 44	41, 65, 120, 165	0

The worst 5 of 32 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	147	PRO	8.0
1	A	141	TYR	6.0
1	B	320	PRO	5.6
1	B	410	LEU	5.1
1	A	405	PRO	4.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	FAD	A	701	53/53	0.98	0.17	44,52,63,70	0
2	FAD	B	701	53/53	0.98	0.20	36,48,57,63	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.