

# Full wwPDB X-ray Structure Validation Report (i)

## Aug 22, 2020 - 06:22 PM BST

PDB ID	:	4KE $8$
Title	:	Crystal structure of Monoglyceride lipase from Bacillus sp. H257 in complex
		with monopalmitoyl glycerol analogue
Authors	:	Rengachari, S.; Aschauer, P.; Gruber, K.; Dreveny, I.; Oberer, M.
Deposited on		
Resolution	:	1.85  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:


Ŭ.	:	4.02b-467 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
$\mathrm{EDS}$	:	2.13.1
$\mathrm{buster}$ -report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
$\operatorname{Refmac}$	:	5.8.0158
$\operatorname{CCP4}$	:	$7.0.044 (\mathrm{Gargrove})$
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.13.1

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries},{ m resolution\ range}({ m \AA}))$
R <sub>free</sub>	130704	2469 (1.86-1.86)
Clashscore	141614	2625(1.86-1.86)
Ramachandran outliers	138981	2592(1.86-1.86)
Sidechain outliers	138945	2592(1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	270	86%	6%	8%
1	В	270	% • 84%	7% •	8%
1	С	270	90%	•	8%
1	D	270	88%	•	8%



# 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 8658 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Λ	248	Total	С	Ν	Ο	$\mathbf{S}$	0	0	0
	A	240	1912	1220	322	360	10	0	0	0
1	В	248	Total	С	Ν	0	S	0	1	0
	D	240	1913	1222	321	360	10	0		
1	С	249	Total	С	Ν	0	S	0	2	0
	U	249	1925	1230	323	362	10	0		0
1	п	240	Total	С	Ν	Ο	S	0	0	0
		249	1914	1220	322	362	10	0	U	0

• Molecule 1 is a protein called Thermostable monoacylglycerol lipase.

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	EXPRESSION TAG	UNP P82597
A	-18	GLY	-	EXPRESSION TAG	UNP P82597
A	-17	SER	-	EXPRESSION TAG	UNP P82597
A	-16	SER	-	EXPRESSION TAG	UNP P82597
A	-15	HIS	-	EXPRESSION TAG	UNP P82597
A	-14	HIS	-	EXPRESSION TAG	UNP P82597
A	-13	HIS	-	EXPRESSION TAG	UNP P82597
A	-12	HIS	-	EXPRESSION TAG	UNP P82597
A	-11	HIS	-	EXPRESSION TAG	UNP P82597
A	-10	HIS	-	EXPRESSION TAG	UNP P82597
A	-9	SER	-	EXPRESSION TAG	UNP P82597
A	-8	SER	-	EXPRESSION TAG	UNP P82597
A	-7	GLY	-	EXPRESSION TAG	UNP P82597
A	-6	LEU	_	EXPRESSION TAG	UNP P82597
A	-5	VAL	-	EXPRESSION TAG	UNP P82597
A	-4	PRO	-	EXPRESSION TAG	UNP P82597
A	-3	ARG	-	EXPRESSION TAG	UNP P82597
А	-2	GLY	-	EXPRESSION TAG	UNP P82597
A	-1	SER	-	EXPRESSION TAG	UNP P82597
А	0	HIS	-	EXPRESSION TAG	UNP P82597
В	-19	MET	-	EXPRESSION TAG	UNP P82597




B B B	-18	GLY			
		GLI	_	EXPRESSION TAG	UNP P82597
В	-17	SER	-	EXPRESSION TAG	UNP P82597
	-16	SER	-	EXPRESSION TAG	UNP P82597
В	-15	HIS	_	EXPRESSION TAG	UNP P82597
В	-14	HIS	-	EXPRESSION TAG	UNP P82597
В	-13	HIS	-	EXPRESSION TAG	UNP P82597
В	-12	HIS	-	EXPRESSION TAG	UNP P82597
В	-11	HIS	-	EXPRESSION TAG	UNP P82597
В	-10	HIS	-	EXPRESSION TAG	UNP P82597
В	-9	SER	-	EXPRESSION TAG	UNP P82597
В	-8	SER	-	EXPRESSION TAG	UNP P82597
В	-7	GLY	-	EXPRESSION TAG	UNP P82597
В	-6	LEU	-	EXPRESSION TAG	UNP P82597
В	-5	VAL	-	EXPRESSION TAG	UNP P82597
В	-4	PRO	-	EXPRESSION TAG	UNP P82597
В	-3	ARG	_	EXPRESSION TAG	UNP P82597
В	-2	GLY	-	EXPRESSION TAG	UNP P82597
В	-1	SER	_	EXPRESSION TAG	UNP P82597
В	0	HIS	_	EXPRESSION TAG	UNP P82597
С	-19	MET	_	EXPRESSION TAG	UNP P82597
С	-18	GLY	_	EXPRESSION TAG	UNP P82597
С	-17	SER	-	EXPRESSION TAG	UNP P82597
С	-16	SER	_	EXPRESSION TAG	UNP P82597
С	-15	HIS	_	EXPRESSION TAG	UNP P82597
С	-14	HIS	_	EXPRESSION TAG	UNP P82597
С	-13	HIS	_	EXPRESSION TAG	UNP P82597
С	-12	HIS	-	EXPRESSION TAG	UNP P82597
С	-11	HIS	_	EXPRESSION TAG	UNP P82597
С	-10	HIS	-	EXPRESSION TAG	UNP P82597
С	-9	SER	_	EXPRESSION TAG	UNP P82597
С	-8	SER	_	EXPRESSION TAG	UNP P82597
С	-7	GLY	-	EXPRESSION TAG	UNP P82597
С	-6	LEU	_	EXPRESSION TAG	UNP P82597
С	-5	VAL	-	EXPRESSION TAG	UNP P82597
С	-4	PRO	-	EXPRESSION TAG	UNP P82597
С	-3	ARG	_	EXPRESSION TAG	UNP P82597
С	-2	GLY	_	EXPRESSION TAG	UNP P82597
С	-1	SER	-	EXPRESSION TAG	UNP P82597
С	0	HIS	-	EXPRESSION TAG	UNP P82597
D	-19	MET	-	EXPRESSION TAG	UNP P82597
D	-18	GLY	-	EXPRESSION TAG	UNP P82597
D	-17	SER	-	EXPRESSION TAG	UNP P82597



Chain	Residue	Modelled	Actual	Comment	Reference
D	-16	SER	-	EXPRESSION TAG	UNP P82597
D	-15	HIS	-	EXPRESSION TAG	UNP P82597
D	-14	HIS	-	EXPRESSION TAG	UNP P82597
D	-13	HIS	-	EXPRESSION TAG	UNP P82597
D	-12	HIS	-	EXPRESSION TAG	UNP P82597
D	-11	HIS	-	EXPRESSION TAG	UNP P82597
D	-10	HIS	-	EXPRESSION TAG	UNP P82597
D	-9	SER	-	EXPRESSION TAG	UNP P82597
D	-8	SER	-	EXPRESSION TAG	UNP P82597
D	-7	GLY	-	EXPRESSION TAG	UNP P82597
D	-6	LEU	-	EXPRESSION TAG	UNP P82597
D	-5	VAL	-	EXPRESSION TAG	UNP P82597
D	-4	PRO	-	EXPRESSION TAG	UNP P82597
D	-3	ARG	-	EXPRESSION TAG	UNP P82597
D	-2	GLY	-	EXPRESSION TAG	UNP P82597
D	-1	SER	-	EXPRESSION TAG	UNP P82597
D	0	HIS	-	EXPRESSION TAG	UNP P82597

• Molecule 2 is tetradecyl hydrogen (R)-(3-azidopropyl)phosphonate (three-letter code: 1QY) (formula:  $C_{17}H_{36}N_3O_3P$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
0	Λ	1	Total	С	Ν	Ο	Р	0	0	
	2 A	L	23	17	3	2	1	0	0	
0	D	1	Total	С	Ν	Ο	Р	0	0	
	D	1	23	17	3	2	1	U	U	



Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
0	C	1	Total	С	Ν	Ο	Р	0	0
		L	23	17	3	2	1	0	0
0	Л	1	Total	С	Ν	Ο	Р	0	0
	D		23	17	3	2	1	0	U

• Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	208	Total O 208 208	0	0
3	В	185	Total O 185 185	0	0
3	С	254	Total         O           254         254	0	0
3	D	255	Total O 255 255	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Thermostable monoacylglycerol lipase





# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	77.02Å $81.12$ Å $85.61$ Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $100.26^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	43.74 - 1.85	Depositor
Resolution (A)	43.74 - 1.85	EDS
% Data completeness	92.0(43.74-1.85)	Depositor
(in resolution range)	$92.0 \ (43.74 - 1.85)$	EDS
R <sub>merge</sub>	(Not available)	Depositor
R <sub>sym</sub>	0.04	Depositor
$< I/\sigma(I) > {}^1$	$4.32 (at 1.86 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.8.2_1309, REFMAC	Depositor
D D.	0.180 , $0.220$	Depositor
$R, R_{free}$	0.180 , $0.220$	DCC
$R_{free}$ test set	1999 reflections $(2.45\%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	19.3	Xtriage
Anisotropy	0.968	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.31, 38.9	EDS
L-test for twinning <sup>2</sup>	$ L  > = 0.49, < L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	8658	wwPDB-VP
Average B, all atoms $(Å^2)$	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 44.55 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.5023e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section:  $1 \rm QY$ 

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
	Cham	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	А	0.34	0/1960	0.50	0/2668	
1	В	0.34	0/1964	0.50	0/2676	
1	С	0.36	0/1979	0.51	0/2693	
1	D	0.35	0/1962	0.49	0/2670	
All	All	0.35	0/7865	0.50	0/10707	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1912	0	1882	9	0
1	В	1913	0	1882	20	0
1	С	1925	0	1903	6	0
1	D	1914	0	1876	8	0
2	А	23	0	35	5	0
2	В	23	0	35	11	0
2	С	23	0	35	3	0
2	D	23	0	35	3	0
3	А	208	0	0	2	0



001000	naca jion	r procto de	$P^{ag}$ $\cdots$			
Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	В	185	0	0	2	0
3	С	254	0	0	1	0
3	D	255	0	0	1	0
All	All	8658	0	7683	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
		distance (Å)	overlap (Å)
1:B:125[B]:ILE:HG21	2:B:301:1QY:H20	1.52	0.90
1:B:125[A]:ILE:HG21	2:B:301:1QY:H20	1.55	0.86
1:B:61:HIS:ND1	1:B:63:GLU:OE1	2.09	0.83
1:B:138:LEU:HD11	2:B:301:1QY:H1	1.61	0.81
2:A:301:1QY:H34	3:A:548:HOH:O	1.79	0.80
1:B:138:LEU:HD11	2:B:301:1QY:C14	2.14	0.78
1:A:61:HIS:ND1	1:A:63:GLU:OE1	2.16	0.74
2:A:301:1QY:H31	3:A:604:HOH:O	1.94	0.66
1:B:226:HIS:CD2	2:B:301:1QY:H33	2.31	0.65
1:D:142:LEU:HD21	2:D:301:1QY:H2	1.80	0.64
1:B:160:GLU:HG3	1:B:161:LYS:HG3	1.78	0.63
1:C:142:LEU:HD21	2:C:301:1QY:H5	1.80	0.63
1:A:131:GLY:HA3	2:A:301:1QY:H6	1.83	0.60
1:B:97:SER:CB	2:B:301:1QY:H35	2.33	0.58
1:B:147:SER:OG	1:B:156:GLU:HB3	2.05	0.57
1:A:220:ARG:HD3	1:A:222:ARG:NH2	2.19	0.56
1:B:226:HIS:NE2	2:B:301:1QY:H33	2.20	0.56
1:C:77[A]:VAL:HG13	1:C:106:LEU:HD11	1.88	0.56
1:D:226:HIS:NE2	2:D:301:1QY:H32	2.20	0.56
1:B:220:ARG:NH2	3:B:421:HOH:O	2.29	0.55
1:C:176:GLN:NE2	3:C:645:HOH:O	2.38	0.55
1:C:142:LEU:HD11	2:C:301:1QY:H9	1.90	0.54
1:C:145:ILE:HD13	2:C:301:1QY:H22	1.93	0.50
1:B:16:ALA:HB3	1:B:50:VAL:HG22	1.95	0.49
1:A:170:LEU:HD21	2:A:301:1QY:H26	1.93	0.48
1:A:128:ILE:O	1:A:132:MET:HG2	2.14	0.47
1:A:132:MET:SD	1:A:138:LEU:HD21	2.55	0.46
1:C:25:LEU:CD1	1:C:50[B]:VAL:HG23	2.46	0.46
1:B:24:LEU:HD21	1:B:77:VAL:HG13	1.98	0.44
1:B:209:GLN:OE1	3:B:494:HOH:O	2.21	0.43



Atom-1	Atom-2	Interatomic	Clash
		distance (Å)	overlap (Å)
1:D:70:PHE:HA	1:D:73:TRP:CE3	2.53	0.43
2:A:301:1QY:H7	2:A:301:1QY:H12	1.87	0.43
1:D:222:ARG:NH2	3:D:610:HOH:O	2.45	0.43
1:B:97:SER:HB2	2:B:301:1QY:H35	2.01	0.42
1:B:151:ASN:HA	1:B:152:PRO:HD3	1.88	0.42
1:D:151:ASN:HA	1:D:152:PRO:HD3	1.87	0.42
2:B:301:1QY:H29	2:B:301:1QY:H31	1.53	0.42
1:B:194:ASP:OD1	1:B:220:ARG:HD3	2.19	0.42
1:A:16:ALA:HB3	1:A:50:VAL:HG22	2.01	0.42
1:B:123:VAL:HB	1:B:178:LYS:HB2	2.03	0.41
1:A:24:LEU:HD21	1:A:77:VAL:HG13	2.03	0.41
1:D:16:ALA:HB3	1:D:50:VAL:HG22	2.02	0.40
1:B:142:LEU:HD22	2:B:301:1QY:H5	2.03	0.40
1:D:195:GLU:O	1:D:225:TYR:HA	2.21	0.40
1:A:5:TYR:CG	1:A:61:HIS:HB2	2.56	0.40
1:B:138:LEU:HD11	2:B:301:1QY:H2	2.01	0.40
1:D:226:HIS:CD2	2:D:301:1QY:H35	2.56	0.40

There are no symmetry-related clashes.

#### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	А	246/270~(91%)	239~(97%)	7 (3%)	0	100	100
1	В	247/270~(92%)	241~(98%)	6(2%)	0	100	100
1	С	249/270~(92%)	242 (97%)	7 (3%)	0	100	100
1	D	247/270~(92%)	240~(97%)	7 (3%)	0	100	100
All	All	989/1080~(92%)	962~(97%)	27 (3%)	0	100	100

There are no Ramachandran outliers to report.



#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	204/222~(92%)	200~(98%)	4 (2%)	55 40
1	В	204/222~(92%)	202~(99%)	2(1%)	76 69
1	С	206/222~(93%)	204~(99%)	2(1%)	76 69
1	D	203/222 (91%)	202~(100%)	1 (0%)	88 86
All	All	817/888~(92%)	808~(99%)	9 (1%)	76 65

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	$\mathbf{Res}$	Type
1	А	2	SER
1	А	61	HIS
1	А	149	LEU
1	А	155	LYS
1	В	142	LEU
1	В	147	SER
1	С	77[A]	VAL
1	С	77[B]	VAL
1	D	85	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	В	248	HIS

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



#### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tune	Chain	Res	Res Link Bond lengths			Bond angles			
	Type	Cham	nes		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	1QY	С	301	1	18,22,23	1.04	1 (5%)	$15,\!22,\!25$	0.82	0
2	1QY	В	301	1	18,22,23	1.04	1 (5%)	$15,\!22,\!25$	0.82	0
2	1QY	D	301	1	$18,\!22,\!23$	1.03	1(5%)	$15,\!22,\!25$	0.82	0
2	1QY	А	301	1	18,22,23	1.03	1 (5%)	$15,\!22,\!25$	0.82	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1QY	С	301	1	-	7/17/21/23	-
2	1QY	В	301	1	-	10/17/21/23	-
2	1QY	D	301	1	-	8/17/21/23	-
2	1QY	А	301	1	-	9/17/21/23	_

All (4) bond length outliers are listed below:

Mol	Chain	$\mathbf{Res}$	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
2	В	301	1QY	N2-N1	-4.14	1.12	1.23
2	С	301	1QY	N2-N1	-4.14	1.12	1.23
2	D	301	1QY	N2-N1	-4.12	1.12	1.23
2	А	301	1QY	N2-N1	-4.10	1.12	1.23

There are no bond angle outliers.



There are no chirality outliers.

All (34) torsion outliers are listed below:

Mol	Chain	$\mathbf{Res}$	Type	Atoms
2	С	301	1QY	C1H-C2H-C3H-N1
2	В	301	1QY	C1H-C2H-C3H-N1
2	D	301	1QY	P1-C1H-C2H-C3H
2	D	301	1QY	C1H-C2H-C3H-N1
2	А	301	1QY	P1-C1H-C2H-C3H
2	А	301	1QY	C1H-C2H-C3H-N1
2	А	301	1QY	C7-C8-C9-C10
2	В	301	1QY	C2-C3-C4-C5
2	В	301	1QY	C11-C10-C9-C8
2	А	301	1QY	C4-C5-C6-C7
2	D	301	1QY	C11-C10-C9-C8
2	D	301	1QY	C5-C6-C7-C8
2	С	301	1QY	C3-C4-C5-C6
2	А	301	1QY	C1-C2-C3-C4
2	D	301	1QY	C4-C5-C6-C7
2	С	301	1QY	C7-C8-C9-C10
2	С	301	1QY	C4-C5-C6-C7
2	D	301	1QY	C3H-N1-N2-N3
2	В	301	1QY	C10-C11-C12-C13
2	А	301	1QY	C11-C12-C13-C14
2	В	301	1QY	C1-C2-C3-C4
2	В	301	1QY	C4-C5-C6-C7
2	А	301	1QY	C5-C6-C7-C8
2	В	301	1QY	C9-C10-C11-C12
2	D	301	1QY	C2-C3-C4-C5
2	А	301	1QY	C3-C4-C5-C6
2	С	301	1QY	C1-C2-C3-C4
2	D	301	1QY	C3-C4-C5-C6
2	С	301	1QY	C9-C10-C11-C12
2	А	301	1QY	C10-C11-C12-C13
2	В	301	1QY	C7-C8-C9-C10
2	В	301	1QY	C6-C7-C8-C9
2	В	301	1QY	C3-C4-C5-C6
2	С	301	1QY	C6-C7-C8-C9

There are no ring outliers.

4 monomers are involved in 22 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	С	301	1QY	3	0




Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	В	301	1QY	11	0
2	D	301	1QY	3	0
2	А	301	1QY	5	0

Continued from previous page...

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\# RSRZ {>}2$	$OWAB(Å^2)$	$\mathbf{Q}{<}0.9$
1	А	248/270~(91%)	-0.39	0 100 100	14, 24, 43, 60	0
1	В	248/270~(91%)	-0.35	2 (0%) 86 86	16, 25, 45, 62	0
1	С	249/270~(92%)	-0.46	0 100 100	14, 23, 37, 54	0
1	D	249/270~(92%)	-0.44	0 100 100	14, 23, 38, 60	0
All	All	994/1080~(92%)	-0.41	2 (0%) 95 94	14, 24, 40, 62	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	145	ILE	4.0
1	В	146	GLY	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	$Q{<}0.9$
2	1QY	D	301	23/24	0.84	0.19	$20,\!30,\!40,\!49$	23
2	1QY	В	301	23/24	0.85	0.24	$23,\!33,\!39,\!42$	23
2	1QY	С	301	23/24	0.86	0.22	$20,\!28,\!35,\!47$	23
2	1QY	А	301	23/24	0.90	0.18	$19,\!35,\!42,\!47$	23


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.











## 6.5 Other polymers (i)

There are no such residues in this entry.

