



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 22, 2020 – 10:37 pm BST

PDB ID : 4KES  
Title : Crystal structure of SsoPox W263T  
Authors : Gotthard, G.; Hiblot, J.; Chabriere, E.; Elias, M.  
Deposited on : 2013-04-26  
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*





The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	314	
1	B	314	
1	C	314	
1	D	314	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	A	409	-	-	X	-
5	GOL	D	406	-	-	X	-
6	EDO	D	407	-	-	X	-
6	EDO	D	408	-	-	X	-

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 10859 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aryldialkylphosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	314	2533	1621	433	471	8	0	3	0
1	B	314	2544	1627	437	472	8	0	4	0
1	C	314	2514	1610	428	469	7	0	1	0
1	D	314	2543	1627	435	474	7	0	4	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	263	THR	TRP	ENGINEERED MUTATION	UNP Q97VT7
B	263	THR	TRP	ENGINEERED MUTATION	UNP Q97VT7
C	263	THR	TRP	ENGINEERED MUTATION	UNP Q97VT7
D	263	THR	TRP	ENGINEERED MUTATION	UNP Q97VT7


- Molecule 2 is FE (II) ION (three-letter code: FE2) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Fe	0	0
			1	1		
2	A	1	Total	Fe	0	0
			1	1		
2	D	1	Total	Fe	0	0
			1	1		
2	C	1	Total	Fe	0	0
			1	1		

- Molecule 3 is COBALT (II) ION (three-letter code: CO) (formula: Co).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Co 1 1	0	0
3	A	1	Total Co 1 1	0	0
3	D	1	Total Co 1 1	0	0
3	C	1	Total Co 1 1	0	0

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 7 4 3	0	0
4	B	1	Total C O 7 4 3	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	C	1	Total C O 6 3 3	0	0
5	D	1	Total C O 6 3 3	0	0
5	D	1	Total C O 6 3 3	0	0
5	D	1	Total C O 6 3 3	0	0

- Molecule 6 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0
6	C	1	Total C O 4 2 2	0	0
6	C	1	Total C O 4 2 2	0	0
6	D	1	Total C O 4 2 2	0	0
6	D	1	Total C O 4 2 2	0	0
6	D	1	Total C O 4 2 2	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	150	Total O 150 150	0	0

*Continued on next page...*

*Continued from previous page...*


<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
7	B	168	Total 168	O 168	0	0
7	C	138	Total 138	O 138	0	0
7	D	123	Total 123	O 123	0	0




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Aryldialkylphosphatase




- Molecule 1: Aryldialkylphosphatase




- Molecule 1: Aryldialkylphosphatase



- Molecule 1: Aryldialkylphosphatase





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	84.20Å 103.60Å 151.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.52 – 2.10 49.52 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.9 (49.52-2.10) 99.9 (49.52-2.10)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.26 (at 2.10Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.163 , 0.205 0.163 , 0.207	Depositor DCC
$R_{free}$ test set	3903 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.6	Xtrriage
Anisotropy	0.086	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 50.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	10859	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.48% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CO, EDO, FE2, PEG, KCX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.37	0/2572	0.48	1/3470 (0.0%)
1	B	0.37	0/2583	0.47	1/3484 (0.0%)
1	C	0.36	0/2553	0.46	1/3446 (0.0%)
1	D	0.36	0/2582	0.46	1/3483 (0.0%)
All	All	0.36	0/10290	0.47	4/13883 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	256	ASP	CB-CG-OD1	6.24	123.92	118.30
1	A	256	ASP	CB-CG-OD1	6.08	123.77	118.30
1	C	256	ASP	CB-CG-OD1	5.42	123.18	118.30
1	D	256	ASP	CB-CG-OD1	5.30	123.07	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2533	0	2574	67	0
1	B	2544	0	2586	90	0
1	C	2514	0	2551	67	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	2543	0	2580	91	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	7	0	10	0	0
4	B	7	0	10	1	0
5	A	30	0	40	11	0
5	B	30	0	40	3	0
5	C	6	0	8	3	0
5	D	18	0	24	7	0
6	A	12	0	18	5	0
6	B	8	0	12	3	0
6	C	8	0	12	3	0
6	D	12	0	18	19	0
7	A	150	0	0	15	0
7	B	168	0	0	17	0
7	C	138	0	0	10	0
7	D	123	0	0	9	0
All	All	10859	0	10483	316	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 316 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1[A]:MET:CE	1:B:1[A]:MET:N	1.89	1.35
1:B:1[A]:MET:N	1:B:1[A]:MET:HE2	0.95	1.25
1:A:16:ILE:HG23	7:A:650:HOH:O	1.38	1.19
1:B:51:ASN:HB3	7:B:579:HOH:O	1.42	1.17
1:D:2[A]:ARG:HG2	1:D:2[A]:ARG:HH11	1.06	1.16

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	313/314 (100%)	300 (96%)	9 (3%)	4 (1%)	12	7
1	B	314/314 (100%)	302 (96%)	11 (4%)	1 (0%)	41	41
1	C	312/314 (99%)	294 (94%)	14 (4%)	4 (1%)	12	7
1	D	315/314 (100%)	302 (96%)	11 (4%)	2 (1%)	25	21
All	All	1254/1256 (100%)	1198 (96%)	45 (4%)	11 (1%)	25	12

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	266	ALA
1	B	273	LYS
1	C	266	ALA
1	D	2[A]	ARG
1	D	2[B]	ARG

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	275/272 (101%)	271 (98%)	4 (2%)	65	71
1	B	276/272 (102%)	271 (98%)	5 (2%)	59	65
1	C	273/272 (100%)	272 (100%)	1 (0%)	91	94
1	D	276/272 (102%)	273 (99%)	3 (1%)	73	79
All	All	1100/1088 (101%)	1087 (99%)	13 (1%)	81	77

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	1[B]	MET
1	B	2[A]	ARG
1	D	2[A]	ARG
1	B	1[A]	MET
1	C	254	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	58	GLN
1	C	127	GLN
1	D	127	GLN
1	C	34	GLN
1	D	131	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	KCX	A	137	1,3	7,11,12	0.51	0	4,12,14	0.69	0
1	KCX	B	137	1,3	7,11,12	0.50	0	4,12,14	0.65	0
1	KCX	C	137	1,3	7,11,12	0.51	0	4,12,14	0.73	0
1	KCX	D	137	1,3	7,11,12	0.46	0	4,12,14	1.09	1 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	A	137	1,3	-	0/7/10/12	-
1	KCX	B	137	1,3	-	0/7/10/12	-
1	KCX	C	137	1,3	-	1/7/10/12	-
1	KCX	D	137	1,3	-	3/7/10/12	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	137	KCX	CE-NZ-CX	2.12	126.54	122.95

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	D	137	KCX	C-CA-CB-CG
1	C	137	KCX	CG-CD-CE-NZ
1	D	137	KCX	CG-CD-CE-NZ
1	D	137	KCX	CA-CB-CG-CD

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	137	KCX	1	0
1	B	137	KCX	1	0
1	C	137	KCX	2	0
1	D	137	KCX	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 34 ligands modelled in this entry, 8 are monoatomic - leaving 26 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul



statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
6	EDO	D	403	-	3,3,3	0.46	0	2,2,2	0.41	0
6	EDO	C	405	-	3,3,3	0.47	0	2,2,2	0.32	0
6	EDO	C	404	-	3,3,3	0.47	0	2,2,2	0.37	0
5	GOL	B	407	-	5,5,5	0.25	0	5,5,5	0.23	0
5	GOL	C	403	-	5,5,5	0.26	0	5,5,5	0.29	0
5	GOL	A	408	-	5,5,5	0.26	0	5,5,5	0.24	0
6	EDO	D	408	-	3,3,3	0.50	0	2,2,2	0.41	0
6	EDO	A	411	-	3,3,3	0.46	0	2,2,2	0.39	0
5	GOL	B	409	-	5,5,5	0.25	0	5,5,5	0.38	0
5	GOL	B	405	-	5,5,5	0.21	0	5,5,5	0.46	0
5	GOL	D	405	-	5,5,5	0.27	0	5,5,5	0.31	0
5	GOL	A	404	-	5,5,5	0.29	0	5,5,5	0.45	0
6	EDO	B	401	-	3,3,3	0.47	0	2,2,2	0.36	0
5	GOL	D	404	-	5,5,5	0.35	0	5,5,5	0.17	0
6	EDO	A	406	-	3,3,3	0.49	0	2,2,2	0.26	0
5	GOL	B	406	-	5,5,5	0.30	0	5,5,5	0.25	0
5	GOL	D	406	-	5,5,5	0.25	0	5,5,5	0.30	0
6	EDO	D	407	-	3,3,3	0.49	0	2,2,2	0.24	0
5	GOL	A	410	-	5,5,5	0.31	0	5,5,5	0.50	0
5	GOL	B	404	-	5,5,5	0.24	0	5,5,5	0.41	0
4	PEG	B	408	-	6,6,6	0.43	0	5,5,5	0.36	0
6	EDO	A	407	-	3,3,3	0.45	0	2,2,2	0.42	0
4	PEG	A	403	-	6,6,6	0.45	0	5,5,5	0.23	0
6	EDO	B	410	-	3,3,3	0.45	0	2,2,2	0.38	0
5	GOL	A	409	-	5,5,5	0.26	0	5,5,5	0.34	0
5	GOL	A	405	-	5,5,5	0.27	0	5,5,5	0.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	D	403	-	-	0/1/1/1	-
6	EDO	C	405	-	-	0/1/1/1	-
6	EDO	C	404	-	-	1/1/1/1	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	B	407	-	-	3/4/4/4	-
5	GOL	C	403	-	-	0/4/4/4	-
5	GOL	A	408	-	-	2/4/4/4	-
6	EDO	D	408	-	-	0/1/1/1	-
6	EDO	A	411	-	-	1/1/1/1	-
5	GOL	B	409	-	-	2/4/4/4	-
5	GOL	B	405	-	-	0/4/4/4	-
5	GOL	D	405	-	-	4/4/4/4	-
5	GOL	A	404	-	-	2/4/4/4	-
6	EDO	B	401	-	-	1/1/1/1	-
5	GOL	D	404	-	-	0/4/4/4	-
6	EDO	A	406	-	-	1/1/1/1	-
5	GOL	B	406	-	-	2/4/4/4	-
5	GOL	D	406	-	-	2/4/4/4	-
6	EDO	D	407	-	-	1/1/1/1	-
5	GOL	A	410	-	-	2/4/4/4	-
5	GOL	B	404	-	-	0/4/4/4	-
4	PEG	B	408	-	-	1/4/4/4	-
6	EDO	A	407	-	-	1/1/1/1	-
4	PEG	A	403	-	-	3/4/4/4	-
6	EDO	B	410	-	-	0/1/1/1	-
5	GOL	A	409	-	-	0/4/4/4	-
5	GOL	A	405	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 31 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	407	GOL	C1-C2-C3-O3
5	A	410	GOL	C1-C2-C3-O3
4	A	403	PEG	O1-C1-C2-O2
6	C	404	EDO	O1-C1-C2-O2
5	A	408	GOL	C1-C2-C3-O3

There are no ring outliers.

17 monomers are involved in 55 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	C	404	EDO	3	0
5	C	403	GOL	3	0
5	A	408	GOL	2	0
6	D	408	EDO	13	0
6	A	411	EDO	2	0
5	A	404	GOL	1	0
6	B	401	EDO	1	0
6	A	406	EDO	2	0
5	B	406	GOL	2	0
5	D	406	GOL	7	0
6	D	407	EDO	6	0
5	A	410	GOL	1	0
5	B	404	GOL	1	0
4	B	408	PEG	1	0
6	A	407	EDO	1	0
6	B	410	EDO	2	0
5	A	409	GOL	7	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	313/314 (99%)	-0.32	5 (1%) 72 75	19, 32, 52, 80	0
1	B	313/314 (99%)	-0.36	4 (1%) 77 80	19, 30, 54, 86	0
1	C	313/314 (99%)	-0.17	8 (2%) 56 61	22, 39, 61, 93	0
1	D	313/314 (99%)	-0.06	5 (1%) 72 75	24, 43, 66, 107	0
All	All	1252/1256 (99%)	-0.23	22 (1%) 68 72	19, 36, 61, 107	0

The worst 5 of 22 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	1	MET	8.7
1	C	1	MET	7.2
1	C	268	PRO	4.7
1	D	268	PRO	4.4
1	C	266	ALA	4.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	KCX	A	137	12/13	0.98	0.11	17,18,21,23	0
1	KCX	B	137	12/13	0.98	0.09	18,19,21,21	0
1	KCX	C	137	12/13	0.98	0.09	24,27,29,31	0
1	KCX	D	137	12/13	0.98	0.08	25,30,34,35	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	PEG	B	408	7/7	0.76	0.18	55,65,69,75	0
5	GOL	A	408	6/6	0.79	0.16	51,60,67,69	0
5	GOL	B	407	6/6	0.81	0.19	43,55,63,73	0
6	EDO	B	410	4/4	0.83	0.17	48,53,55,64	0
5	GOL	D	405	6/6	0.84	0.13	64,66,72,73	0
6	EDO	B	401	4/4	0.84	0.15	53,54,56,60	0
6	EDO	A	406	4/4	0.85	0.16	43,51,52,56	0
5	GOL	D	406	6/6	0.87	0.17	55,62,65,66	0
6	EDO	A	407	4/4	0.89	0.18	58,58,66,68	0
6	EDO	D	408	4/4	0.89	0.82	89,94,97,102	0
5	GOL	A	409	6/6	0.89	0.39	50,61,62,66	0
6	EDO	D	407	4/4	0.90	0.69	54,61,63,69	0
6	EDO	C	404	4/4	0.91	0.12	52,53,53,55	0
4	PEG	A	403	7/7	0.91	0.15	56,57,69,71	0
5	GOL	A	410	6/6	0.91	0.13	30,35,52,57	0
5	GOL	B	409	6/6	0.91	0.17	50,57,64,67	0
6	EDO	A	411	4/4	0.92	0.16	40,59,65,65	0
6	EDO	D	403	4/4	0.93	0.08	44,47,55,62	0
5	GOL	C	403	6/6	0.94	0.13	46,58,61,64	0
6	EDO	C	405	4/4	0.94	0.15	41,45,47,50	0
5	GOL	B	404	6/6	0.94	0.10	43,48,50,51	0
5	GOL	A	404	6/6	0.94	0.11	36,40,47,49	0
5	GOL	A	405	6/6	0.94	0.17	33,35,44,48	0
5	GOL	B	406	6/6	0.95	0.11	30,40,46,53	0
5	GOL	B	405	6/6	0.97	0.14	33,34,42,58	0
5	GOL	D	404	6/6	0.97	0.15	26,36,44,51	0
2	FE2	C	401	1/1	0.98	0.10	24,24,24,24	0
2	FE2	D	401	1/1	0.99	0.09	23,23,23,23	0
2	FE2	A	401	1/1	0.99	0.10	19,19,19,19	0
3	CO	B	403	1/1	0.99	0.11	22,22,22,22	0
3	CO	C	402	1/1	0.99	0.07	30,30,30,30	0
2	FE2	B	402	1/1	0.99	0.10	18,18,18,18	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CO	D	402	1/1	0.99	0.11	25,25,25,25	0
3	CO	A	402	1/1	1.00	0.11	22,22,22,22	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.