



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 16, 2023 – 02:19 AM JST

PDB ID : 6KFV  
Title : GroEL from *Xanthomonas oryzae* pv. *oryzae*  
Authors : Tran, H.T.; Lee, J.H.; Kang, L.W.  
Deposited on : 2019-07-08  
Resolution : 3.22 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.22 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.











Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1335 (3.24-3.20)
Clashscore	141614	1460 (3.24-3.20)
Ramachandran outliers	138981	1437 (3.24-3.20)
Sidechain outliers	138945	1436 (3.24-3.20)
RSRZ outliers	127900	1291 (3.24-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	525	
1	B	525	
1	C	525	
1	D	525	
1	E	525	
1	F	525	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	G	525	 73% 22% 5%
1	H	525	 68% 25% 6% .
1	I	525	 65% 29% 6%
1	J	525	 72% 25% .
1	K	525	 69% 24% 6% .
1	L	525	 68% 26% 6% .
1	M	525	 71% 24% . .
1	N	525	 63% 29% 7% .

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 53933 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 60 kDa chaperonin.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	525	3862	2396	677	775	14	0	0	0
1	B	525	3858	2393	676	775	14	0	0	0
1	C	525	3862	2396	677	775	14	0	0	0
1	D	525	3854	2392	677	771	14	0	0	0
1	E	525	3854	2392	677	771	14	0	0	0
1	F	525	3839	2383	672	770	14	0	0	0
1	G	525	3858	2393	676	775	14	0	0	0
1	H	525	3856	2393	674	775	14	0	0	0
1	I	525	3842	2384	669	775	14	0	0	0
1	J	525	3854	2391	675	774	14	0	0	0
1	K	518	3809	2364	667	764	14	0	0	0
1	L	525	3859	2393	677	775	14	0	0	0
1	M	525	3854	2390	675	775	14	0	0	0
1	N	524	3811	2362	668	767	14	0	0	0

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	C O	0	0
			6	3 3		
2	F	1	Total	C O	0	0
			6	3 3		
2	F	1	Total	C O	0	0
			6	3 3		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	I	1	Total	O	S	0	0
			5	4	1		
3	K	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	6	Total	O	0	0
			6	6		
4	C	3	Total	O	0	0
			3	3		
4	D	3	Total	O	0	0
			3	3		
4	E	2	Total	O	0	0
			2	2		
4	F	1	Total	O	0	0
			1	1		
4	G	4	Total	O	0	0
			4	4		
4	H	1	Total	O	0	0
			1	1		
4	I	2	Total	O	0	0
			2	2		
4	J	2	Total	O	0	0
			2	2		
4	K	3	Total	O	0	0
			3	3		
4	M	5	Total	O	0	0
			5	5		
4	N	1	Total	O	0	0
			1	1		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 60 kDa chaperonin







- Molecule 1: 60 kDa chaperonin








• Molecule 1: 60 kDa chaperonin




• Molecule 1: 60 kDa chaperonin




• Molecule 1: 60 kDa chaperonin









• Molecule 1: 60 kDa chaperonin




• Molecule 1: 60 kDa chaperonin




• Molecule 1: 60 kDa chaperonin




• Molecule 1: 60 kDa chaperonin




• Molecule 1: 60 kDa chaperonin






• Molecule 1: 60 kDa chaperonin



• Molecule 1: 60 kDa chaperonin







## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	137.09Å 239.50Å 278.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.36 – 3.22 47.36 – 3.22	Depositor EDS
% Data completeness (in resolution range)	91.7 (47.36-3.22) 91.8 (47.36-3.22)	Depositor EDS
$R_{merge}$	0.19	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.12 (at 3.19Å)	Xtrriage
Refinement program	REFMAC 5.8.0238	Depositor
R, $R_{free}$	0.219 , 0.304 0.215 , 0.300	Depositor DCC
$R_{free}$ test set	6843 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	68.9	Xtrriage
Anisotropy	0.110	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 46.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.44$ , $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	53933	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	82.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.22% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.71	0/3892	0.82	1/5251 (0.0%)
1	B	0.70	0/3888	0.82	0/5247
1	C	0.71	0/3892	0.82	0/5251
1	D	0.72	0/3884	0.82	0/5241
1	E	0.71	0/3884	0.83	0/5241
1	F	0.71	0/3869	0.83	0/5223
1	G	0.71	0/3888	0.83	0/5247
1	H	0.71	0/3886	0.83	0/5244
1	I	0.71	0/3872	0.83	0/5229
1	J	0.72	0/3884	0.80	0/5242
1	K	0.72	0/3838	0.83	0/5177
1	L	0.72	0/3889	0.82	0/5247
1	M	0.73	0/3884	0.85	1/5243 (0.0%)
1	N	0.72	0/3840	0.82	0/5184
All	All	0.72	0/54290	0.83	2/73267 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	M	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	M	245	LYS	C-N-CD	-15.61	86.26	120.60
1	A	453	GLU	CB-CA-C	5.65	121.69	110.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	M	245	LYS	Mainchain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3862	0	3973	71	0
1	B	3858	0	3962	94	0
1	C	3862	0	3973	84	0
1	D	3854	0	3965	66	0
1	E	3854	0	3965	69	0
1	F	3839	0	3932	112	0
1	G	3858	0	3962	76	0
1	H	3856	0	3962	101	0
1	I	3842	0	3929	109	0
1	J	3854	0	3956	73	0
1	K	3809	0	3916	95	0
1	L	3859	0	3964	83	0
1	M	3854	0	3951	129	0
1	N	3811	0	3874	152	0
2	B	6	0	8	3	0
2	F	12	0	16	0	0
3	I	5	0	0	0	0
3	K	5	0	0	0	0
4	A	6	0	0	0	0
4	C	3	0	0	0	0
4	D	3	0	0	0	0
4	E	2	0	0	0	0
4	F	1	0	0	0	0
4	G	4	0	0	0	0
4	H	1	0	0	0	0
4	I	2	0	0	1	0
4	J	2	0	0	0	0
4	K	3	0	0	1	0
4	M	5	0	0	0	0

*Continued on next page...*



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	N	1	0	0	0	0
All	All	53933	0	55308	1262	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 1262 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:M:234:LEU:HA	1:M:237:LEU:CD2	1.34	1.55
1:N:228:SER:O	1:N:258:ALA:CB	1.63	1.42
1:F:234:LEU:CD2	1:F:235:PRO:HD3	1.52	1.40
1:M:234:LEU:CA	1:M:237:LEU:CD2	2.00	1.39
1:F:234:LEU:HD23	1:F:235:PRO:CD	1.53	1.38

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	523/525 (100%)	438 (84%)	65 (12%)	20 (4%)	<b>3</b>   21
1	B	523/525 (100%)	441 (84%)	66 (13%)	16 (3%)	<b>4</b>   26
1	C	523/525 (100%)	441 (84%)	59 (11%)	23 (4%)	<b>2</b>   18
1	D	523/525 (100%)	451 (86%)	60 (12%)	12 (2%)	<b>6</b>   33
1	E	523/525 (100%)	434 (83%)	72 (14%)	17 (3%)	<b>4</b>   24
1	F	523/525 (100%)	432 (83%)	69 (13%)	22 (4%)	<b>3</b>   19
1	G	523/525 (100%)	452 (86%)	62 (12%)	9 (2%)	<b>9</b>   40
1	H	523/525 (100%)	424 (81%)	75 (14%)	24 (5%)	<b>2</b>   17

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	I	523/525 (100%)	436 (83%)	62 (12%)	25 (5%)	2	16
1	J	523/525 (100%)	437 (84%)	65 (12%)	21 (4%)	3	20
1	K	514/525 (98%)	420 (82%)	69 (13%)	25 (5%)	2	16
1	L	523/525 (100%)	409 (78%)	86 (16%)	28 (5%)	2	13
1	M	523/525 (100%)	433 (83%)	65 (12%)	25 (5%)	2	16
1	N	520/525 (99%)	403 (78%)	90 (17%)	27 (5%)	2	14
All	All	7310/7350 (100%)	6051 (83%)	965 (13%)	294 (4%)	3	20

5 of 294 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	183	LEU
1	A	184	GLU
1	A	256	GLY
1	A	383	ALA
1	A	426	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	404/404 (100%)	353 (87%)	51 (13%)	4	20
1	B	403/404 (100%)	339 (84%)	64 (16%)	2	11
1	C	404/404 (100%)	361 (89%)	43 (11%)	6	27
1	D	402/404 (100%)	351 (87%)	51 (13%)	4	19
1	E	402/404 (100%)	348 (87%)	54 (13%)	4	17
1	F	398/404 (98%)	327 (82%)	71 (18%)	2	9
1	G	403/404 (100%)	341 (85%)	62 (15%)	2	12
1	H	403/404 (100%)	341 (85%)	62 (15%)	2	12
1	I	400/404 (99%)	345 (86%)	55 (14%)	3	16
1	J	402/404 (100%)	357 (89%)	45 (11%)	6	24

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	K	399/404 (99%)	341 (86%)	58 (14%)	3	14
1	L	403/404 (100%)	351 (87%)	52 (13%)	4	19
1	M	402/404 (100%)	354 (88%)	48 (12%)	5	22
1	N	392/404 (97%)	333 (85%)	59 (15%)	3	13
All	All	5617/5656 (99%)	4842 (86%)	775 (14%)	3	16

5 of 775 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	I	58	LYS
1	K	171	LYS
1	I	166	MET
1	I	50	THR
1	J	101	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 91 such sidechains are listed below:

Mol	Chain	Res	Type
1	I	208	GLN
1	K	206	ASN
1	I	265	ASN
1	J	348	GLN
1	K	475	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SO4	I	601	-	4,4,4	0.36	0	6,6,6	0.06	0
2	GOL	F	601	-	5,5,5	0.14	0	5,5,5	0.33	0
3	SO4	K	601	-	4,4,4	0.38	0	6,6,6	0.05	0
2	GOL	B	601	-	5,5,5	0.16	0	5,5,5	0.48	0
2	GOL	F	602	-	5,5,5	0.09	0	5,5,5	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	F	601	-	-	2/4/4/4	-
2	GOL	F	602	-	-	2/4/4/4	-
2	GOL	B	601	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 8 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	F	601	GOL	O1-C1-C2-C3
2	F	602	GOL	O1-C1-C2-O2
2	F	602	GOL	O1-C1-C2-C3
2	B	601	GOL	O1-C1-C2-C3
2	B	601	GOL	C1-C2-C3-O3


There are no ring outliers.


1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	601	GOL	3	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	525/525 (100%)	-0.31	2 (0%) 92 89	37, 68, 104, 128	0
1	B	525/525 (100%)	-0.22	3 (0%) 89 84	36, 68, 104, 124	0
1	C	525/525 (100%)	-0.20	3 (0%) 89 84	34, 73, 128, 168	0
1	D	525/525 (100%)	-0.20	7 (1%) 77 66	29, 78, 146, 173	0
1	E	525/525 (100%)	-0.12	3 (0%) 89 84	34, 79, 136, 183	0
1	F	525/525 (100%)	-0.21	3 (0%) 89 84	30, 73, 113, 137	0
1	G	525/525 (100%)	-0.26	2 (0%) 92 89	32, 65, 98, 127	0
1	H	525/525 (100%)	-0.18	7 (1%) 77 66	39, 75, 119, 156	0
1	I	525/525 (100%)	-0.25	1 (0%) 95 94	39, 74, 117, 162	0
1	J	525/525 (100%)	-0.07	8 (1%) 73 62	37, 90, 140, 165	0
1	K	518/525 (98%)	-0.04	8 (1%) 73 62	43, 84, 140, 179	0
1	L	525/525 (100%)	-0.09	5 (0%) 82 73	40, 82, 129, 148	0
1	M	525/525 (100%)	0.04	16 (3%) 50 36	41, 94, 146, 174	0
1	N	524/525 (99%)	-0.08	10 (1%) 66 54	38, 91, 149, 220	0
All	All	7342/7350 (99%)	-0.16	78 (1%) 80 70	29, 76, 133, 220	0

The worst 5 of 78 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	N	223	HIS	5.6
1	K	263	VAL	5.4
1	L	525	PRO	5.0
1	K	203	TYR	4.2
1	A	357	SER	3.9

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GOL	B	601	6/6	0.78	0.25	74,80,83,85	0
2	GOL	F	602	6/6	0.81	0.37	82,88,89,92	0
3	SO4	K	601	5/5	0.81	0.34	126,129,133,134	0
3	SO4	I	601	5/5	0.82	0.31	126,128,133,134	0
2	GOL	F	601	6/6	0.88	0.26	66,79,80,80	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around GOL F 602:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around SO4 K 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around SO4 I 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [i](#)

There are no such residues in this entry.