



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 19, 2023 – 09:51 PM EDT

PDB ID : 5KGM
Title : 2.95Å resolution structure of Apo independent phosphoglycerate mutase from *C. elegans* (monoclinic form)
Authors : Lovell, S.; Mehzabeen, N.; Battaile, K.P.; Yu, H.; Dranchak, P.; MacArthur, R.; Li, Z.; Carlow, T.; Suga, H.; Inglese, J.
Deposited on : 2016-06-13
Resolution : 2.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3104 (3.00-2.92)
Clashscore	141614	3462 (3.00-2.92)
Ramachandran outliers	138981	3340 (3.00-2.92)
Sidechain outliers	138945	3343 (3.00-2.92)
RSRZ outliers	127900	2986 (3.00-2.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	552	
1	B	552	

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 7845 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2,3-bisphosphoglycerate-independent phosphoglycerate mutase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	519	3933	2483	681	741	28	0	0	0
1	B	518	3904	2466	672	738	28	0	0	0

There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	540	LYS	-	expression tag	UNP G5EFZ1
A	541	LEU	-	expression tag	UNP G5EFZ1
A	542	ALA	-	expression tag	UNP G5EFZ1
A	543	ALA	-	expression tag	UNP G5EFZ1
A	544	ALA	-	expression tag	UNP G5EFZ1
A	545	LEU	-	expression tag	UNP G5EFZ1
A	546	GLU	-	expression tag	UNP G5EFZ1
A	547	HIS	-	expression tag	UNP G5EFZ1
A	548	HIS	-	expression tag	UNP G5EFZ1
A	549	HIS	-	expression tag	UNP G5EFZ1
A	550	HIS	-	expression tag	UNP G5EFZ1
A	551	HIS	-	expression tag	UNP G5EFZ1
A	552	HIS	-	expression tag	UNP G5EFZ1
B	540	LYS	-	expression tag	UNP G5EFZ1
B	541	LEU	-	expression tag	UNP G5EFZ1
B	542	ALA	-	expression tag	UNP G5EFZ1
B	543	ALA	-	expression tag	UNP G5EFZ1
B	544	ALA	-	expression tag	UNP G5EFZ1
B	545	LEU	-	expression tag	UNP G5EFZ1
B	546	GLU	-	expression tag	UNP G5EFZ1
B	547	HIS	-	expression tag	UNP G5EFZ1
B	548	HIS	-	expression tag	UNP G5EFZ1
B	549	HIS	-	expression tag	UNP G5EFZ1
B	550	HIS	-	expression tag	UNP G5EFZ1
B	551	HIS	-	expression tag	UNP G5EFZ1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	552	HIS	-	expression tag	UNP G5EFZ1

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Cl 2 2	0	0
2	B	2	Total Cl 2 2	0	0

- Molecule 3 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mn 1 1	0	0
3	B	1	Total Mn 1 1	0	0

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Zn 1 1	0	0
4	B	1	Total Zn 1 1	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	67.85Å 94.38Å 102.20Å 90.00° 96.57° 90.00°	Depositor
Resolution (Å)	42.98 – 2.95 46.48 – 2.95	Depositor EDS
% Data completeness (in resolution range)	99.6 (42.98-2.95) 99.6 (46.48-2.95)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.24 (at 2.96Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.184 , 0.237 0.187 , 0.234	Depositor DCC
R_{free} test set	1353 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	48.4	Xtrriage
Anisotropy	0.472	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 43.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7845	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MN, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/4021	0.62	0/5459
1	B	0.39	0/3992	0.62	1/5424 (0.0%)
All	All	0.39	0/8013	0.62	1/10883 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	426	ASP	CB-CG-OD1	7.50	125.05	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3933	0	3776	15	0
1	B	3904	0	3726	25	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	7845	0	7502	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:465:THR:HG22	1:B:466:ALA:H	1.58	0.68
1:B:476:ALA:HB1	1:B:477:PRO:HD2	1.88	0.55
1:A:233:GLY:N	1:A:271:ARG:HD2	2.21	0.55
1:A:24:ASN:O	1:A:27:VAL:HG12	2.12	0.49
1:A:145:GLY:O	1:A:283:TYR:OH	2.30	0.49
1:A:423:ALA:N	1:A:424:PRO:CD	2.76	0.48
1:B:465:THR:HG22	1:B:466:ALA:N	2.27	0.48
1:A:351:GLN:HA	1:A:417:PHE:O	2.14	0.48
1:B:74:LEU:HD11	1:B:80:GLU:HB2	1.95	0.48
1:B:24:ASN:O	1:B:27:VAL:HG12	2.14	0.47
1:B:356:GLU:HA	1:B:383:VAL:O	2.14	0.46
1:B:21:MET:SD	1:B:376:GLU:HG2	2.57	0.45
1:B:145:GLY:O	1:B:283:TYR:OH	2.34	0.45
1:B:352:PHE:CE2	1:B:354:CYS:HB2	2.51	0.45
1:A:85:ASN:ND2	1:B:387:LYS:HG2	2.31	0.45
1:A:267:GLY:O	1:A:271:ARG:NH1	2.50	0.45
1:B:36:ILE:O	1:B:466:ALA:HA	2.17	0.45
1:A:30:LYS:HB3	1:A:460:TYR:CE1	2.52	0.43
1:B:423:ALA:N	1:B:424:PRO:CD	2.81	0.43
1:B:179:THR:O	1:B:179:THR:HG23	2.19	0.43
1:B:179:THR:O	1:B:180:SER:O	2.37	0.42
1:B:265:LEU:O	1:B:266:GLN:HB2	2.19	0.42
1:B:278:ILE:CG2	1:B:315:VAL:HG22	2.49	0.42
1:B:232:GLY:HA3	1:B:307:LEU:HD13	2.00	0.42
1:B:232:GLY:HA3	1:B:307:LEU:CD1	2.50	0.41
1:B:465:THR:HG23	1:B:491:PRO:O	2.20	0.41
1:B:173:TYR:HA	1:B:208:VAL:O	2.21	0.41
1:B:282:ASP:HB2	1:B:288:MET:HE3	2.01	0.41
1:B:351:GLN:HA	1:B:417:PHE:O	2.21	0.41
1:A:406:MET:SD	1:A:420:CYS:HB3	2.61	0.41
1:A:27:VAL:HG11	1:A:416:PRO:HB3	2.03	0.41
1:A:282:ASP:HB2	1:A:288:MET:HE3	2.02	0.41
1:A:342:GLU:HB2	1:A:375:PHE:CZ	2.56	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:411:GLU:HG2	1:A:458:HIS:NE2	2.36	0.41
1:B:406:MET:SD	1:B:420:CYS:HB3	2.60	0.41
1:A:224:ASN:OD1	1:A:305:SER:HB2	2.20	0.40
1:B:27:VAL:HG11	1:B:416:PRO:HB3	2.03	0.40
1:A:424:PRO:HB2	1:A:425:PRO:HD3	2.03	0.40
1:B:500:LYS:HE2	1:B:539:ILE:HG21	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	515/552 (93%)	504 (98%)	8 (2%)	3 (1%)	25 60
1	B	514/552 (93%)	500 (97%)	11 (2%)	3 (1%)	25 60
All	All	1029/1104 (93%)	1004 (98%)	19 (2%)	6 (1%)	25 60

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	180	SER
1	A	470	ASN
1	B	470	ASN
1	B	468	HIS
1	A	200	GLY
1	A	468	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	403/445 (91%)	403 (100%)	0	100	100
1	B	398/445 (89%)	393 (99%)	5 (1%)	69	87
All	All	801/890 (90%)	796 (99%)	5 (1%)	86	94

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	25	SER
1	B	271	ARG
1	B	298	ASP
1	B	311	SER
1	B	500	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	519/552 (94%)	-0.05	2 (0%) 92 84	29, 45, 60, 95	0
1	B	518/552 (93%)	0.13	19 (3%) 41 27	33, 50, 75, 98	0
All	All	1037/1104 (93%)	0.04	21 (2%) 65 48	29, 47, 68, 98	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	252	ALA	3.7
1	B	476	ALA	3.3
1	B	185	VAL	3.3
1	B	182	ASN	3.2
1	B	183	SER	3.1
1	B	251	TYR	3.0
1	B	179	THR	2.7
1	B	479	GLY	2.7
1	B	254	ASP	2.6
1	B	256	THR	2.5
1	B	130	ASN	2.5
1	B	481	LYS	2.5
1	B	432	GLY	2.4
1	B	477	PRO	2.4
1	B	257	ASP	2.4
1	A	479	GLY	2.2
1	B	480	GLY	2.2
1	B	478	ASP	2.2
1	B	189	GLU	2.1
1	B	181	PRO	2.1
1	A	131	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CL	A	602	1/1	0.91	0.28	44,44,44,44	0
3	MN	B	603	1/1	0.94	0.18	45,45,45,45	0
3	MN	A	603	1/1	0.95	0.20	52,52,52,52	0
2	CL	A	601	1/1	0.96	0.21	34,34,34,34	0
2	CL	B	601	1/1	0.96	0.20	37,37,37,37	0
2	CL	B	602	1/1	0.98	0.37	49,49,49,49	0
4	ZN	A	604	1/1	0.99	0.18	54,54,54,54	0
4	ZN	B	604	1/1	1.00	0.08	41,41,41,41	0

6.5 Other polymers [i](#)

There are no such residues in this entry.