

wwPDB X-ray Structure Validation Summary Report (i)

Oct 11, 2023 – 03:14 AM EDT

PDB ID : 7KLS

Title: Dihydrodipicolinate synthase (DHDPS) from C.jejuni, H59A mutant with

pyruvate bound in the active site and L-lysine bound at the allosteric site

Authors: Saran, S.; Sanders, D.A.R.

Deposited on : 2020-10-31

Resolution : 2.59 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35.1

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}({\rm \AA})) \end{array}$
R_{free}	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	310	81%	14%	5%
1	В	310	79%	5%	• 5%
1	С	310		14%	• 5%
1	D	310	77% 18	3%	5%
1	Е	310	81%	14%	5%



Continued from previous page...

Mol	Chain	Length	Quality of chain		
			<u>%</u>		
1	F	310	80%	15%	• 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PGE	F	301	-	=	-	X



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 13985 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called 4-hydroxy-tetrahydrodipicolinate synthase.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	A	295	Total	С	N	Ο	S	0	1	0
1	A	290	2250	1431	369	437	13	U	1	
1	В	296	Total	С	N	О	S	0	0	0
1	Ъ	290	2266	1441	375	437	13	U	U	
1	С	295	Total	С	N	О	S	0	0	0
1		290	2259	1437	373	436	13	U		
1	D	296	Total	С	N	О	S	0	0	0
1	D	290	2255	1433	372	437	13	U	U	
1	Е	296	Total	С	N	О	S	0	0	0
1	l L	290	2266	1441	375	437	13	U	0	
1	F	296	Total	С	N	О	S	0	1	0
1	I.	290	2270	1446	374	437	13	U	0 1	0

There are 78 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	MET	-	expression tag	UNP Q9PPB4
A	-10	ARG	-	expression tag	UNP Q9PPB4
A	-9	GLY	-	expression tag	UNP Q9PPB4
A	-8	SER	-	expression tag	UNP Q9PPB4
A	-7	HIS	-	expression tag	UNP Q9PPB4
A	-6	HIS	-	expression tag	UNP Q9PPB4
A	-5	HIS	-	expression tag	UNP Q9PPB4
A	-4	HIS	-	expression tag	UNP Q9PPB4
A	-3	HIS	-	expression tag	UNP Q9PPB4
A	-2	HIS	-	expression tag	UNP Q9PPB4
A	-1	GLY	-	expression tag	UNP Q9PPB4
A	0	SER	-	expression tag	UNP Q9PPB4
A	59	ALA	HIS	engineered mutation	UNP Q9PPB4
В	-11	MET		expression tag	UNP Q9PPB4
В	-10	ARG	=	expression tag	UNP Q9PPB4
В	-9	GLY	-	expression tag	UNP Q9PPB4
В	-8	SER	-	expression tag	UNP Q9PPB4



Continued from previous page...


Chain	Residue	Modelled Modelled	Actual	Comment	Reference
В	-7	HIS	-	expression tag	UNP Q9PPB4
В	-6	HIS	-	expression tag	UNP Q9PPB4
В	-5	HIS	-	expression tag	UNP Q9PPB4
В	-4	HIS	_	expression tag	UNP Q9PPB4
В	-3	HIS	-	expression tag	UNP Q9PPB4
В	-2	HIS	-	expression tag	UNP Q9PPB4
В	-1	GLY	-	expression tag	UNP Q9PPB4
В	0	SER	-	expression tag	UNP Q9PPB4
В	59	ALA	HIS	engineered mutation	UNP Q9PPB4
С	-11	MET	_	expression tag	UNP Q9PPB4
С	-10	ARG	_	expression tag	UNP Q9PPB4
С	-9	GLY	-	expression tag	UNP Q9PPB4
С	-8	SER	-	expression tag	UNP Q9PPB4
С	-7	HIS	_	expression tag	UNP Q9PPB4
С	-6	HIS	_	expression tag	UNP Q9PPB4
С	-5	HIS	_	expression tag	UNP Q9PPB4
С	-4	HIS	-	expression tag	UNP Q9PPB4
С	-3	HIS	-	expression tag	UNP Q9PPB4
С	-2	HIS	_	expression tag	UNP Q9PPB4
С	-1	GLY	-	expression tag	UNP Q9PPB4
С	0	SER	-	expression tag	UNP Q9PPB4
С	59	ALA	HIS	engineered mutation	UNP Q9PPB4
D	-11	MET	-	expression tag	UNP Q9PPB4
D	-10	ARG	-	expression tag	UNP Q9PPB4
D	-9	GLY	-	expression tag	UNP Q9PPB4
D	-8	SER	-	expression tag	UNP Q9PPB4
D	-7	HIS	-	expression tag	UNP Q9PPB4
D	-6	HIS	-	expression tag	UNP Q9PPB4
D	-5	HIS	-	expression tag	UNP Q9PPB4
D	-4	HIS	-	expression tag	UNP Q9PPB4
D	-3	HIS	-	expression tag	UNP Q9PPB4
D	-2	HIS	_	expression tag	UNP Q9PPB4
D	-1	GLY	-	expression tag	UNP Q9PPB4
D	0	SER	_	expression tag	UNP Q9PPB4
D	59	ALA	HIS	engineered mutation	UNP Q9PPB4
Е	-11	MET	-	expression tag	UNP Q9PPB4
Е	-10	ARG	-	expression tag	UNP Q9PPB4
Е	-9	GLY	-	expression tag	UNP Q9PPB4
Е	-8	SER	-	expression tag	UNP Q9PPB4
Е	-7	HIS	-	expression tag	UNP Q9PPB4
Е	-6	HIS	-	expression tag	UNP Q9PPB4
Е	-5	HIS	-	expression tag	UNP Q9PPB4



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
Е	-4	HIS	-	expression tag	UNP Q9PPB4
Е	-3	HIS	-	expression tag	UNP Q9PPB4
Е	-2	HIS	-	expression tag	UNP Q9PPB4
Е	-1	GLY	-	expression tag	UNP Q9PPB4
E	0	SER	-	expression tag	UNP Q9PPB4
Е	59	ALA	HIS	engineered mutation	UNP Q9PPB4
F	-11	MET	-	expression tag	UNP Q9PPB4
F	-10	ARG	-	expression tag	UNP Q9PPB4
F	-9	GLY	-	expression tag	UNP Q9PPB4
F	-8	SER	-	expression tag	UNP Q9PPB4
F	-7	HIS	-	expression tag	UNP Q9PPB4
F	-6	HIS	-	expression tag	UNP Q9PPB4
F	-5	HIS	-	expression tag	UNP Q9PPB4
F	-4	HIS	-	expression tag	UNP Q9PPB4
F	-3	HIS		expression tag	UNP Q9PPB4
F	-2	HIS	-	expression tag	UNP Q9PPB4
F	-1	GLY	-	expression tag	UNP Q9PPB4
F	0	SER	-	expression tag	UNP Q9PPB4
F	59	ALA	HIS	engineered mutation	UNP Q9PPB4

• Molecule 2 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: $C_6H_{14}O_4$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 10 6 4	0	0




Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	В	1	Total C O	0	0
2	Б	1	10 6 4		
2	C	1	Total C O	0	0
2		10 6 4	U	U	
2	D	1	Total C O	0	0
	D	1	10 6 4	0	
2	Е	1	Total C O	0	0
	ינו	1	10 6 4	U	U
2	F	1	Total C O	0	0
	1'	1	10 6 4	U	0

• Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mg 1 1	0	0
3	С	1	Total Mg 1 1	0	0
3	E	1	Total Mg 1 1	0	0


• Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	С	1	Total C O 4 2 2	0	0
4	С	1	Total C O 4 2 2	0	0
4	D	1	Total C O 4 2 2	0	0
4	Е	1	Total C O 4 2 2	0	0
4	F	1	Total C O 4 2 2	0	0
4	F	1	Total C O 4 2 2	0	0

 \bullet Molecule 5 is LYSINE (three-letter code: LYS) (formula: $C_6H_{15}N_2O_2)$ (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C N O 10 6 2 2	0	0
5	В	1	Total C N O 10 6 2 2	0	0
5	С	1	Total C N O 10 6 2 2	0	0
5	D	1	Total C N O 10 6 2 2	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	E	1	Total 10			0	0
5	F	1	Total 10			0	0

• Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	44	Total O 44 44	0	0
6	В	39	Total O 39 39	0	0
6	С	47	Total O 47 47	0	0
6	D	43	Total O 43 43	0	0
6	Е	45	Total O 45 45	0	0
6	F	50	Total O 50 50	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 4-hydroxy-tetrahydrodipicolinate synthase









4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants	85.08Å 231.29Å 201.54Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.70 - 2.59	Depositor
resolution (A)	49.70 - 2.59	EDS
% Data completeness	99.5 (49.70-2.59)	Depositor
(in resolution range)	99.6 (49.70-2.59)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.20 (at 2.58Å)	Xtriage
Refinement program	PHENIX 1.17.1_3660	Depositor
P.P.	0.211 , 0.253	Depositor
R, R_{free}	0.211 , 0.250	DCC
R_{free} test set	3097 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	43.4	Xtriage
Anisotropy	0.343	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.35, 41.2	EDS
L-test for twinning ²	$ < L > = 0.50, < L^2> = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	13985	wwPDB-VP
Average B, all atoms $(Å^2)$	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.67% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PGE, KPI, MG, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
MIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.50	0/2276	0.54	0/3083
1	В	0.37	0/2289	0.53	0/3097
1	С	0.40	0/2282	0.56	0/3088
1	D	0.46	0/2278	0.55	0/3086
1	Е	0.38	0/2289	0.52	0/3097
1	F	0.47	0/2297	0.51	0/3109
All	All	0.43	0/13711	0.54	0/18560

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintenain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	Е	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	Е	166	KPI	Mainchain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within



the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2250	0	2264	36	0
1	В	2266	0	2294	32	0
1	С	2259	0	2285	30	0
1	D	2255	0	2263	33	1
1	Ε	2266	0	2294	24	0
1	F	2270	0	2292	31	1
2	A	10	0	14	0	0
2	В	10	0	14	1	0
2	С	10	0	14	0	0
2	D	10	0	14	0	0
2	Ε	10	0	14	0	0
2	F	10	0	14	2	0
3	A	1	0	0	0	0
3	С	1	0	0	0	0
3	Ε	1	0	0	0	0
4	A	4	0	3	0	0
4	С	8	0	6	1	0
4	D	4	0	3	1	0
4	Ε	4	0	3	0	0
4	F	8	0	6	0	0
5	A	10	0	12	1	0
5	В	10	0	12	1	0
5	С	10	0	12	1	0
5	D	10	0	12	0	0
5	Е	10	0	12	0	0
5	F	10	0	12	1	0
6	A	44	0	0	3	0
6	В	39	0	0	0	0
6	С	47	0	0	1	0
6	D	43	0	0	0	0
6	Ε	45	0	0	1	0
6	F	50	0	0	1	0
All	All	13985	0	13869	182	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 182 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:A:95:PHE:CE2	1:A:99:HIS:CD2	2.13	1.37
1:A:95:PHE:HE2	1:A:99:HIS:NE2	1.57	1.03
1:A:95:PHE:CD2	1:A:99:HIS:CD2	2.47	1.02
1:A:95:PHE:CE2	1:A:99:HIS:CG	2.56	0.93
1:A:95:PHE:HE2	1:A:99:HIS:CD2	1.71	0.86

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:D:298:PHE:O	1:F:184:ARG:NH2[3_655]	2.17	0.03

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	\mathbf{ntiles}
1	A	293/310~(94%)	283 (97%)	10 (3%)	0	100	100
1	В	293/310 (94%)	283 (97%)	10 (3%)	0	100	100
1	С	292/310 (94%)	284 (97%)	8 (3%)	0	100	100
1	D	293/310 (94%)	282 (96%)	11 (4%)	0	100	100
1	E	293/310 (94%)	284 (97%)	9 (3%)	0	100	100
1	F	294/310 (95%)	284 (97%)	9 (3%)	1 (0%)	41	64
All	All	1758/1860 (94%)	1700 (97%)	57 (3%)	1 (0%)	51	75

All (1) Ramachandran outliers are listed below:

\mathbf{Mol}	Chain	Res	Type
1	F	268	SER



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	243/259 (94%)	241 (99%)	2 (1%)	81	92
1	В	245/259 (95%)	241 (98%)	4 (2%)	62	82
1	С	244/259 (94%)	241 (99%)	3 (1%)	71	87
1	D	242/259 (93%)	239 (99%)	3 (1%)	71	87
1	${ m E}$	245/259 (95%)	241 (98%)	4 (2%)	62	82
1	F	245/259 (95%)	243 (99%)	2 (1%)	81	92
All	All	1464/1554 (94%)	1446 (99%)	18 (1%)	69	87

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Ε	32	LYS
1	F	99	HIS
1	F	26	SER
1	С	268	SER
1	Е	18	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	99	HIS
1	A	161	ASN
1	A	201	ASN
1	В	99	HIS
1	D	99	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Type	Chain	Res	Link	Во	ond leng	ths	В	ond ang	les
MIOI	Type	Chain	nes	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	KPI	A	166	1	11,13,14	1.83	2 (18%)	10,15,17	3.69	5 (50%)
1	KPI	В	166	1	11,13,14	1.85	2 (18%)	10,15,17	3.71	6 (60%)
1	KPI	Е	166	1	11,13,14	1.54	2 (18%)	10,15,17	3.15	4 (40%)
1	KPI	F	166	1	11,13,14	0.86	1 (9%)	10,15,17	3.40	5 (50%)
1	KPI	D	166	1	11,13,14	0.90	1 (9%)	10,15,17	3.46	5 (50%)
1	KPI	С	166	1	11,13,14	1.87	2 (18%)	10,15,17	3.87	6 (60%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KPI	A	166	1	-	1/13/14/16	-
1	KPI	В	166	1	-	1/13/14/16	-
1	KPI	Е	166	1	-	1/13/14/16	-
1	KPI	F	166	1	-	1/13/14/16	-
1	KPI	D	166	1	-	2/13/14/16	-
1	KPI	С	166	1	-	1/13/14/16	-

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	Ideal(A)
1	В	166	KPI	O2-CX2	5.31	1.36	1.22
1	С	166	KPI	O2-CX2	5.24	1.36	1.22
1	A	166	KPI	O2-CX2	5.18	1.36	1.22
1	Е	166	KPI	O-C	4.17	1.36	1.19
1	A	166	KPI	O1-CX2	-2.39	1.23	1.30

The worst 5 of 31 bond angle outliers are listed below:



Mol	Chain	Res	Type	$\Gamma \text{ype} \hspace{0.2cm} \hspace{0.2cm} \text{Atoms} \hspace{0.2cm} $		$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	С	166	KPI	C1-CX1-CX2	-8.55	109.86	118.17
1	D	166	KPI	C1-CX1-CX2	-8.17	110.23	118.17
1	A	166	KPI	C1-CX1-CX2	-7.59	110.79	118.17
1	F	166	KPI	C1-CX1-CX2	-7.41	110.96	118.17
1	В	166	KPI	C1-CX1-CX2	-7.26	111.11	118.17

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	D	166	KPI	C-CA-CB-CG
1	A	166	KPI	C1-CX1-NZ-CE
1	В	166	KPI	C1-CX1-NZ-CE
1	С	166	KPI	C1-CX1-NZ-CE
1	D	166	KPI	C1-CX1-NZ-CE

There are no ring outliers.

5 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	166	KPI	1	0
1	Е	166	KPI	2	0
1	F	166	KPI	1	0
1	D	166	KPI	2	0
1	С	166	KPI	1	0

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 22 ligands modelled in this entry, 3 are monoatomic - leaving 19 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Type	Chain	Res	Link	В	ond leng	$_{ m gths}$	В	ond ang	gles
IVIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	ACT	С	304	-	3,3,3	1.38	1 (33%)	3,3,3	1.37	0
5	LYS	A	304	-	8,9,9	1.11	1 (12%)	9,10,10	1.60	2 (22%)
2	PGE	Е	301	-	9,9,9	0.32	0	8,8,8	0.30	0
4	ACT	F	302	-	3,3,3	1.34	0	3,3,3	1.38	0
5	LYS	D	303	-	8,9,9	0.83	1 (12%)	9,10,10	1.28	2 (22%)
4	ACT	Е	303	-	3,3,3	1.31	0	3,3,3	1.54	0
5	LYS	В	302	-	8,9,9	0.84	1 (12%)	9,10,10	1.26	2 (22%)
4	ACT	F	303	-	3,3,3	1.23	0	3,3,3	1.37	0
4	ACT	С	303	-	3,3,3	1.34	0	3,3,3	1.52	0
4	ACT	D	302	-	3,3,3	1.36	0	3,3,3	1.49	0
2	PGE	В	301	_	9,9,9	0.31	0	8,8,8	0.23	0
5	LYS	F	304	-	8,9,9	0.82	1 (12%)	9,10,10	1.24	2 (22%)
2	PGE	A	301	-	9,9,9	0.31	0	8,8,8	0.27	0
5	LYS	С	305	-	8,9,9	0.82	1 (12%)	9,10,10	1.22	2 (22%)
2	PGE	F	301	-	9,9,9	0.30	0	8,8,8	0.38	0
5	LYS	Е	304	-	8,9,9	0.82	1 (12%)	9,10,10	1.12	1 (11%)
2	PGE	С	301	-	9,9,9	0.33	0	8,8,8	0.20	0
2	PGE	D	301	-	9,9,9	0.31	0	8,8,8	0.25	0
4	ACT	A	303	-	3,3,3	1.22	0	3,3,3	1.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PGE	С	301	-	-	1/7/7/7	-
2	PGE	В	301	ı	ı	0/7/7/7	-
2	PGE	D	301	-	-	3/7/7/7	-
5	LYS	D	303	-	-	2/9/9/9	-
5	LYS	F	304	-	-	0/9/9/9	-
5	LYS	A	304	ı	-	4/9/9/9	-
2	PGE	Е	301	-	-	2/7/7/7	-
2	PGE	A	301	-	-	1/7/7/7	-
5	LYS	В	302	-	-	3/9/9/9	-
5	LYS	С	305	-	-	3/9/9/9	_
2	PGE	F	301	-	-	6/7/7/7	_
5	LYS	Е	304	-	-	2/9/9/9	-



TT1	_	. C	$\overline{}$	1 . 1	1 1			1 1	1 1 .
The worst 5)	OŤ.	(bond	length	outliers	are	listed	below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$Ideal(\AA)$
5	A	304	LYS	OXT-C	-2.84	1.21	1.30
5	В	302	LYS	OXT-C	-2.17	1.23	1.30
5	Е	304	LYS	OXT-C	-2.14	1.23	1.30
5	F	304	LYS	OXT-C	-2.13	1.23	1.30
5	С	305	LYS	OXT-C	-2.13	1.23	1.30

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$Ideal(^{o})$
5	A	304	LYS	OXT-C-O	-3.73	115.61	124.09
5	В	302	LYS	OXT-C-O	-2.96	117.37	124.09
5	С	305	LYS	OXT-C-O	-2.79	117.76	124.09
5	D	303	LYS	OXT-C-O	-2.76	117.81	124.09
5	F	304	LYS	OXT-C-O	-2.72	117.91	124.09

There are no chirality outliers.

5 of 27 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	304	LYS	C-CA-CB-CG
5	D	303	LYS	C-CA-CB-CG
2	A	301	PGE	O1-C1-C2-O2
2	F	301	PGE	O3-C5-C6-O4
5	A	304	LYS	CA-CB-CG-CD

There are no ring outliers.

8 monomers are involved in 9 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	304	LYS	1	0
5	В	302	LYS	1	0
4	С	303	ACT	1	0
4	D	302	ACT	1	0
2	В	301	PGE	1	0
5	F	304	LYS	1	0
5	С	305	LYS	1	0
2	F	301	PGE	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will





also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





























































5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$	$OWAB(Å^2)$	Q<0.9
1	A	$294/310\ (94\%)$	0.17	3 (1%) 82 80	33, 43, 60, 71	0
1	В	$295/310\ (95\%)$	0.20	4 (1%) 75 71	32, 40, 55, 67	0
1	С	294/310~(94%)	0.22	7 (2%) 59 53	33, 41, 60, 72	0
1	D	$295/310\ (95\%)$	0.31	6 (2%) 65 60	34, 42, 60, 73	0
1	E	$295/310\ (95\%)$	0.24	2 (0%) 87 86	34, 40, 55, 67	0
1	F	295/310~(95%)	0.27	3 (1%) 82 80	34, 43, 60, 71	0
All	All	1768/1860 (95%)	0.23	25 (1%) 75 71	32, 42, 59, 73	0

The worst 5 of 25 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	4	ASN	4.0
1	С	268	SER	3.3
1	F	95[A]	PHE	3.3
1	В	98	GLU	3.0
1	С	60	ARG	3.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathring{\mathbf{A}}^2)$	Q < 0.9
1	KPI	В	166	14/15	0.91	0.20	31,34,40,40	0
1	KPI	D	166	14/15	0.91	0.21	39,42,44,45	0
1	KPI	С	166	14/15	0.93	0.17	34,35,46,48	0
1	KPI	A	166	14/15	0.93	0.18	38,39,43,45	0



Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
1	KPI	Е	166	14/15	0.94	0.15	35,38,43,43	0
1	KPI	F	166	14/15	0.95	0.16	37,39,44,46	0

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathbf{A}}^2)$	Q < 0.9
3	MG	A	302	1/1	0.71	0.14	50,50,50,50	0
2	PGE	F	301	10/10	0.75	0.42	50,55,58,60	0
4	ACT	С	304	4/4	0.78	0.27	44,48,49,51	0
2	PGE	С	301	10/10	0.85	0.22	40,41,43,44	0
3	MG	E	302	1/1	0.87	0.31	48,48,48,48	0
2	PGE	Е	301	10/10	0.89	0.16	41,44,49,51	0
4	ACT	D	302	4/4	0.89	0.20	46,46,49,51	0
5	LYS	D	303	10/10	0.89	0.21	43,47,50,54	0
5	LYS	В	302	10/10	0.90	0.16	40,44,49,50	0
5	LYS	A	304	10/10	0.91	0.24	20,20,20,20	0
2	PGE	В	301	10/10	0.91	0.21	42,45,53,54	0
4	ACT	F	302	4/4	0.91	0.15	44,44,52,52	0
4	ACT	A	303	4/4	0.92	0.19	38,42,45,46	0
5	LYS	С	305	10/10	0.92	0.19	40,46,53,53	0
4	ACT	E	303	4/4	0.92	0.15	41,42,44,44	0
5	LYS	F	304	10/10	0.92	0.20	44,46,52,57	0
2	PGE	A	301	10/10	0.93	0.18	$44,\!49,\!51,\!52$	0
2	PGE	D	301	10/10	0.93	0.17	44,46,50,51	0
5	LYS	E	304	10/10	0.94	0.19	40,44,50,52	0
4	ACT	С	303	4/4	0.95	0.18	42,45,46,50	0
4	ACT	F	303	4/4	0.96	0.13	45,48,48,49	0
3	MG	С	302	1/1	0.97	0.13	44,44,44,44	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different




orientation to approximate a three-dimensional view.





















Electron density around MG E 302:


 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)




































Electron density around ACT F 302: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)




Electron density around ACT A 303: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)





















Electron density around PGE D 301:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around LYS E 304:


 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)



















Electron density around ACT F 303:


 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)











6.5 Other polymers (i)

There are no such residues in this entry.

