



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 9, 2021 – 06:11 PM EST

PDB ID : 7KV4
Title : Surface glycan-binding protein A from Bacteroides fluxus
Authors : Tamura, K.; Brumer, H.; Van Petegem, F.
Deposited on : 2020-11-26
Resolution : 1.84 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.17
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.17


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4003 (1.86-1.82)
Clashscore	141614	4233 (1.86-1.82)
Ramachandran outliers	138981	4185 (1.86-1.82)
Sidechain outliers	138945	4186 (1.86-1.82)
RSRZ outliers	127900	3957 (1.86-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	510	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7333 atoms, of which 3448 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SusD family protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	AAA	465	7092	2293	3448	624	707	20	198	0	0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	AAA	241	Total	O	0	0
			241	241		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: SusD family protein



4 Data and refinement statistics i

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	63.85Å 116.50Å 139.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.43 – 1.84 36.40 – 1.84	Depositor EDS
% Data completeness (in resolution range)	99.8 (36.43-1.84) 99.9 (36.40-1.84)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.46 (at 1.84Å)	Xtrriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.204 , 0.240 0.211 , 0.247	Depositor DCC
R_{free} test set	2369 reflections (5.20%)	wwPDB-VP
Wilson B-factor (Å ²)	35.4	Xtrriage
Anisotropy	0.192	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 43.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.016 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.031 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7333	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	AAA	0.76	2/3721 (0.1%)	0.88	4/5026 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	AAA	269	GLU	CD-OE2	9.61	1.36	1.25
1	AAA	269	GLU	CB-CG	5.16	1.61	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AAA	227	ARG	CG-CD-NE	7.25	127.02	111.80
1	AAA	269	GLU	OE1-CD-OE2	-5.39	116.83	123.30
1	AAA	269	GLU	CG-CD-OE2	5.33	128.97	118.30
1	AAA	372	PRO	N-CA-CB	-5.21	96.87	102.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	3644	3448	3412	21	0
2	AAA	241	0	0	1	0
All	All	3885	3448	3412	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:85:GLY:O	1:AAA:353:ARG:HB3	2.08	0.54
1:AAA:295:GLY:HA3	1:AAA:380:ASN:HA	1.96	0.47
1:AAA:62:TYR:CE1	1:AAA:65:ASN:HA	2.50	0.46
1:AAA:327:ALA:O	1:AAA:330:MET:HG2	2.16	0.46
1:AAA:488:PRO:HG3	1:AAA:510:TYR:CD1	2.51	0.45
1:AAA:369:GLY:C	1:AAA:371:ASN:H	2.19	0.45
1:AAA:194:LEU:HD13	1:AAA:225:LYS:HA	1.99	0.43
1:AAA:260:PHE:O	1:AAA:263:MET:HG2	2.18	0.43
1:AAA:137:GLU:OE1	1:AAA:137:GLU:HA	2.18	0.43
1:AAA:406:VAL:HG12	1:AAA:407:ALA:O	2.18	0.42
1:AAA:82:TYR:O	1:AAA:321:ALA:HA	2.20	0.42
1:AAA:318:TRP:CD1	1:AAA:318:TRP:N	2.87	0.42
1:AAA:273:GLU:HG2	1:AAA:389:ARG:HB2	2.01	0.42
1:AAA:56:ILE:HG13	1:AAA:118:GLY:HA2	2.01	0.42
1:AAA:305:ASN:HA	1:AAA:317:GLY:O	2.19	0.41
1:AAA:361:GLN:HB3	2:AAA:798:HOH:O	2.20	0.41
1:AAA:42:THR:HG23	1:AAA:45:THR:HG22	2.01	0.41
1:AAA:84:GLY:HA2	1:AAA:354:PHE:O	2.21	0.41
1:AAA:69:SER:O	1:AAA:73:MET:HG2	2.20	0.40
1:AAA:42:THR:HG23	1:AAA:45:THR:H	1.87	0.40
1:AAA:321:ALA:N	1:AAA:322:PRO:CD	2.84	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	AAA	459/510 (90%)	438 (95%)	21 (5%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	AAA	366/418 (88%)	361 (99%)	5 (1%)	67 55

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	42	THR
1	AAA	156	HIS
1	AAA	266	ASP
1	AAA	428	ILE
1	AAA	430	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	465/510 (91%)	0.47	51 (10%) 5 4	22, 41, 71, 90	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	499	GLY	5.1
1	AAA	71	LEU	4.3
1	AAA	46	ILE	4.1
1	AAA	310	PRO	3.9
1	AAA	502	PHE	3.6
1	AAA	45	THR	3.6
1	AAA	41	ALA	3.5
1	AAA	312	GLY	3.4
1	AAA	498	LYS	3.4
1	AAA	135	VAL	3.4
1	AAA	130	ALA	3.4
1	AAA	503	GLU	3.2
1	AAA	491	LEU	3.2
1	AAA	455	TYR	3.0
1	AAA	131	ASN	3.0
1	AAA	42	THR	2.9
1	AAA	171	PRO	2.9
1	AAA	177	MET	2.8
1	AAA	388	PHE	2.8
1	AAA	178	THR	2.8
1	AAA	120	ALA	2.7
1	AAA	313	VAL	2.6
1	AAA	138	ARG	2.6
1	AAA	48	SER	2.6
1	AAA	492	ASN	2.6
1	AAA	495	GLU	2.6
1	AAA	387	ILE	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	AAA	448	LEU	2.5
1	AAA	228	VAL	2.5
1	AAA	170	GLU	2.5
1	AAA	368	GLU	2.5
1	AAA	209	SER	2.4
1	AAA	72	LEU	2.4
1	AAA	128	ALA	2.4
1	AAA	501	GLU	2.3
1	AAA	166	PRO	2.3
1	AAA	69	SER	2.3
1	AAA	176	TYR	2.2
1	AAA	211	ASP	2.3
1	AAA	347	GLY	2.2
1	AAA	258	ASP	2.2
1	AAA	161	PHE	2.2
1	AAA	494	ILE	2.1
1	AAA	449	GLY	2.1
1	AAA	157	LEU	2.1
1	AAA	452	MET	2.1
1	AAA	385	LEU	2.1
1	AAA	497	THR	2.0
1	AAA	133	GLU	2.0
1	AAA	447	PHE	2.0
1	AAA	496	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.