

Full wwPDB X-ray Structure Validation Report (i)

Nov 5, 2023 – 09:03 AM EST

PDB ID	:	4KX8
Title	:	Crystal structure of human aminopeptidase A complexed with amastatin
Authors	:	Yang, Y.; Liu, C.; Lin, Y.Y.; Li, F.
Deposited on	:	2013-05-24
Resolution	:	2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Motria	Whole archive	Similar resolution		
	$(\# {\rm Entries})$	$(\# { m Entries, resolution range}({ m \AA}))$		
R _{free}	130704	3907 (2.40-2.40)		
Clashscore	141614	4398 (2.40-2.40)		
Ramachandran outliers	138981	4318 (2.40-2.40)		
Sidechain outliers	138945	4319 (2.40-2.40)		
RSRZ outliers	127900	3811 (2.40-2.40)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
1	Δ	888	<u>2</u> %	010/		150/		
1	Λ	000	25%	81%		15% ••		
2	С	4	25%	25%	25%	25%		
3	В	2		10	0%			
3	F	2		10	0%			
3	G	2		10	0%			



Mol	Chain	Length	Quality of chain
4	D	3	100%
4	Е	3	100%



4KX8

2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 7942 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Glutamyl aminopeptidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	А	875	Total 7151	C 4580	N 1190	O 1356	$\begin{array}{c} \mathrm{S} \\ \mathrm{25} \end{array}$	0	1	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	213	ARG	GLN	variant	UNP Q07075
А	218	ALA	VAL	variant	UNP Q07075
А	958	HIS	-	expression tag	UNP Q07075
А	959	HIS	-	expression tag	UNP Q07075
А	960	HIS	-	expression tag	UNP Q07075
А	961	HIS	-	expression tag	UNP Q07075
A	962	HIS	-	expression tag	UNP Q07075
А	963	HIS	-	expression tag	UNP Q07075

• Molecule 2 is a protein called amastatin.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	С	4	Total 33	C 21	N 4	0 8	0	0	0

• Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
3	В	2	Total C N O 28 16 2 10	0	0	0



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
3	F	2	Total C N O 28 16 2 10	0	0	0
3	G	2	Total C N O 28 16 2 10	0	0	0

• Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace
4	D	3	Total C N 42 24 3	NO 815	0	0	0
4	Е	3	Total C N 42 24 3	N O B 15	0	0	0

• Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	1	Total Zn 1 1	0	0

• Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	А	1	Total C N O 14 8 1 5	0	0
6	А	1	Total C N O 14 8 1 5	0	0
6	А	1	Total C N O 14 8 1 5	0	0
6	А	1	Total C N O 14 8 1 5	0	0

• Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	А	531	Total O 531 531	0	0
7	С	2	Total O 2 2	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Glutamyl aminopeptidase

• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B:

100%

NAG1 NAG2

• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



Chain F:	100%
NAG1 NAG2	
• Molecule 3: 2-acetamido-2-deoxy-beta opyranose	a-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-gluc
Chain G:	100%
NAG 2 NAG 2	
• Molecule 4: 2-acetamido-2-deoxy-beta opyranose-(1-4)-2-acetamido-2-deoxy-be	a-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-gluc eta-D-glucopyranose
Chain D:	100%
NA G1 NA G2 NA G3	
• Molecule 4: 2-acetamido-2-deoxy-beta opyranose-(1-4)-2-acetamido-2-deoxy-be	a-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-gluc eta-D-glucopyranose

Chain E:

100%

NAG1 NAG2 NAG3 NAG3



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 64 2 2	Depositor
Cell constants	142.74Å 142.74Å 237.81Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
$Resolution(\AA)$	41.21 - 2.40	Depositor
Resolution (A)	41.21 - 2.40	EDS
% Data completeness	95.0 (41.21-2.40)	Depositor
(in resolution range)	95.0 (41.21-2.40)	EDS
R _{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$6.20 (at 2.39 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
D D .	0.210 , 0.276	Depositor
n, n_{free}	0.214 , 0.273	DCC
R_{free} test set	2713 reflections (5.05%)	wwPDB-VP
Wilson B-factor $(Å^2)$	34.5	Xtriage
Anisotropy	0.613	Xtriage
Bulk solvent $k_{sol}(e/A^3), B_{sol}(A^2)$	0.32 , 56.3	EDS
L-test for $twinning^2$	$ < L >=0.53, < L^2>=0.37$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7942	wwPDB-VP
Average B, all atoms $(Å^2)$	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.62% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: L2O, NAG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bo	nd lengths	Bond angles		
		RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.45	1/7336~(0.0%)	0.61	1/9974~(0.0%)	
2	С	1.51	0/22	1.72	0/28	
All	All	0.45	1/7358~(0.0%)	0.61	1/10002~(0.0%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	С	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\mathrm{Ideal}(\mathrm{\AA})$
1	А	158	ARG	CG-CD	5.11	1.64	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	А	903	LEU	CA-CB-CG	5.30	127.49	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	С	1	L2O	Peptide



5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	7151	0	6943	97	0
2	С	33	0	29	5	0
3	В	28	0	25	1	0
3	F	28	0	25	0	0
3	G	28	0	25	0	0
4	D	42	0	37	1	0
4	Е	42	0	37	0	0
5	А	1	0	0	0	0
6	А	56	0	52	2	0
7	А	531	0	0	18	0
7	C	2	0	0	0	0
All	All	7942	0	7173	99	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (99) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (\AA)	overlap (Å)
1:A:907:GLN:HB3	7:A:1482:HOH:O	1.43	1.16
1:A:244:THR:HB	7:A:1595:HOH:O	1.62	0.98
1:A:175:GLU:O	1:A:176:GLU:HB3	1.68	0.94
1:A:761:LEU:HD22	1:A:793:ASN:HD21	1.41	0.83
1:A:423:GLU:O	1:A:427:VAL:HG12	1.81	0.81
1:A:356:THR:HG21	2:C:1:L2O:H8	1.68	0.75
1:A:793:ASN:HB3	7:A:1501:HOH:O	1.90	0.72
1:A:501:GLY:HA3	1:A:523:LEU:HD23	1.72	0.72
1:A:736:ALA:O	1:A:745:ARG:NH1	2.20	0.70
1:A:120:SER:OG	1:A:189:THR:HG22	1.92	0.70
1:A:146:GLU:HA	7:A:1563:HOH:O	1.91	0.70
1:A:834:ARG:HD2	7:A:1612:HOH:O	1.91	0.70
1:A:100:VAL:O	1:A:242:THR:HG22	1.92	0.69
1:A:386:ARG:NH2	2:C:4:ASP:OD1	2.26	0.68
1:A:118:THR:HG22	1:A:191:GLU:HG2	1.74	0.68
1:A:943:ARG:HH12	1:A:947:ARG:HH11	1.42	0.68



	louo pugom	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:674:LYS:HD3	4:D:1:NAG:H62	1.77	0.67
1:A:703:ILE:HD12	1:A:713:TYR:CE2	2.29	0.67
1:A:511:PHE:CZ	7:A:1411:HOH:O	2.47	0.66
6:A:1004:NAG:H83	7:A:1591:HOH:O	1.98	0.64
1:A:244:THR:HG22	1:A:278:LYS:HA	1.79	0.64
1:A:735:ASP:OD1	1:A:781:ASN:HB2	1.98	0.63
1:A:101:HIS:CD2	1:A:242:THR:HG23	2.33	0.62
1:A:224:PRO:HD2	7:A:1241:HOH:O	2.01	0.60
1:A:137:ARG:NH2	1:A:226:ASP:OD1	2.34	0.60
1:A:103:ASP:HB2	1:A:120:SER:HB2	1.84	0.59
1:A:923:LYS:HG3	7:A:1409:HOH:O	2.02	0.58
1:A:869:ASN:HB2	7:A:1515:HOH:O	2.03	0.58
1:A:597:ASN:HD22	6:A:1017:NAG:H83	1.68	0.57
1:A:552:VAL:HG12	1:A:617:LEU:HD12	1.86	0.56
1:A:703:ILE:CD1	1:A:713:TYR:CE2	2.89	0.56
1:A:717:GLU:HG2	1:A:755:MET:O	2.06	0.56
1:A:119:VAL:O	1:A:189:THR:HA	2.06	0.55
1:A:175:GLU:CG	1:A:176:GLU:N	2.71	0.54
1:A:689:PHE:HA	1:A:744:LEU:HD13	1.90	0.54
1:A:713:TYR:HB3	1:A:714:PRO:HD3	1.90	0.53
1:A:365:ILE:HG22	1:A:367:TYR:CE1	2.44	0.53
1:A:177:LEU:CD1	1:A:178:THR:H	2.23	0.52
1:A:565:ARG:CZ	1:A:937:GLU:HG3	2.40	0.52
1:A:128:PRO:HB3	1:A:176:GLU:OE2	2.11	0.51
1:A:273:ARG:HD2	7:A:1508:HOH:O	2.09	0.51
1:A:402:ASN:N	1:A:402:ASN:HD22	2.07	0.51
1:A:278:LYS:HB2	7:A:1596:HOH:O	2.10	0.51
1:A:616:PHE:HE2	1:A:618:LYS:HG3	1.76	0.51
1:A:197:ASN:O	3:B:1:NAG:H82	2.10	0.51
1:A:202:GLY:O	1:A:220:THR:HA	2.11	0.50
1:A:131:TYR:CD2	1:A:172:GLU:HB3	2.46	0.50
1:A:397:HIS:HA	1:A:400:PHE:O	2.12	0.50
1:A:341:TYR:HA	7:A:1483:HOH:O	2.12	0.50
1:A:131:TYR:CE2	1:A:172:GLU:HB3	2.47	0.49
1:A:361:ASN:HB2	1:A:364:LEU:O	2.13	0.49
1:A:511:PHE:HZ	7:A:1411:HOH:O	1.92	0.49
1:A:175:GLU:O	1:A:176:GLU:CB	2.49	0.49
1:A:174:GLU:O	1:A:175:GLU:CB	2.61	0.48
1:A:493:ILE:O	1:A:494:LYS:HB2	2.14	0.48
1:A:585:TRP:CD1	1:A:617:LEU:HD23	2.47	0.48
1:A:175:GLU:CG	1:A:176:GLU:H	2.26	0.48



		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:356:THR:HG21	2:C:1:L2O:H12	1.96	0.48
1:A:493:ILE:HG13	1:A:529:LEU:HD12	1.95	0.47
1:A:461:ILE:HD12	1:A:539:THR:HG23	1.95	0.47
1:A:460:ILE:HD11	1:A:481:LYS:HG2	1.96	0.47
1:A:145:PRO:HB3	1:A:190:MET:HB3	1.97	0.47
1:A:101:HIS:HD2	1:A:242:THR:HG23	1.78	0.47
1:A:943:ARG:HH22	1:A:947:ARG:NH1	2.13	0.47
1:A:411:ASP:OD1	1:A:516:THR:OG1	2.31	0.46
1:A:348:LYS:HZ3	1:A:365:ILE:HD13	1.79	0.46
1:A:177:LEU:HD12	1:A:178:THR:H	1.81	0.46
1:A:348:LYS:HE3	1:A:391:VAL:HG13	1.97	0.46
1:A:360:GLU:O	1:A:361:ASN:C	2.52	0.46
1:A:863:GLY:HA2	1:A:866:MET:HB2	1.97	0.46
1:A:259:MET:SD	1:A:279:SER:HA	2.56	0.46
1:A:348:LYS:NZ	1:A:365:ILE:HD13	2.32	0.45
1:A:527:SER:C	1:A:529:LEU:H	2.19	0.45
1:A:175:GLU:HG2	1:A:176:GLU:H	1.82	0.45
1:A:588:ASP:O	1:A:589:ASN:HB2	2.15	0.45
1:A:77:ILE:HD11	1:A:144:LEU:N	2.32	0.45
7:A:1264:HOH:O	2:C:3:VAL:HG13	2.17	0.45
1:A:177:LEU:HB3	1:A:186:TYR:OH	2.18	0.44
1:A:86:GLY:HA3	1:A:88:TRP:CZ3	2.53	0.44
1:A:371:ASN:HD21	2:C:4:ASP:CG	2.21	0.44
1:A:105:HIS:HE1	1:A:248:THR:OG1	2.00	0.44
1:A:360:GLU:HG3	1:A:397:HIS:HB3	1.99	0.44
1:A:510:GLN:HE21	1:A:510:GLN:HB2	1.65	0.43
1:A:407:ASP:HB3	7:A:1603:HOH:O	2.18	0.43
1:A:596:PHE:C	1:A:596:PHE:CD1	2.92	0.43
1:A:175:GLU:HG3	1:A:176:GLU:N	2.34	0.43
1:A:174:GLU:O	1:A:175:GLU:CG	2.67	0.42
1:A:788:ARG:HD2	7:A:1141:HOH:O	2.20	0.42
1:A:222:HIS:HA	1:A:226:ASP:HB2	2.01	0.42
1:A:666:ALA:HA	1:A:671:LEU:O	2.20	0.42
1:A:415:ASN:N	1:A:415:ASN:HD22	2.18	0.41
1:A:806:GLN:NE2	7:A:1328:HOH:O	2.53	0.41
1:A:858:SER:O	1:A:864:LYS:HA	2.20	0.41
1:A:175:GLU:HG3	1:A:176:GLU:O	2.20	0.41
1:A:923:LYS:HD2	1:A:923:LYS:HA	1.85	0.41
1:A:797:GLU:HG3	1:A:831:LEU:HD21	2.03	0.40
1:A:149:ARG:CZ	1:A:155:VAL:HG22	2.51	0.40
1:A:174:GLU:O	1:A:175:GLU:HG2	2.21	0.40



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:120:SER:CB	1:A:189:THR:HG22	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	872/888~(98%)	838~(96%)	29~(3%)	5 (1%)	25 36
2	С	2/4~(50%)	2(100%)	0	0	100 100
All	All	874/892~(98%)	840 (96%)	29~(3%)	5 (1%)	25 36

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	175	GLU
1	А	176	GLU
1	А	613	GLY
1	А	494	LYS
1	А	552	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Analysed Rotameric		Percentiles
Mol	Chain	Analysed	Botameric	Outliers	Percentiles
1					17 00
1	А	(86/ (97 (99%))	736 (94%)	50 (6%)	17 28
2	\mathbf{C}	3/3~(100%)	2~(67%)	1 (33%)	0
All	All	789/800~(99%)	738 (94%)	51 (6%)	17 27

All (51) residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
1	А	77	ILE
1	А	81	SER
1	А	84	GLU
1	А	95	ASP
1	А	98	ASN
1	А	132	LEU
1	А	136	LEU
1	А	148	LYS
1	А	156	GLN
1	А	177	LEU
1	А	183	ASP
1	А	190	MET
1	А	200	LEU
1	А	278	LYS
1	А	298	LYS
1	А	303	SER
1	А	345	LYS
1	А	356	THR
1	А	362	TRP
1	А	394	GLU
1	А	402	ASN
1	А	406	MET
1	А	427	VAL
1	А	462	VAL
1	А	493	ILE
1	А	508	LYS
1	А	510	GLN
1	А	524	GLU
1	A	527	SER
1	A	528	ARG
1	А	539	THR
1	А	575	LEU



Mol	Chain	Res	Type
1	А	590	ILE
1	А	596	PHE
1	А	605	THR
1	А	617	LEU
1	А	658	LEU
1	А	665	LEU
1	А	677	LEU
1	А	703	ILE
1	А	724	VAL
1	А	738	ASP
1	А	748	VAL
1	А	765	SER
1	А	778	LEU
1	А	794	SER
1	А	821	TYR
1	А	844	LEU
1	А	903	LEU
1	А	953	LEU
2	С	4	ASP

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such side chains are listed below:

Mol	Chain	Res	Type
1	А	105	HIS
1	А	402	ASN
1	А	415	ASN
1	А	510	GLN
1	А	570	GLN
1	А	792	GLN
1	А	793	ASN
1	А	843	ASN
1	А	874	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.



In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type Cha	Type	Chain	Dog	Tink	B	ond leng	gths	E	Bond ang	gles
	Ullalli	Chann Res		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2	
2	L2O	С	1	5,2	7,9,10	0.98	1 (14%)	6,11,13	2.26	3 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	L2O	С	1	5,2	-	5/9/10/12	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	Observed(Å)	$\mathrm{Ideal}(\mathrm{\AA})$
2	С	1	L2O	C2-CA	2.00	1.56	1.53

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	С	1	L2O	O1-C6-CA	4.04	113.73	107.31
2	С	1	L2O	O1-C6-C	-3.09	102.79	110.08
2	С	1	L2O	C3-C2-CA	-2.11	111.28	115.82

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	С	1	L2O	C-C6-CA-N
2	С	1	L2O	C-C6-CA-C2
2	С	1	L2O	CA-C2-C3-C4
2	С	1	L2O	CA-C2-C3-C5
2	С	1	L2O	O1-C6-CA-C2

There are no ring outliers.



1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	С	1	L2O	2	0

5.5 Carbohydrates (i)

12 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Tuno	Chain	Dog	Tink	Bo	ond leng	$_{\rm ths}$	Bond angles			
	Type	Ullalli	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
3	NAG	В	1	1,3	14,14,15	0.50	0	17,19,21	0.82	0	
3	NAG	В	2	3	14,14,15	0.46	0	17,19,21	0.94	1 (5%)	
4	NAG	D	1	4,1	14,14,15	0.51	0	17,19,21	1.10	0	
4	NAG	D	2	4	$14,\!14,\!15$	0.61	0	17,19,21	0.96	1(5%)	
4	NAG	D	3	4	14,14,15	0.51	0	17,19,21	1.24	2 (11%)	
4	NAG	Е	1	4,1	14,14,15	0.54	0	17,19,21	1.38	2 (11%)	
4	NAG	Е	2	4	14,14,15	0.65	0	17,19,21	2.17	7 (41%)	
4	NAG	Е	3	4	14,14,15	1.39	1 (7%)	17,19,21	1.97	4 (23%)	
3	NAG	F	1	1,3	14,14,15	0.59	0	17,19,21	1.17	2 (11%)	
3	NAG	F	2	3	14,14,15	0.46	0	17,19,21	1.01	1 (5%)	
3	NAG	G	1	1,3	14,14,15	0.60	0	17,19,21	0.62	0	
3	NAG	G	2	3	14,14,15	0.50	0	17,19,21	0.97	0	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	В	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	В	2	3	-	1/6/23/26	0/1/1/1
4	NAG	D	1	4,1	-	2/6/23/26	0/1/1/1
4	NAG	D	2	4	-	1/6/23/26	0/1/1/1
4	NAG	D	3	4	-	0/6/23/26	0/1/1/1



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	Е	1	4,1	-	1/6/23/26	0/1/1/1
4	NAG	Е	2	4	-	1/6/23/26	0/1/1/1
4	NAG	Е	3	4	-	4/6/23/26	0/1/1/1
3	NAG	F	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	F	2	3	-	3/6/23/26	0/1/1/1
3	NAG	G	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	G	2	3	-	2/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	Observed(Å)	Ideal(Å)
4	Ε	3	NAG	O6-C6	4.13	1.59	1.42

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
4	Е	2	NAG	O5-C1-C2	-4.92	103.53	111.29
4	Е	3	NAG	C4-C3-C2	4.89	118.18	111.02
4	Е	1	NAG	C1-O5-C5	4.14	117.80	112.19
4	Е	2	NAG	C1-O5-C5	3.72	117.23	112.19
4	Е	3	NAG	C2-N2-C7	3.50	127.89	122.90
4	Е	3	NAG	C3-C4-C5	3.39	116.29	110.24
4	Е	2	NAG	C2-N2-C7	3.34	127.66	122.90
4	D	2	NAG	O5-C1-C2	-2.77	106.92	111.29
4	Е	3	NAG	C1-O5-C5	2.75	115.91	112.19
3	F	1	NAG	O5-C1-C2	-2.73	106.97	111.29
4	D	3	NAG	C1-O5-C5	2.60	115.71	112.19
3	F	1	NAG	C4-C3-C2	2.54	114.74	111.02
4	Е	2	NAG	C1-C2-N2	2.53	114.81	110.49
3	F	2	NAG	C2-N2-C7	2.39	126.30	122.90
4	Е	2	NAG	O4-C4-C3	2.36	115.82	110.35
3	В	2	NAG	O5-C1-C2	-2.28	107.69	111.29
4	D	3	NAG	O5-C5-C6	2.25	110.73	107.20
4	Е	2	NAG	O5-C5-C4	2.22	116.24	110.83
4	Е	2	NAG	C4-C3-C2	-2.15	107.86	111.02
4	Е	1	NAG	O5-C1-C2	-2.11	107.96	111.29

There are no chirality outliers.

All (17) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
4	Е	3	NAG	O5-C5-C6-O6
4	Е	3	NAG	C1-C2-N2-C7
4	Е	3	NAG	C4-C5-C6-O6
4	D	1	NAG	C8-C7-N2-C2
4	D	1	NAG	O7-C7-N2-C2
3	G	2	NAG	O5-C5-C6-O6
3	F	2	NAG	O5-C5-C6-O6
3	F	2	NAG	C4-C5-C6-O6
4	D	2	NAG	C4-C5-C6-O6
3	G	2	NAG	C4-C5-C6-O6
3	В	2	NAG	O5-C5-C6-O6
3	F	2	NAG	C3-C2-N2-C7
4	Е	2	NAG	C3-C2-N2-C7
4	Е	3	NAG	C3-C2-N2-C7
4	Е	1	NAG	C4-C5-C6-O6
3	F	1	NAG	C4-C5-C6-O6
3	F	1	NAG	O5-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	1	NAG	1	0
3	В	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

























5.6 Ligand geometry (i)

Torsions

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Rings

Mol Type Ch	Chain	Dec	Tiple	Bo	ond leng	$_{\rm sths}$	Bond angles			
MOI	туре	Unain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	NAG	А	1005	1	14,14,15	0.58	0	17,19,21	1.24	1 (5%)
6	NAG	А	1017	1	14,14,15	0.53	0	17,19,21	1.34	4 (23%)
6	NAG	А	1004	1	14,14,15	0.52	0	17,19,21	1.59	4 (23%)
6	NAG	А	1006	1	14,14,15	0.51	0	17,19,21	1.88	5 (29%)



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	А	1005	1	-	1/6/23/26	0/1/1/1
6	NAG	А	1017	1	-	3/6/23/26	0/1/1/1
6	NAG	А	1004	1	-	3/6/23/26	0/1/1/1
6	NAG	А	1006	1	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
6	А	1006	NAG	C2-N2-C7	5.18	130.28	122.90
6	А	1005	NAG	C1-O5-C5	4.44	118.21	112.19
6	А	1004	NAG	C1-O5-C5	3.64	117.13	112.19
6	А	1004	NAG	O7-C7-C8	-2.88	116.70	122.06
6	А	1004	NAG	C8-C7-N2	2.67	120.62	116.10
6	А	1017	NAG	C1-O5-C5	2.32	115.34	112.19
6	А	1017	NAG	C1-C2-N2	2.29	114.39	110.49
6	А	1006	NAG	C1-O5-C5	2.27	115.27	112.19
6	А	1006	NAG	C1-C2-N2	2.27	114.37	110.49
6	А	1004	NAG	C2-N2-C7	2.26	126.12	122.90
6	А	1006	NAG	C3-C4-C5	2.23	114.22	110.24
6	А	1006	NAG	O5-C5-C6	2.21	110.66	107.20
6	А	1017	NAG	C2-N2-C7	2.16	125.98	122.90
6	A	1017	NAG	O5-C5-C6	2.16	110.59	107.20

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	А	1004	NAG	C8-C7-N2-C2
6	А	1004	NAG	O7-C7-N2-C2
6	А	1017	NAG	C8-C7-N2-C2
6	А	1017	NAG	O7-C7-N2-C2
6	А	1017	NAG	O5-C5-C6-O6
6	А	1005	NAG	O5-C5-C6-O6
6	А	1004	NAG	O5-C5-C6-O6
6	А	1006	NAG	C3-C2-N2-C7
6	А	1006	NAG	C4-C5-C6-O6



There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	А	1017	NAG	1	0
6	А	1004	NAG	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	875/888~(98%)	-0.10	22 (2%) 57 55	20, 51, 97, 139	0
2	С	3/4~(75%)	1.26	1 (33%) 0 0	22, 22, 31, 57	0
All	All	878/892~(98%)	-0.10	23 (2%) 56 54	20, 51, 97, 139	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	183	ASP	5.4
1	А	184	GLY	5.3
1	А	615	ALA	4.3
1	А	182	GLY	3.7
1	А	175	GLU	3.5
1	А	181	SER	3.5
1	А	185	LEU	3.4
1	А	128	PRO	3.4
1	А	173	ALA	3.3
1	А	76	ASP	2.9
1	А	84	GLU	2.9
1	А	528	ARG	2.9
1	А	951	PHE	2.8
1	А	151	SER	2.8
1	А	304	GLY	2.8
1	А	602	GLU	2.7
1	А	530	PRO	2.7
1	А	180	SER	2.6
1	А	599	SER	2.5
2	С	4	ASP	2.3
1	А	80	ALA	2.2
1	А	82	GLU	2.2
1	А	614	ASN	2.1



6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
2	L2O	C	1	10/11	0.85	0.23	13,16,17,18	0

6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
3	NAG	F	2	14/15	0.77	0.36	106,120,137,139	0
3	NAG	В	2	14/15	0.84	0.36	106,135,149,150	0
4	NAG	D	3	14/15	0.84	0.26	113,148,164,177	0
4	NAG	Е	3	14/15	0.84	0.23	$95,\!137,\!155,\!160$	0
4	NAG	Е	1	14/15	0.87	0.17	61,82,94,130	0
3	NAG	G	2	14/15	0.88	0.29	100,117,131,135	0
3	NAG	В	1	14/15	0.88	0.16	$61,\!85,\!105,\!112$	0
4	NAG	Е	2	14/15	0.90	0.22	96,110,126,129	0
4	NAG	D	2	14/15	0.90	0.20	112,135,161,175	0
4	NAG	D	1	14/15	0.91	0.13	67,92,107,110	0
3	NAG	F	1	14/15	0.93	0.26	50,77,90,92	0
3	NAG	G	1	14/15	0.94	0.11	43,71,81,97	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.













6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	NAG	А	1005	14/15	0.74	0.35	106,139,160,168	0
6	NAG	А	1017	14/15	0.76	0.38	90,122,144,150	0
6	NAG	А	1006	14/15	0.77	0.30	111,131,154,171	0
5	ZN	А	1001	1/1	0.83	0.10	76,76,76,76	0
6	NAG	А	1004	14/15	0.92	0.14	63,88,99,100	0

6.5 Other polymers (i)

There are no such residues in this entry.

