



# Full wwPDB X-ray Structure Validation Report i

Sep 23, 2023 – 03:53 PM EDT

PDB ID : 5KYV  
Title : Structure of Photinus pyralis Luciferase green shifted light emitting variant  
Authors : Gulick, A.M.  
Deposited on : 2016-07-22  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	TLA	A	602	-	X	-	-

## 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 8314 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Luciferin 4-monooxygenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	530	4017	2595	654	750	18	0	2	0
1	B	543	4136	2666	685	768	17	0	0	0

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	241	ILE	VAL	engineered mutation	UNP P08659
A	246	ALA	GLY	engineered mutation	UNP P08659
A	250	SER	PHE	engineered mutation	UNP P08659
B	241	ILE	VAL	engineered mutation	UNP P08659
B	246	ALA	GLY	engineered mutation	UNP P08659
B	250	SER	PHE	engineered mutation	UNP P08659

- Molecule 2 is 5'-O-[N-(DEHYDROLUCIFERYL)-SULFAMOYL] ADENOSINE (three-letter code: SLU) (formula: C<sub>21</sub>H<sub>18</sub>N<sub>8</sub>O<sub>8</sub>S<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
2	A	1	40	21	8	8	3	0	0
2	B	1	40	21	8	8	3	0	0

- Molecule 3 is L(+)-TARTARIC ACID (three-letter code: TLA) (formula: C<sub>4</sub>H<sub>6</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	A	1	10	4	6	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	39	39	39	0	0
4	B	32	32	32	0	0


### 3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Luciferin 4-monooxygenase



- Molecule 1: Luciferin 4-monooxygenase



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	93.08Å 93.08Å 293.64Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.06 – 2.50 29.83 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.1 (29.06-2.50) 99.1 (29.83-2.50)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	1.79 (at 2.51Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
$R$ , $R_{free}$	0.194 , 0.247 0.204 , 0.250	Depositor DCC
$R_{free}$ test set	2634 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	51.5	Xtriage
Anisotropy	0.260	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 46.4	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.46$ , $< L^2 > = 0.29$	Xtriage
Estimated twinning fraction	0.039 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	8314	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.92% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [\(i\)](#)

### 5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TLA, SLU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.43	0/4112	0.58	0/5598
1	B	0.41	0/4229	0.56	0/5748
All	All	0.42	0/8341	0.57	0/11346

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4017	0	3904	51	0
1	B	4136	0	4066	62	0
2	A	40	0	17	2	0
2	B	40	0	17	0	0
3	A	10	0	4	0	0
4	A	39	0	0	0	0
4	B	32	0	0	2	0
All	All	8314	0	8008	111	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (111) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:353:PRO:HG2	1:A:356:ASP:HB3	1.62	0.78
1:A:455:GLU:HG3	1:A:469:VAL:HG23	1.65	0.77
1:A:288:VAL:HG23	1:A:290:THR:HG22	1.70	0.72
1:A:187:ASP:OD1	1:A:190:LYS:HE2	1.92	0.69
1:A:298:SER:HB2	1:A:330:ARG:HH11	1.54	0.69
1:B:423:ILE:HD11	1:B:438:LEU:HD13	1.74	0.69
1:B:390:LEU:O	1:B:420:SER:HB3	1.96	0.66
1:A:476:ASP:OD1	1:A:476:ASP:N	2.31	0.64
1:A:298:SER:HB2	1:A:330:ARG:NH1	2.14	0.61
1:A:346:THR:HB	1:A:396:MET:HG2	1.84	0.60
1:A:495:GLU:HG3	1:A:516:VAL:HB	1.85	0.59
1:A:322:GLU:H	1:A:322:GLU:CD	2.07	0.58
1:A:467:ALA:HB2	1:A:485:VAL:HG13	1.86	0.58
1:A:338:GLN:HE21	2:A:601:SLU:HG381	1.51	0.57
1:B:77:ARG:HG2	1:B:101:ALA:HB3	1.86	0.57
1:B:325:GLU:HA	1:B:328:ALA:HB3	1.84	0.57
1:B:375:ASP:HB2	1:B:382:LEU:HD21	1.87	0.56
1:A:512:LEU:O	1:A:514:GLY:N	2.37	0.56
1:B:32:ARG:NH2	4:B:703:HOH:O	2.39	0.56
1:B:240:VAL:HG11	1:B:291:LEU:HD11	1.89	0.55
1:A:79:VAL:HG13	1:A:123:PRO:HB3	1.89	0.53
1:A:230:GLN:NE2	1:B:231:ILE:HD12	2.24	0.52
1:A:296:ALA:C	1:A:330:ARG:HH12	2.13	0.52
1:B:265:MET:HE2	1:B:268:PHE:HA	1.93	0.51
1:B:524:LYS:HG2	1:B:530:LEU:HD23	1.93	0.51
1:B:277:LEU:HA	1:B:282:ILE:HG13	1.91	0.51
1:B:444:TYR:HB2	1:B:479:GLU:O	2.11	0.51
1:B:151:ILE:HD13	1:B:158:TYR:HB3	1.93	0.50
1:B:151:ILE:HD11	1:B:161:PHE:HB2	1.93	0.50
1:B:288:VAL:HG22	1:B:289:PRO:HD2	1.93	0.50
1:A:384:VAL:HB	1:A:426:TRP:CZ3	2.47	0.50
1:B:321:LYS:O	1:B:325:GLU:HG2	2.12	0.49
1:B:455:GLU:HG3	1:B:469:VAL:HG23	1.94	0.49
1:B:440:SER:HB3	1:B:451:PRO:HG2	1.94	0.49
1:A:79:VAL:HG22	1:A:126:VAL:HG13	1.95	0.48
1:A:474:ASP:OD2	1:A:475:ASP:N	2.47	0.48
1:A:349:ILE:HD12	1:A:392:VAL:HG11	1.95	0.48
1:B:349:ILE:HD12	1:B:392:VAL:HG11	1.94	0.48
1:B:193:ALA:HB2	1:B:212:HIS:CD2	2.48	0.48
1:B:423:ILE:HG13	1:B:441:LEU:HD11	1.96	0.48

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:225:PRO:HB3	1:A:230:GLN:HB2	1.96	0.47
1:A:273:PHE:CZ	1:A:287:LEU:HD21	2.49	0.47
1:B:482:ALA:HB2	1:B:514:GLY:HA3	1.96	0.47
1:B:77:ARG:HH21	1:B:122:GLN:CB	2.28	0.47
1:A:88:PHE:O	1:A:91:PRO:HD2	2.14	0.47
1:B:118:MET:HG2	1:B:123:PRO:HG3	1.96	0.47
1:B:125:VAL:HG22	1:B:148:LYS:HB3	1.97	0.47
1:A:111:GLU:OE2	1:A:138:ASN:ND2	2.38	0.47
1:B:151:ILE:CD1	1:B:161:PHE:HB2	2.44	0.47
1:A:373:VAL:HG12	1:A:382:LEU:HD12	1.97	0.46
1:B:121:SER:HB2	1:B:123:PRO:HD3	1.98	0.45
1:B:78:ILE:HG23	1:B:100:VAL:HG11	1.97	0.45
1:B:79:VAL:HG23	1:B:123:PRO:HB3	1.98	0.45
1:B:549:LYS:O	1:B:550:LEU:HB2	2.17	0.45
1:B:153:ASP:OD1	1:B:153:ASP:N	2.45	0.45
1:A:298:SER:O	1:A:300:LEU:N	2.45	0.45
1:B:394:GLY:O	1:B:397:ILE:HG13	2.17	0.44
1:B:344:GLU:OE1	1:B:344:GLU:N	2.42	0.44
1:A:522:VAL:O	1:A:524:LYS:N	2.50	0.44
1:A:258:CYS:HB3	1:A:260:PHE:CE1	2.53	0.44
1:A:79:VAL:HG13	1:A:123:PRO:CB	2.48	0.44
1:A:338:GLN:HG2	1:A:362:VAL:HA	2.00	0.44
1:B:76:HIS:HA	4:B:714:HOH:O	2.17	0.43
1:B:384:VAL:O	1:B:386:GLN:HG3	2.18	0.43
1:A:230:GLN:HE21	1:B:231:ILE:HD12	1.83	0.43
1:B:172:LEU:HA	1:B:173:PRO:HD2	1.88	0.43
1:B:364:LYS:HG2	1:B:430:GLU:O	2.18	0.43
1:B:524:LYS:CG	1:B:530:LEU:HD23	2.49	0.43
1:A:254:GLY:O	1:A:257:ILE:HG22	2.18	0.43
1:B:459:LEU:HD21	1:B:466:ASP:HA	2.00	0.43
1:B:98:ILE:HG13	1:B:100:VAL:HG23	1.99	0.43
1:B:506:VAL:HG23	1:B:507:THR:O	2.18	0.43
1:A:289:PRO:HG2	1:A:317:ALA:HB3	2.00	0.43
1:B:378:THR:HB	1:B:380:LYS:H	1.83	0.43
1:B:516:VAL:O	1:B:543:LYS:NZ	2.51	0.43
1:A:114:LEU:HD21	1:A:136:ILE:HD13	2.00	0.43
1:B:77:ARG:HH21	1:B:122:GLN:HB2	1.84	0.43
1:A:339:GLY:HA3	2:A:601:SLU:C5	2.49	0.43
1:B:278:GLN:NE2	1:B:304:TYR:HD2	2.17	0.42
1:A:480:LEU:HB3	1:A:513:ARG:O	2.18	0.42
1:B:271:GLU:O	1:B:275:ARG:HB2	2.19	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:444:TYR:HA	1:B:445:LYS:HA	1.86	0.42
1:A:295:PHE:O	1:A:330:ARG:NH1	2.52	0.42
1:B:83:GLU:OE1	1:B:131:LYS:HE3	2.19	0.42
1:B:273:PHE:HE1	1:B:277:LEU:HD11	1.84	0.42
1:B:445:LYS:HA	1:B:477:ALA:O	2.20	0.42
1:B:289:PRO:HG2	1:B:317:ALA:HB3	2.01	0.42
1:B:524:LYS:HG2	1:B:530:LEU:HA	2.01	0.42
1:A:89:PHE:CE1	1:A:252:THR:HG21	2.55	0.42
1:A:199:SER:HA	1:A:205:PRO:HA	2.02	0.42
1:A:349:ILE:CD1	1:A:392:VAL:HG21	2.49	0.42
1:A:340:TYR:HB3	1:A:350:LEU:HB2	2.02	0.41
1:A:301:ILE:HG23	1:A:302:ASP:OD1	2.20	0.41
1:B:349:ILE:H	1:B:349:ILE:HG12	1.73	0.41
1:B:520:ASP:OD1	1:B:520:ASP:N	2.53	0.41
1:A:65:GLU:HG2	1:A:176:PHE:CE1	2.54	0.41
1:A:182:VAL:HA	1:A:183:PRO:HD2	1.95	0.41
1:B:102:VAL:O	1:B:104:PRO:HD3	2.20	0.41
1:A:177:ASN:ND2	1:A:180:ASP:OD2	2.53	0.41
1:B:523:PRO:HB2	1:B:531:ASP:HB3	2.02	0.41
1:A:369:PHE:HE1	1:A:396:MET:HB2	1.84	0.41
1:A:522:VAL:HA	1:A:523:PRO:HD3	1.94	0.41
1:B:349:ILE:HD12	1:B:392:VAL:HG21	2.03	0.41
1:A:508:THR:HG23	1:A:509:ALA:H	1.86	0.41
1:A:79:VAL:CG2	1:A:126:VAL:HG13	2.51	0.40
1:B:18:GLU:O	1:B:188:ARG:NH2	2.53	0.40
1:A:482:ALA:HB3	1:A:539:LEU:HD13	2.03	0.40
1:B:208:VAL:HG13	1:B:398:MET:SD	2.62	0.40
1:B:241:ILE:HA	1:B:242:PRO:HD3	1.92	0.40
1:B:474:ASP:OD1	1:B:476:ASP:N	2.46	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	524/550 (95%)	490 (94%)	25 (5%)	9 (2%)	9 16
1	B	539/550 (98%)	510 (95%)	27 (5%)	2 (0%)	34 54
All	All	1063/1100 (97%)	1000 (94%)	52 (5%)	11 (1%)	15 28

All (11) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	299	THR
1	B	506	VAL
1	A	513	ARG
1	A	526	LEU
1	A	298	SER
1	A	475	ASP
1	A	516	VAL
1	A	520	ASP
1	A	523	PRO
1	B	298	SER
1	A	508	THR

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	416/463 (90%)	410 (99%)	6 (1%)	67 86
1	B	432/463 (93%)	428 (99%)	4 (1%)	78 92
All	All	848/926 (92%)	838 (99%)	10 (1%)	73 88

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	56	TYR
1	A	300	LEU
1	A	310[A]	HIS
1	A	310[B]	HIS
1	A	320	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	322	GLU
1	B	320	SER
1	B	340	TYR
1	B	420	SER
1	B	506	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [\(i\)](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SLU	B	601	-	35,45,45	1.86	7 (20%)	41,68,68	2.50	9 (21%)
3	TLA	A	602	-	9,9,9	1.40	1 (11%)	12,12,12	1.81	5 (41%)
2	SLU	A	601	-	35,45,45	1.59	6 (17%)	41,68,68	2.47	10 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SLU	B	601	-	-	3/10/39/39	0/6/6/6
3	TLA	A	602	-	-	10/12/12/12	-
2	SLU	A	601	-	-	1/10/39/39	0/6/6/6

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	SLU	C4-S9	6.17	1.78	1.74
2	B	601	SLU	O26-C25	4.04	1.46	1.41
2	A	601	SLU	C16-N40	-3.87	1.34	1.39
2	B	601	SLU	C16-N40	-3.86	1.34	1.39
2	A	601	SLU	O20-S17	-3.83	1.51	1.59
2	A	601	SLU	O26-C25	3.41	1.45	1.41
2	B	601	SLU	O20-S17	-3.28	1.52	1.59
2	A	601	SLU	C4-S9	3.18	1.76	1.74
2	B	601	SLU	S17-N40	2.79	1.64	1.59
2	B	601	SLU	C13-S12	2.69	1.74	1.70
2	A	601	SLU	S17-N40	2.62	1.64	1.59
3	A	602	TLA	C2-C1	-2.60	1.49	1.52
2	B	601	SLU	C36-N37	-2.29	1.30	1.34
2	A	601	SLU	C13-S12	2.19	1.73	1.70

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	SLU	C14-C13-S12	-8.33	101.56	111.79
2	B	601	SLU	C14-C13-S12	-8.30	101.60	111.79
2	B	601	SLU	C3-C4-S9	-6.94	102.64	111.85
2	A	601	SLU	C3-C4-S9	-6.66	103.02	111.85
2	B	601	SLU	C5-C4-S9	5.42	135.95	125.10
2	B	601	SLU	C13-S12-C11	4.53	96.59	91.12
2	A	601	SLU	C13-S12-C11	4.46	96.50	91.12
2	B	601	SLU	C21-O20-S17	-4.43	108.22	117.37
2	A	601	SLU	C5-C4-S9	4.41	133.92	125.10
2	A	601	SLU	N31-C30-N29	-3.98	122.46	128.68
2	B	601	SLU	N31-C30-N29	-3.89	122.60	128.68
2	A	601	SLU	C21-O20-S17	-3.84	109.45	117.37
2	A	601	SLU	O19-S17-O18	3.50	126.21	120.76
2	B	601	SLU	O19-S17-O18	3.41	126.06	120.76

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	SLU	C4-C3-N7	3.24	115.58	108.04
2	B	601	SLU	C4-C3-N7	3.19	115.46	108.04
3	A	602	TLA	O41-C4-C3	3.03	121.47	113.27
3	A	602	TLA	O2-C2-C1	-2.76	104.89	110.66
2	A	601	SLU	C32-C33-N37	-2.72	106.56	109.40
3	A	602	TLA	O2-C2-C3	2.60	115.40	110.23
2	B	601	SLU	C16-N40-S17	-2.58	120.13	123.38
2	A	601	SLU	O39-C16-N40	2.28	123.88	121.08
3	A	602	TLA	C2-C3-C4	2.17	114.71	109.87
3	A	602	TLA	O4-C4-C3	-2.07	116.19	121.63

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	SLU	C16-N40-S17-O19
2	B	601	SLU	C16-N40-S17-O19
3	A	602	TLA	C1-C2-C3-C4
3	A	602	TLA	O2-C2-C3-O3
3	A	602	TLA	C2-C3-C4-O4
3	A	602	TLA	C2-C3-C4-O41
3	A	602	TLA	C1-C2-C3-O3
3	A	602	TLA	O2-C2-C3-C4
3	A	602	TLA	O1-C1-C2-O2
3	A	602	TLA	O11-C1-C2-O2
3	A	602	TLA	O3-C3-C4-O4
3	A	602	TLA	O3-C3-C4-O41
2	B	601	SLU	O20-C21-C22-O26
2	B	601	SLU	O20-C21-C22-C23


There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	SLU	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be

highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [\(i\)](#)

### 6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	530/550 (96%)	0.04	33 (6%) 20 21	33, 47, 91, 103	0
1	B	543/550 (98%)	0.05	32 (5%) 22 23	38, 56, 79, 90	0
All	All	1073/1100 (97%)	0.04	65 (6%) 21 22	33, 51, 86, 103	0

All (65) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	92	VAL	4.8
1	A	175	GLY	4.7
1	A	493	MET	4.6
1	A	174	PRO	4.5
1	A	518	PHE	4.2
1	B	175	GLY	4.2
1	A	527	THR	4.0
1	B	527	THR	3.9
1	A	447	TYR	3.4
1	A	92	VAL	3.2
1	B	95	ALA	3.2
1	A	301	ILE	3.1
1	A	492	THR	3.1
1	A	538	ILE	3.0
1	B	89	PHE	3.0
1	A	476	ASP	3.0
1	B	476	ASP	2.9
1	A	477	ALA	2.9
1	B	447	TYR	2.9
1	A	93	LEU	2.9
1	B	266	TYR	2.9
1	A	519	VAL	2.9
1	A	540	ILE	2.8
1	B	253	LEU	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	96	LEU	2.8
1	A	504	SER	2.8
1	B	334	PRO	2.8
1	A	204	LEU	2.7
1	A	516	VAL	2.7
1	A	199	SER	2.7
1	B	444	TYR	2.7
1	B	405	PRO	2.7
1	A	94	GLY	2.6
1	A	521	GLU	2.6
1	B	249	MET	2.6
1	B	251	THR	2.6
1	A	302	ASP	2.6
1	B	91	PRO	2.5
1	B	93	LEU	2.5
1	B	102	VAL	2.5
1	A	494	THR	2.5
1	A	539	LEU	2.4
1	A	173	PRO	2.4
1	B	489	HIS	2.3
1	B	488	GLU	2.3
1	A	486	VAL	2.3
1	A	499	VAL	2.3
1	B	250	SER	2.3
1	B	526	LEU	2.3
1	B	100	VAL	2.2
1	A	95	ALA	2.2
1	A	91	PRO	2.2
1	A	542	ALA	2.2
1	B	294	PHE	2.2
1	B	174	PRO	2.2
1	A	478	GLY	2.1
1	B	94	GLY	2.1
1	A	487	LEU	2.1
1	A	500	ASP	2.1
1	B	204	LEU	2.1
1	B	332	HIS	2.1
1	B	406	GLU	2.1
1	B	111	GLU	2.1
1	B	109	TYR	2.0
1	B	490	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)


There are no monosaccharides in this entry.


## 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	TLA	A	602	10/10	0.75	0.23	51,63,64,65	0
2	SLU	B	601	40/40	0.98	0.12	34,44,50,55	0
2	SLU	A	601	40/40	0.98	0.11	30,37,42,50	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.