



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 6, 2020 – 11:38 AM BST

PDB ID : 5KZQ
Title : Metabotropic Glutamate Receptor in complex with antagonist (1 {S},2 {R},3 {S},4 {S},5 {R},6 {R})-2-azanyl-3-[[3,4-bis(fluoranyl)phenyl]sulfanylmethyl]-4-oxidanyl-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid
Authors : Chappell, M.D.; Li, R.; Smith, S.C.; Dressman, B.A.; Tromiczak, E.G.; Tripp, A.E.; Blanco, M.-J.; Vetman, T.; Quimby, S.J.; Matt, J.; Britton, T.; Fivush, A.M.; Schkeryantz, J.M.; Mayhugh, D.; Erickson, J.A.; Bures, M.; Jaramillo, C.; Carpintero, M.; de Diego, J.E.; Barberis, M.; Garcia-Cerrada, S.; Soriano, J.F.; Antonysamy, S.; Atwell, S.; MacEwan, I.; Condon, B.; Bradley, C.; Wang, J.; Zhang, A.; Connors, K.; Groshong, C.; Wasserman, S.R.; Koss, J.W.; Witkin, J.M.; Li, X.; Overshiner, C.; Wafford, K.A.; Seidel, W.; Wang, X.-S.; Heinz, B.A.; Swanson, S.; Catlow, J.; Bedwell, D.; Monn, J.A.; Mitch, C.H.; Ornstein, P.
Deposited on : 2016-07-25
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	570	

Refmac : 5.8.0158
 CCP4 : 7.0.044 (Gargrove)
 Ideal geometry (proteins) : Engh & Huber (2001)
 Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
 Validation Pipeline (wwPDB-VP) : 2.13.1

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 3747 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Metabotropic glutamate receptor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	503	3702	2360	621	699	22	0	0	0

There are 7 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	563	GLU	TRP	conflict	UNP Q14416
A	565	HIS	-	expression tag	UNP Q14416
A	566	HIS	-	expression tag	UNP Q14416
A	567	HIS	-	expression tag	UNP Q14416
A	568	HIS	-	expression tag	UNP Q14416
A	569	HIS	-	expression tag	UNP Q14416
A	570	HIS	-	expression tag	UNP Q14416

- Molecule 2 is (1 {S},2 {R},3 {S},4 {S},5 {R},6 {R})-2-azanyl-3-[[3,4-bis(fluoranyl)phenyl]sulfanylmethyl]-4-oxidanyl-bicyclo[3.1.0]hexane-2,6-dicarboxylic acid (three-letter code: 6YS) (formula: C₁₅H₁₅F₂NO₅S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	F	N	O			S
2	A	1	24	15	2	1	5	1	0	0

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	14	8	1	5	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	7	7	7	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	166.30Å 166.30Å 79.78Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	32.80 – 2.80 47.33 – 2.80	Depositor EDS
% Data completeness (in resolution range)	91.1 (32.80-2.80) 91.1 (47.33-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.13	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.80 (at 2.81Å)	Xtrriage
Refinement program	BUSTER-TNT 2.11.6	Depositor
R, R_{free}	0.208 , 0.245 0.230 , 0.264	Depositor DCC
R_{free} test set	1278 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	85.3	Xtrriage
Anisotropy	0.090	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 100.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3747	wwPDB-VP
Average B, all atoms (Å ²)	121.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 6YS, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.52	0/3796	0.74	0/5187

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3702	0	3363	54	0
2	A	24	0	0	0	0
3	A	14	0	13	0	0
4	A	7	0	0	0	0
All	All	3747	0	3376	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:324:ILE:HG13	1:A:432:PHE:HB2	1.58	0.86
1:A:186:PRO:HG3	1:A:451:GLY:HA2	1.68	0.75
1:A:142:GLY:HA3	1:A:148:SER:OG	1.99	0.62
1:A:183:ARG:HG3	1:A:185:VAL:O	1.99	0.62
1:A:234:CYS:HB2	1:A:518:CYS:SG	2.41	0.61
1:A:151:VAL:HG12	1:A:155:LEU:HD22	1.84	0.60
1:A:204:TRP:CD1	1:A:497:ALA:HA	2.39	0.58
1:A:322:TYR:HE1	1:A:324:ILE:HD12	1.69	0.57
1:A:110:LEU:HD13	1:A:158:PHE:HE2	1.69	0.57
1:A:181:PHE:HE2	1:A:183:ARG:HD2	1.72	0.55
1:A:324:ILE:HD13	1:A:379:MET:HA	1.89	0.54
1:A:207:VAL:HG12	1:A:265:VAL:HB	1.89	0.54
1:A:377:LYS:HA	1:A:380:PHE:HD2	1.74	0.53
1:A:65:MET:HE3	1:A:140:ILE:HB	1.91	0.53
1:A:279:LEU:HD21	1:A:311:ALA:HB2	1.90	0.52
1:A:293:ALA:HB3	1:A:316:THR:HG22	1.92	0.52
1:A:322:TYR:CE1	1:A:324:ILE:HD12	2.45	0.52
1:A:141:GLY:HA2	1:A:165:TYR:CE2	2.45	0.51
1:A:399:LEU:HD12	1:A:410:MET:HG3	1.94	0.50
1:A:46:PRO:HB2	1:A:48:GLU:HG3	1.93	0.50
1:A:386:TYR:CG	1:A:431:PRO:HG3	2.46	0.49
1:A:419:TYR:CE1	1:A:424:LEU:HD11	2.48	0.49
1:A:65:MET:CE	1:A:140:ILE:HB	2.43	0.49
1:A:139:VAL:HG21	1:A:155:LEU:HD21	1.94	0.48
1:A:36:GLY:HA2	1:A:140:ILE:O	2.12	0.48
1:A:34:LEU:O	1:A:86:ALA:HA	2.13	0.48
1:A:191:GLN:HB2	1:A:317:ILE:HD13	1.96	0.48
1:A:199:LEU:HD13	1:A:207:VAL:HG11	1.96	0.47
1:A:210:VAL:HB	1:A:268:LEU:HD22	1.95	0.47
1:A:199:LEU:HB2	1:A:233:ILE:HD13	1.96	0.47
1:A:375:GLU:HB3	1:A:378:ILE:HG12	1.97	0.47
1:A:212:SER:O	1:A:217:GLY:HA3	2.16	0.46
1:A:251:GLU:HG3	1:A:281:ALA:HB1	1.98	0.44
1:A:270:THR:O	1:A:296:GLY:HA3	2.18	0.44
1:A:161:PRO:HG2	1:A:418:LEU:HD23	1.99	0.44
1:A:26:LEU:HB3	1:A:88:ILE:HB	1.99	0.44
1:A:186:PRO:HD2	1:A:319:LEU:HD11	1.99	0.43
1:A:221:ILE:HD13	1:A:221:ILE:HA	1.85	0.43
1:A:199:LEU:HD22	1:A:204:TRP:HE3	1.83	0.43
1:A:420:LYS:HA	1:A:424:LEU:HD12	2.00	0.43
1:A:204:TRP:CD2	1:A:265:VAL:HG21	2.54	0.43
1:A:188:ASP:O	1:A:191:GLN:HG2	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:508:GLU:HA	1:A:524:PRO:HA	2.01	0.42
1:A:210:VAL:HB	1:A:268:LEU:CD2	2.50	0.42
1:A:89:LEU:HD12	1:A:105:PHE:CZ	2.54	0.42
1:A:106:VAL:HG11	1:A:154:LEU:HG	2.01	0.42
1:A:141:GLY:HA2	1:A:165:TYR:HE2	1.84	0.42
1:A:297:TRP:HA	1:A:303:VAL:HG21	2.01	0.42
1:A:110:LEU:HD13	1:A:158:PHE:CE2	2.52	0.41
1:A:181:PHE:CE2	1:A:183:ARG:HD2	2.54	0.41
1:A:185:VAL:HG23	1:A:319:LEU:HD21	2.02	0.41
1:A:100:GLU:O	1:A:103:LEU:HB2	2.21	0.41
1:A:443:PHE:HB3	1:A:447:GLY:HA2	2.03	0.41
1:A:159:GLN:O	1:A:415:GLY:HA3	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	495/570 (87%)	453 (92%)	39 (8%)	3 (1%)	25 56

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	243	ARG
1	A	497	ALA
1	A	437	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	356/463 (77%)	324 (91%)	32 (9%)	9 28

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	LEU
1	A	48	GLU
1	A	95	ASP
1	A	150	GLN
1	A	155	LEU
1	A	159	GLN
1	A	175	LYS
1	A	183	ARG
1	A	221	ILE
1	A	237	THR
1	A	241	VAL
1	A	262	SER
1	A	268	LEU
1	A	269	PHE
1	A	302	SER
1	A	308	GLU
1	A	321	SER
1	A	324	ILE
1	A	346	ARG
1	A	348	PHE
1	A	362	CYS
1	A	410	MET
1	A	416	ARG
1	A	417	ARG
1	A	426	VAL
1	A	429	ASP
1	A	446	PHE
1	A	452	ARG
1	A	457	THR
1	A	459	LEU
1	A	500	CYS
1	A	513	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	513	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	A	602	1	14,14,15	0.36	0	17,19,21	0.65	1 (5%)
2	6YS	A	601	-	16,26,26	0.76	0	21,41,41	1.23	3 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	602	1	-	0/6/23/26	0/1/1/1
2	6YS	A	601	-	-	3/5/44/44	0/3/3/3

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	A	601	6YS	C6-C3-C5	2.86	120.59	118.18
2	A	601	6YS	C3-C5-C4	-2.30	118.41	121.03
3	A	602	NAG	C1-O5-C5	2.12	115.07	112.19
2	A	601	6YS	C11-C10-C13	2.11	110.83	106.31

There are no chirality outliers.


All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	6YS	C13-C12-C15-S24
2	A	601	6YS	C2-C6-S24-C15
2	A	601	6YS	C3-C6-S24-C15

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	503/570 (88%)	0.47	43 (8%) 10 5	62, 114, 197, 222	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	525	CYS	5.5
1	A	530	TYR	5.0
1	A	507	ASN	4.9
1	A	538	ALA	4.5
1	A	532	LEU	4.4
1	A	536	THR	4.2
1	A	506	GLN	4.1
1	A	531	ARG	4.0
1	A	353	PHE	3.5
1	A	514	PRO	3.4
1	A	527	PRO	3.2
1	A	524	PRO	3.2
1	A	523	ILE	3.2
1	A	349	TRP	3.1
1	A	515	GLY	3.0
1	A	539	ASP	2.8
1	A	509	VAL	2.8
1	A	535	PHE	2.8
1	A	537	CYS	2.8
1	A	196	ALA	2.8
1	A	542	LEU	2.7
1	A	503	PRO	2.7
1	A	511	SER	2.6
1	A	489	SER	2.6
1	A	504	CYS	2.6
1	A	281	ALA	2.6
1	A	233	ILE	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	204	TRP	2.5
1	A	289	PHE	2.5
1	A	285	LEU	2.5
1	A	231	ARG	2.4
1	A	533	ASP	2.4
1	A	218	GLU	2.4
1	A	500	CYS	2.4
1	A	541	GLY	2.3
1	A	357	PHE	2.3
1	A	350	GLU	2.2
1	A	512	VAL	2.2
1	A	545	TRP	2.2
1	A	315	ILE	2.1
1	A	496	PRO	2.0
1	A	359	GLN	2.0
1	A	435	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	A	602	14/15	0.90	0.19	150,157,160,160	0
2	6YS	A	601	24/24	0.95	0.26	80,87,102,108	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.