

# Full wwPDB X-ray Structure Validation Report (i)

#### Oct 31, 2023 – 03:53 PM EDT

PDB ID	:	3L4Q
Title	:	Structural insights into phosphoinositide 3-kinase activation by the influenza
		A virus NS1 protein
Authors	:	Hale, B.G.; Kerry, P.S.; Jackson, D.; Precious, B.L.; Gray, A.; Killip, M.J.;
		Randall, R.E.; Russell, R.J.
Deposited on	:	2009-12-21
Resolution	:	2.30  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
$\mathrm{EDS}$	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575(2.30-2.30)
Sidechain outliers	138945	5575(2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain				
1	А	164	<sup>2%</sup> 60%	12% •	27%		
1	В	164	53%	18% •	27%		
2	С	170	17%		16% • •		
2	D	170	66%		30% •		



#### $\mathbf{2}$ Entry composition (i)

67

68

69

70

71

72

HIS

HIS

HIS

HIS

HIS

HIS

В

В

В

В

В

В

There are 4 unique types of molecules in this entry. The entry contains 4923 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

		1				1			
Mol	Chain	Residues		At	oms			ZeroOcc	AltConf
1	٨	190	Total	С	Ν	0	S	0	0
T	A	120	933	593	158	176	6		0

С

Ν

Ο

expression tag

expression tag

expression tag

expression tag

expression tag

expression tag

S

UNP P03496

UNP P03496

UNP P03496

UNP P03496

UNP P03496

UNP P03496

• Molecule 1 is a protein called Non-structural protein 1.

Total

1	В	119	927	590	157	174	5 6	0	0	
There are 12 discrepancies between the modelled and reference sequences:										
Chair	ı Resid	due Mode	elled A	ctual	C	omme	ent	Referen	ce	
А	67	HI	S	-	exp	ressio	n tag	UNP P03	496	
А	68	HI	S	-	exp	ressio	n tag	UNP P03	496	
А	69	HI	S	-	exp	ressio	n tag	UNP P03	496	
А	70	HI	S	-	exp	ressio	n tag	UNP P03	496	
А	71	HI	S	-	exp	ressio	n tag	UNP P03	496	
A	72	HI	S	-	exp	ressio	n tag	UNP P03	496	

\_

\_

\_

\_

-

\_

• Molecule 2 is a protein called Phosphatidylinositol 3-kinase regulatory subunit beta.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
2	С	163	Total	С	Ν	0	$\mathbf{S}$	0	0	0
			1385	855	256	270	4	0		0
9	Л	163	Total	С	Ν	0	S	0	0	0
2 D	105	1385	855	256	270	4	0	0	U	

There are 2 discrepancies between the modelled and reference sequences:

Trace

0

0



$3L_2$	4Q
--------	----

Chain	Residue	Modelled	Actual	Comment	Reference
С	495	SER	CYS	engineered mutation	UNP P23726
D	495	SER	CYS	engineered mutation	UNP P23726



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	С	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0
3	D	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	72	Total O 72 72	0	0
4	С	76	Total         O           76         76	0	0
4	В	68	Total         O           68         68	0	0
4	D	65	$\begin{array}{cc} \text{Total} & \text{O} \\ 65 & 65 \end{array}$	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Non-structural protein 1





# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	57.95Å 98.67Å 149.94Å	Deperitor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
$\mathbf{B}_{\mathrm{ascolution}}(\hat{\boldsymbol{\lambda}})$	49.97 - 2.30	Depositor
Resolution (A)	49.98 - 2.30	EDS
% Data completeness	99.3 (49.97-2.30)	Depositor
(in resolution range)	95.2 (49.98-2.30)	EDS
$R_{merge}$	(Not available)	Depositor
R <sub>sym</sub>	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.16 (at 2.29 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.5.0070	Depositor
D D.	0.232 , $0.292$	Depositor
$n, n_{free}$	0.236 , (Not available)	DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor $(Å^2)$	35.5	Xtriage
Anisotropy	0.372	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.34, $58.6$	EDS
L-test for twinning <sup>2</sup>	$ < L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	4923	wwPDB-VP
Average B, all atoms $(Å^2)$	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 14.42% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

# 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Chain	Bond lengths		Bond angles	
1VIOI		RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	А	1.08	0/948	0.94	1/1285~(0.1%)
1	В	1.06	0/942	0.94	1/1277~(0.1%)
2	С	0.96	0/1401	0.85	0/1868
2	D	0.85	0/1401	0.82	3/1868~(0.2%)
All	All	0.98	0/4692	0.88	5/6298~(0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	А	0	2
2	С	0	1
All	All	0	3

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms		$Observed(^{o})$	$Ideal(^{o})$
2	D	561	LEU	CA-CB-CG	7.64	132.88	115.30
1	А	90	LEU	CA-CB-CG	-5.96	101.58	115.30
1	В	98	MET	CG-SD-CE	-5.49	91.42	100.20
2	D	560	ASP	CB-CG-OD1	5.28	123.05	118.30
2	D	500	LEU	CA-CB-CG	5.02	126.84	115.30

There are no chirality outliers.

All (3) planarity outliers are listed below:



Mol	Chain	Res	Type	Group
1	А	165	SER	Peptide
1	А	86	ALA	Peptide
2	С	568	ARG	Sidechain

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	933	0	955	29	0
1	В	927	0	950	46	0
2	С	1385	0	1385	31	0
2	D	1385	0	1385	53	0
3	С	6	0	8	0	0
3	D	6	0	8	3	0
4	А	72	0	0	5	0
4	В	68	0	0	8	0
4	С	76	0	0	3	0
4	D	65	0	0	15	0
All	All	4923	0	4691	149	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (149) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:165:SER:HB3	1:A:166:LEU:CD2	1.68	1.22
2:C:506:GLU:CB	2:C:507:GLY:HA2	1.65	1.20
1:A:165:SER:CB	1:A:166:LEU:HD22	1.74	1.18
1:A:165:SER:HB3	1:A:166:LEU:HD22	1.20	1.15
1:B:84:VAL:HB	1:B:85:PRO:HD3	1.14	1.11
2:C:506:GLU:HB3	2:C:507:GLY:HA2	1.31	1.08
2:D:449:ASP:OD1	2:D:452:ARG:NH1	1.97	0.97
2:C:506:GLU:HB2	2:C:507:GLY:HA2	1.44	0.97
1:B:134:PHE:CZ	1:B:197:THR:CG2	2.52	0.92
1:B:84:VAL:CB	1:B:85:PRO:HD3	2.00	0.90
1:B:134:PHE:CZ	1:B:197:THR:HG22	2.08	0.88



		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
2:C:506:GLU:HB3	2:C:507:GLY:CA	2.03	0.88
2:C:506:GLU:CB	2:C:507:GLY:CA	2.51	0.88
1:B:84:VAL:HB	1:B:85:PRO:CD	2.00	0.88
2:C:585:ILE:O	2:C:589:LEU:HD13	1.78	0.83
1:B:134:PHE:CD1	1:B:141:LEU:HD13	2.14	0.83
2:D:590:GLY:O	2:D:591:ILE:HG22	1.79	0.81
1:B:134:PHE:HZ	1:B:197:THR:CG2	1.94	0.78
1:B:163:LEU:HD12	1:B:166:LEU:CD1	2.13	0.78
2:C:489:GLY:HA2	2:C:522:LEU:HD21	1.65	0.78
1:A:165:SER:OG	1:A:166:LEU:HD22	1.85	0.77
1:A:165:SER:HB3	1:A:166:LEU:HD23	1.63	0.77
1:B:134:PHE:CE2	1:B:197:THR:HG22	2.20	0.76
2:C:571:TYR:CE2	1:B:167:PRO:HB2	2.20	0.76
1:B:114:PRO:HA	1:B:166:LEU:HD13	1.68	0.75
2:C:489:GLY:CA	2:C:522:LEU:HD21	2.16	0.75
2:D:469:GLN:OE1	4:D:233:HOH:O	2.05	0.73
1:B:134:PHE:CD1	1:B:141:LEU:CD1	2.72	0.73
2:D:590:GLY:O	2:D:591:ILE:CG2	2.39	0.70
2:C:505:ARG:O	4:C:45:HOH:O	2.10	0.69
1:A:189:ASP:OD2	4:A:265:HOH:O	2.09	0.69
2:D:524:SER:HB2	4:D:67:HOH:O	1.91	0.69
2:D:590:GLY:O	2:D:591:ILE:CB	2.40	0.69
1:B:163:LEU:HD12	1:B:166:LEU:HD12	1.73	0.69
1:B:125:ASP:HB3	4:B:251:HOH:O	1.94	0.68
2:D:441:LYS:HG3	2:D:588:TRP:O	1.92	0.68
2:D:503:PHE:CZ	2:D:512:MET:SD	2.87	0.68
1:B:163:LEU:HB2	1:B:166:LEU:HD12	1.75	0.67
2:C:475:ILE:HG13	2:C:536:LEU:HD23	1.76	0.66
1:B:134:PHE:CE2	1:B:197:THR:CG2	2.79	0.66
1:A:165:SER:CB	1:A:166:LEU:CD2	2.49	0.65
1:B:163:LEU:CD1	1:B:166:LEU:HD12	2.27	0.65
2:D:524:SER:CB	4:D:67:HOH:O	2.45	0.64
1:B:134:PHE:HZ	1:B:197:THR:HG22	1.57	0.63
2:D:590:GLY:O	2:D:591:ILE:HB	1.99	0.62
1:A:138:PHE:CE2	2:D:559:PRO:HB3	2.36	0.61
1:B:163:LEU:CD1	1:B:166:LEU:CD1	2.79	0.60
2:D:439:GLN:NE2	4:D:188:HOH:O	2.34	0.59
1:A:166:LEU:HD12	2:D:436:VAL:CG2	2.32	0.59
2:C:508:ASN:O	2:C:512:MET:HB3	2.02	0.59
1:B:165:SER:O	1:B:165:SER:OG	2.20	0.59
1:B:134:PHE:HZ	1:B:197:THR:HG23	1.67	0.58



		Interatomic	Clash	
Atom-1	Atom-2	distance $(\text{\AA})$	overlap (Å)	
1:B:95:LEU:HG	3:D:1:GOL:H2	1.86	0.58	
2:D:496:SER:HA	2:D:499:TYR:HD2	1.67	0.58	
1:B:134:PHE:CZ	1:B:197:THR:HG23	2.38	0.57	
2:D:530:HIS:CE1	4:D:129:HOH:O	2.58	0.57	
2:D:442:VAL:HB	4:D:188:HOH:O	2.04	0.57	
2:D:444:HIS:HD2	2:D:589:LEU:O	1.88	0.56	
1:A:124:MET:HE3	4:A:265:HOH:O	2.05	0.56	
2:C:583:LYS:NZ	4:C:78:HOH:O	2.25	0.56	
1:B:188:ASN:O	1:B:189:ASP:HB2	2.06	0.56	
1:A:166:LEU:CD1	2:D:436:VAL:CG2	2.83	0.55	
2:D:444:HIS:CD2	2:D:589:LEU:O	2.59	0.55	
1:B:125:ASP:CB	4:B:251:HOH:O	2.52	0.55	
2:D:472:ARG:O	2:D:476:GLU:HG2	2.06	0.55	
1:B:85:PRO:O	1:B:86:ALA:C	2.44	0.55	
2:C:440:LEU:HD21	2:C:589:LEU:CD1	2.36	0.54	
2:C:458:TYR:O	2:C:462:THR:HG23	2.07	0.54	
2:D:503:PHE:CE1	2:D:512:MET:SD	3.01	0.54	
1:A:147:LEU:C	1:A:147:LEU:HD23	2.29	0.53	
2:D:541:ARG:CD	4:D:160:HOH:O	2.56	0.53	
2:C:489:GLY:HA3	2:C:522:LEU:HD21	1.89	0.53	
2:D:530:HIS:HE1	4:D:129:HOH:O	1.89	0.53	
1:A:166:LEU:CD1	2:D:436:VAL:HG21	2.39	0.53	
1:A:86:ALA:HA	4:A:240:HOH:O	2.08	0.52	
1:A:88:ARG:HA	2:C:562:MET:CE	2.39	0.52	
2:C:537:GLU:OE2	2:C:540:LEU:HD23	2.10	0.52	
2:D:590:GLY:C	2:D:591:ILE:HG22	2.31	0.52	
2:C:496:SER:O	2:C:500:LEU:HG	2.11	0.51	
1:B:84:VAL:CB	1:B:85:PRO:CD	2.71	0.51	
2:D:499:TYR:O	2:D:503:PHE:HB2	2.11	0.50	
2:D:520:GLU:OE1	2:D:521:ARG:HG3	2.12	0.50	
2:D:449:ASP:HA	2:D:452:ARG:NH1	2.27	0.49	
1:B:134:PHE:HD1	1:B:141:LEU:CD1	2.23	0.49	
4:B:270:HOH:O	3:D:1:GOL:H31	2.12	0.49	
2:D:496:SER:HA	2:D:499:TYR:CD2	2.47	0.49	
1:B:166:LEU:C	1:B:167:PRO:O	2.46	0.49	
1:B:139:ASP:HA	4:B:250:HOH:O	2.13	0.49	
1:A:166:LEU:HD12	2:D:436:VAL:HG22	1.95	0.48	
2:C:436:VAL:HG13	2:C:571:TYR:CD1	2.48	0.48	
1:A:137:ILE:HG12	1:A:142:GLU:HB2	1.96	0.48	
1:A:164:PRO:O	1:A:165:SER:C	2.50	0.48	
2:C:440:LEU:HD21	2:C:589:LEU:HD12	1.94	0.48	



	1 · · · · ·	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
2:C:505:ARG:HA	4:C:30:HOH:O	2.14	0.48
1:B:137:ILE:O	1:B:138:PHE:HB2	2.14	0.48
2:D:470:MET:O	2:D:473:THR:HG22	2.13	0.47
1:A:138:PHE:O	4:A:240:HOH:O	2.20	0.47
2:D:541:ARG:HD3	4:D:160:HOH:O	2.13	0.47
1:A:188:ASN:O	1:A:189:ASP:HB2	2.14	0.47
1:A:108:LYS:HD2	1:A:121:GLN:CD	2.35	0.47
1:B:114:PRO:CA	1:B:166:LEU:HD13	2.42	0.47
1:B:108:LYS:HD3	1:B:121:GLN:CD	2.35	0.47
2:D:473:THR:HA	2:D:476:GLU:CG	2.45	0.47
2:D:505:ARG:HG3	2:D:506:GLU:HG2	1.97	0.47
2:D:473:THR:HA	2:D:476:GLU:HG3	1.96	0.47
1:A:90:LEU:HD12	1:A:136:VAL:HG13	1.97	0.46
2:C:464:THR:OG1	2:C:547:ASN:OD1	2.32	0.46
1:B:85:PRO:HG2	1:B:88:ARG:HD3	1.98	0.46
2:D:466:GLN:HB2	4:D:261:HOH:O	2.16	0.46
2:D:504:ARG:HG3	4:D:81:HOH:O	2.15	0.46
2:D:568:ARG:HH22	3:D:1:GOL:H11	1.81	0.46
2:D:486:GLU:O	2:D:490:GLN:HG2	2.16	0.46
2:D:555:ASN:HA	2:D:558:LYS:HD2	1.98	0.46
2:C:490:GLN:O	2:C:494:LYS:HB2	2.15	0.45
2:D:574:TRP:CH2	2:D:578:LYS:HG3	2.51	0.45
2:C:506:GLU:HB3	2:C:508:ASN:H	1.81	0.45
2:C:571:TYR:CD2	1:B:167:PRO:HB2	2.51	0.45
1:A:88:ARG:HA	2:C:562:MET:SD	2.56	0.45
1:A:165:SER:HA	4:A:2:HOH:O	2.16	0.45
1:B:168:GLY:HA2	4:B:236:HOH:O	2.17	0.45
2:D:583:LYS:O	2:D:587:GLU:HG3	2.17	0.44
1:A:166:LEU:HD12	2:D:436:VAL:HG21	1.99	0.44
2:C:475:ILE:HG13	2:C:536:LEU:CD2	2.45	0.44
2:D:442:VAL:CB	4:D:188:HOH:O	2.65	0.44
2:D:439:GLN:HG2	2:D:571:TYR:CE1	2.53	0.44
1:B:95:LEU:HD23	1:B:98:MET:HE3	1.99	0.44
2:D:581:ARG:NH2	4:D:52:HOH:O	2.50	0.44
2:D:507:GLY:O	2:D:511:GLU:HB2	2.18	0.43
1:A:182:ILE:O	1:A:186:GLU:HG3	2.19	0.43
2:D:503:PHE:HZ	2:D:512:MET:SD	2.39	0.43
2:C:447:TYR:CD1	2:C:568:ARG:NH1	2.87	0.43
1:B:193:ARG:HG2	4:B:249:HOH:O	2.19	0.43
2:D:574:TRP:CZ3	2:D:578:LYS:HG3	2.54	0.43
2:D:584:LYS:HA	2:D:584:LYS:HD2	1.84	0.42



Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:B:165:SER:HB3	4:B:33:HOH:O	2.19	0.42
2:D:478:PHE:CD1	2:D:529:ILE:HD11	2.54	0.42
1:B:147:LEU:HD23	1:B:147:LEU:C	2.40	0.42
2:C:489:GLY:HA2	2:C:522:LEU:CD2	2.43	0.42
1:B:137:ILE:HD12	1:B:137:ILE:HA	1.94	0.42
1:B:127:ASN:ND2	1:B:152:GLU:OE2	2.51	0.42
2:D:541:ARG:HD2	4:D:160:HOH:O	2.17	0.41
1:B:106:ILE:HD12	1:B:122:ALA:HB2	2.02	0.41
1:B:140:ARG:HB3	4:B:275:HOH:O	2.19	0.41
1:A:145:ILE:HD11	2:C:566:LYS:HD2	2.02	0.41
2:D:503:PHE:HB3	4:D:81:HOH:O	2.21	0.41
1:B:102:TRP:CZ3	1:B:104:MET:HG3	2.56	0.41
1:B:134:PHE:HB2	1:B:141:LEU:HD11	2.03	0.40
1:A:108:LYS:HD2	1:A:121:GLN:NE2	2.37	0.40
1:A:123:ILE:HG13	1:A:157:VAL:CG1	2.52	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	118/164~(72%)	116~(98%)	2(2%)	0	100	100
1	В	117/164~(71%)	114 (97%)	3~(3%)	0	100	100
2	$\mathbf{C}$	161/170~(95%)	158~(98%)	3~(2%)	0	100	100
2	D	161/170~(95%)	155~(96%)	6 (4%)	0	100	100
All	All	557/668~(83%)	543 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.



#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	104/143~(73%)	103~(99%)	1 (1%)	76 87
1	В	103/143~(72%)	102 (99%)	1 (1%)	76 87
2	С	150/157~(96%)	149 (99%)	1 (1%)	84 92
2	D	150/157~(96%)	148 (99%)	2(1%)	69 82
All	All	507/600~(84%)	502 (99%)	5 (1%)	76 87

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	139	ASP
2	С	568	ARG
1	В	165	SER
2	D	450	LYS
2	D	522	LEU

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such side chains are listed below:

Mol	Chain	Res	Type
2	С	513	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	True	Chain	Dog Link		B	ond leng	$\mathbf{gths}$	E	Bond ang	gles
	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z  > 2
3	GOL	С	1	-	5,5,5	0.48	0	$5,\!5,\!5$	0.72	0
3	GOL	D	1	-	$5,\!5,\!5$	0.37	0	$5,\!5,\!5$	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	С	1	-	-	4/4/4/4	-
3	GOL	D	1	-	-	1/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	С	1	GOL	O1-C1-C2-C3
3	С	1	GOL	C1-C2-C3-O3
3	С	1	GOL	O1-C1-C2-O2
3	С	1	GOL	O2-C2-C3-O3
3	D	1	GOL	O1-C1-C2-O2

There are no ring outliers.



1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	1	GOL	3	0

# 5.7 Other polymers (i)

There are no such residues in this entry.

# 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

# 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	120/164~(73%)	0.37	3 (2%) 57 64	17, 30, 57, 75	0
1	В	119/164~(72%)	0.35	3 (2%) 57 64	13, 28, 57, 68	0
2	С	163/170~(95%)	1.07	29 (17%) 1 1	19, 48, 137, 142	0
2	D	163/170~(95%)	1.31	35 (21%) 0 1	17, 53, 137, 141	0
All	All	565/668~(84%)	0.84	70 (12%) 4 5	13, 41, 131, 142	0

All (70) RSRZ outliers are listed below:

Mol	Chain	$\mathbf{Res}$	Type	RSRZ
2	D	507	GLY	11.7
2	С	505	ARG	9.4
2	D	500	LEU	8.4
2	С	503	PHE	8.0
2	D	518	ASN	7.7
2	D	506	GLU	7.6
2	С	507	GLY	7.1
2	С	501	GLU	6.9
2	D	509	GLU	6.7
2	D	517	LEU	6.5
2	С	510	LYS	6.5
2	D	505	ARG	6.3
2	С	513	GLN	6.3
2	D	521	ARG	6.1
2	D	503	PHE	5.9
2	С	514	ARG	5.5
2	D	508	ASN	5.5
2	С	516	LEU	5.3
1	А	83	SER	5.2
2	D	504	ARG	5.2
2	D	512	MET	5.2



3L4Q

Mol	Chain	Res	Type	RSRZ
2	D	514	ARG	5.0
2	D	499	TYR	4.9
2	С	506	GLU	4.8
2	С	497	LYS	4.7
2	С	515	ILE	4.6
2	С	504	ARG	4.6
2	С	470	MET	4.4
2	С	511	GLU	4.3
2	D	502	ARG	4.2
2	D	495	SER	4.2
2	С	499	TYR	4.1
2	С	495	SER	4.0
2	D	510	LYS	4.0
1	А	84	VAL	3.8
2	D	430	GLU	3.7
2	D	496	SER	3.6
2	D	591	ILE	3.6
2	D	523	LYS	3.6
2	D	494	LYS	3.6
2	С	512	MET	3.5
2	С	498	GLU	3.5
2	С	430	GLU	3.5
1	В	138	PHE	3.4
2	D	525	ARG	3.4
2	D	490	GLN	3.3
2	С	508	ASN	3.0
2	D	429	LYS	3.0
2	С	522	LEU	2.9
2	D	466	GLN	2.9
2	С	502	ARG	2.9
2	D	562	MET	2.7
1	В	202	ALA	2.6
2	D	529	ILE	2.6
2	D	497	LYS	2.6
2	С	518	ASN	2.6
1	А	165	SER	2.5
2	С	509	GLU	2.5
2	D	589	LEU	2.5
2	D	501	GLU	2.4
2	С	492	GLN	2.3
2	D	470	MET	2.3
2	D	511	GLU	2.2



Mol	Chain	Res	Type	RSRZ
2	С	525	ARG	2.2
2	D	515	ILE	2.2
1	В	200	ARG	2.2
2	С	521	ARG	2.2
2	С	459	GLU	2.1
2	D	493	GLU	2.1
2	С	589	LEU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
3	GOL	С	1	6/6	0.55	0.30	20,20,20,20	0
3	GOL	D	1	6/6	0.90	0.18	20,20,20,20	0

#### 6.5 Other polymers (i)

There are no such residues in this entry.

