



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 6, 2020 – 05:36 PM BST

PDB ID : 1L5R  
Title : Human liver glycogen phosphorylase a complexed with riboflavin, N-Acetyl-b  
eta-D-Glucopyranosylamine and CP-403,700  
Authors : Ekstrom, J.L.; Pauly, T.A.; Carty, M.D.; Soeller, W.C.; Culp, J.; Danley,  
D.E.; Hoover, D.J.; Treadway, J.L.; Gibbs, E.M.; Fletterick, R.J.; Day, Y.S.N.;  
Myszka, D.G.; Rath, V.L.  
Deposited on : 2002-03-07  
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	847	 65% 26% • 7%
1	B	847	 65% 27% • 7%

## 2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 13497 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called glycogen phosphorylase, liver form.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	790	Total 6417	C 4125	N 1089	O 1174	S 29	0	0	0
1	B	791	Total 6423	C 4128	N 1090	O 1176	S 29	0	0	0

- Molecule 2 is N-acetyl-beta-D-glucopyranosylamine (three-letter code: NBG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	Total 15	C 8	N 1	O 6	0	0

- Molecule 3 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: C<sub>8</sub>H<sub>10</sub>NO<sub>6</sub>P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	15	8	1	5	1	0	0
3	B	1	15	8	1	5	1	0	0

- Molecule 4 is [5-CHLORO-1H-INDOL-2-CARBONYL-PHENYLALANINYL]-AZETIDINE-3-CARBOXYLIC ACID (three-letter code: 700) (formula: C<sub>22</sub>H<sub>20</sub>ClN<sub>3</sub>O<sub>4</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Cl	N	O		
4	A	1	30	22	1	3	4	0	0

- Molecule 5 is RIBOFLAVIN (three-letter code: RBF) (formula:  $C_{17}H_{20}N_4O_6$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	N			O
5	A	1	27	17	4	6	0	0

- Molecule 6 is (4R)-2-METHYLPENTANE-2,4-DIOL (three-letter code: MRD) (formula:  $C_6H_{14}O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
6	B	1	8	6	2	0	0

- Molecule 7 is water.


<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
7	A	307	Total 307	O 307	0	0
7	B	240	Total 240	O 240	0	0

### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: glycogen phosphorylase, liver form



- Molecule 1: glycogen phosphorylase, liver form



D785	T681	V565	C445	F329	T637	L102
K786	M692	Q566	G448	P330	V238	M106
Q789	D693	V567	V455	Q336	L243	E110
M792	A695	H571	H459	L337	W244	Y113
M793	M696	E572	V463	H341	S245	D118
P794	V697	Y573	K469	P342	A246	E121
L802	E701	R574	D470	A345	R247	E124
D814	L708	R575	K469	I346	N250	I125
R815	F709	Q576	D470	E382	PHE	E126
R818	I710	L577	E473	W361	ASN	A129
W825	M713	D593	L474	Y374	LEU	ASN
N826	R714	P594	T487	E386	ARG	GLY
W827	I715	V599	P488	R386	ASP	G135
E828	D716	P600	R489	V389	ASN	L136
P829	D717	I604	L492	K394	VAL	S146
ASP	K724	I605	L493	H399	GLY	L150
LEU	A728	G606	L494	I402	P281	Y155
LYS	Y732	Y613	L499	I403	W282	Y157
ILE	E733	H614	L502	I406	M284	R160
SER	A734	M615	E505	K409	F285	Q168
LEU	F736	M618	K506	R410	V300	K169
SER	E737	L622	V512	R411	I170	I170
ASN	L738	D628	K513	L412	A301	Q175
GLU	K739	D633	F524	R413	S302	S192
SER	L740	P634	L525	I414	T303	M197
ASN	V741	W636	D528	V415	L304	L198
LYS	D746	G637	R522	F418	Q305	Y203
VAL	P752	L645	K538	R419	I308	G204
ASN	K753	W646	K542	K420	R309	K205
ASN	Q754	M647	L543	D421	R310	H208
LYS	P755	Y648	R544	V422	F311	T213
VAL	D756	Y648	F545	R423	K312	I216
ASN	L757	Y648	S546	R424	K314	L224
GLY	F758	Y648	Q547	L425	A313	P225
ASN	K759	L652	F548	R426	S314	Y233
LYS	D760	L652	F548	R427	K315	N236
ASN	I761	K655	T551	E432	F316	
ASN	I762	K655	K554	E433	GLY	
ASN	N763	I657	V555	E433	SER	
ASN	M764	P658	W556	E433	THR	
ASN	L765	I666	K556	E433	ARG	
ASN	D769	S674	I557	E433	GLY	
ASN	F774	L683	N558	E433	ALA	
ASN	Y777	I689	I558	E433	GLY	
ASN	C783	Q784	I558	E433	THR	
ASN	Q784		S561	E433	THR	



## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	124.42Å 124.42Å 124.01Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	55.61 – 2.10	Depositor
% Data completeness (in resolution range)	95.5 (55.61-2.10)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.09	Depositor
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.247 , 0.283	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	13497	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MRD, RBF, 700, NBG, PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.44	0/6561	0.65	1/8873 (0.0%)
1	B	0.43	0/6567	0.65	1/8881 (0.0%)
All	All	0.43	0/13128	0.65	2/17754 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	129	ALA	N-CA-C	-5.82	95.29	111.00
1	A	129	ALA	N-CA-C	-5.38	96.47	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6417	0	6412	191	0
1	B	6423	0	6417	183	0
2	A	15	0	15	0	0
3	A	15	0	7	0	0
3	B	15	0	7	0	0
4	A	30	0	18	0	0
5	A	27	0	20	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	8	0	14	0	0
7	A	307	0	0	28	0
7	B	240	0	0	18	0
All	All	13497	0	12910	368	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 368 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:110:GLU:HA	7:B:2505:HOH:O	1.65	0.93
1:B:113:TYR:HB3	7:B:2505:HOH:O	1.69	0.93
1:A:213:THR:HB	7:A:2504:HOH:O	1.70	0.91
1:A:710:ILE:HD13	7:A:2205:HOH:O	1.72	0.89
1:A:798:ASN:HB3	7:A:2061:HOH:O	1.72	0.87

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	784/847 (93%)	742 (95%)	39 (5%)	3 (0%)	34 32
1	B	785/847 (93%)	742 (94%)	39 (5%)	4 (0%)	29 26
All	All	1569/1694 (93%)	1484 (95%)	78 (5%)	7 (0%)	34 32

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	555	VAL

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	555	VAL
1	A	554	LYS
1	B	554	LYS
1	A	342	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	692/740 (94%)	662 (96%)	30 (4%)	29	29
1	B	693/740 (94%)	661 (95%)	32 (5%)	27	26
All	All	1385/1480 (94%)	1323 (96%)	62 (4%)	27	27

5 of 62 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	815	ARG
1	B	102	LEU
1	B	683	LEU
1	B	78	CYS
1	B	243	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	459	HIS
1	A	576	GLN
1	B	369	GLN
1	A	369	GLN
1	B	250	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	PLP	A	860	1	15,15,16	2.11	4 (26%)	20,22,23	1.34	3 (15%)
3	PLP	B	860	1	15,15,16	2.00	3 (20%)	20,22,23	0.91	0
2	NBG	A	861	-	15,15,15	1.32	3 (20%)	21,21,21	1.24	2 (9%)
6	MRD	B	902	-	7,7,7	0.71	0	9,10,10	0.72	0
5	RBF	A	859	-	27,29,29	2.48	12 (44%)	33,43,43	3.01	11 (33%)
4	700	A	862	-	28,33,33	1.91	9 (32%)	32,47,47	1.52	4 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PLP	A	860	1	-	0/6/6/8	0/1/1/1
3	PLP	B	860	1	-	2/6/6/8	0/1/1/1
2	NBG	A	861	-	-	0/6/26/26	0/1/1/1
6	MRD	B	902	-	-	2/5/5/5	-
5	RBF	A	859	-	-	1/14/14/14	0/3/3/3
4	700	A	862	-	-	0/15/32/32	0/4/4/4

The worst 5 of 31 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	859	RBF	C4-C4A	6.33	1.52	1.41
3	B	860	PLP	C4A-C4	-5.49	1.40	1.51
3	A	860	PLP	C4A-C4	-5.05	1.41	1.51
5	A	859	RBF	C1'-N10	5.00	1.53	1.48
5	A	859	RBF	C4A-C10	4.37	1.43	1.38

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	859	RBF	C4-N3-C2	8.15	122.03	115.14
5	A	859	RBF	C4-C4A-C10	-6.80	115.45	119.95
5	A	859	RBF	C10-C4A-N5	6.44	125.71	121.26
5	A	859	RBF	O5'-C5'-C4'	-5.76	98.52	111.07
5	A	859	RBF	O4'-C4'-C3'	4.96	121.16	109.10

There are no chirality outliers.


All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	860	PLP	C6-C5-C5A-O4P
3	B	860	PLP	C4-C5-C5A-O4P
6	B	902	MRD	C2-C3-C4-C5
6	B	902	MRD	C2-C3-C4-O4
5	A	859	RBF	O3'-C3'-C4'-C5'

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.