



wwPDB NMR Structure Validation Summary Report ⓘ

Nov 7, 2023 – 07:59 AM EST

PDB ID : 2L5E
BMRB ID : 17270
Title : Complex between BD1 of Brd3 and GATA-1 C-tail
Authors : Gamsjaeger, R.; Webb, S.R.; Lamonica, J.M.; Blobel, G.A.; Mackay, J.P.
Deposited on : 2010-10-31

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<https://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 87%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	128	
2	B	13	

2 Ensemble composition and analysis

This entry contains 20 models. Model 20 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:26-A:147 (122)	0.68	20

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 4 single-model clusters were found.

Cluster number	Models
1	2, 8, 12, 13, 16, 18, 20
2	3, 6, 10, 14
3	5, 7, 19
4	4, 15
Single-model clusters	1; 9; 11; 17

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2202 atoms, of which 1102 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Bromodomain-containing protein 3.

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		S
1	A	127	2092	677	1042	172	192	9	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	20	GLY	-	expression tag	UNP Q3TUI3
A	21	PRO	-	expression tag	UNP Q3TUI3
A	22	LEU	-	expression tag	UNP Q3TUI3
A	23	GLY	-	expression tag	UNP Q3TUI3
A	24	SER	-	expression tag	UNP Q3TUI3

- Molecule 2 is a protein called GATA-1.


Mol	Chain	Residues	Atoms					Trace
			Total	C	H	N	O	
2	B	6	110	32	60	10	8	0

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Bromodomain-containing protein 3




- Molecule 2: GATA-1



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 20. Colouring as in section 4.1 above.

- Molecule 1: Bromodomain-containing protein 3



- Molecule 2: GATA-1



LYS	
ALA	
SER	6311
	6312
	6313
	6314
	6315
	6316
ARG	
GLY	
SER	
ASN	

5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics*.

Of the 20 calculated structures, 20 were deposited, based on the following criterion: *all calculated structures submitted*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	refinement	1.1

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1615
Number of shifts mapped to atoms	1565
Number of unparsed shifts	0
Number of shifts with mapping errors	50
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	87%

6 Model quality i

6.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: ALY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.38±0.06	1±1/1043 (0.1± 0.1%)	0.41±0.01	0±0/1412 (0.0± 0.0%)
All	All	0.39	19/20860 (0.1%)	0.41	0/28240 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.1±0.5
All	All	0	3

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	95	TYR	CE2-CZ	-7.71	1.28	1.38	12	2
1	A	60	TYR	CE2-CZ	-7.62	1.28	1.38	3	7
1	A	60	TYR	CE1-CZ	7.29	1.48	1.38	3	7
1	A	95	TYR	CE1-CZ	7.26	1.48	1.38	7	2
1	A	94	TYR	CE2-CZ	-5.08	1.31	1.38	12	1

There are no bond-angle outliers.

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	95	TYR	Sidechain	2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group	Models (Total)
1	A	94	TYR	Sidechain	1

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1016	1009	1009	11±3
2	B	0	0	0	0±0
All	All	20320	20180	20180	229

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

5 of 83 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:36:THR:HG22	1:A:146:GLU:HG3	0.78	1.53	19	10
1:A:118:PRO:HA	1:A:123:VAL:HG11	0.69	1.63	12	6
1:A:39:LEU:HA	1:A:98:ALA:HB2	0.62	1.70	17	5
1:A:90:LEU:HG	1:A:101:CYS:SG	0.61	2.36	3	1
1:A:89:ARG:HB3	1:A:95:TYR:CE1	0.61	2.31	7	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	121/128 (95%)	110±2 (91±2%)	11±2 (9±2%)	1±1 (1±1%)	29 74
2	B	0	-	-	-	-
All	All	2420/2820 (86%)	2194 (91%)	212 (9%)	14 (1%)	29 74

5 of 8 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	82	ASP	5
1	A	144	GLU	2
1	A	71	PRO	2
1	A	80	PRO	1
1	A	33	GLY	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	114/118 (97%)	80±4 (70±3%)	34±4 (30±3%)	1 17
2	B	0	-	-	-
All	All	2280/2500 (91%)	1600 (70%)	680 (30%)	1 17

5 of 79 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	50	LEU	20
1	A	119	THR	20
1	A	39	LEU	19
1	A	87	LYS	19
1	A	105	PHE	19

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types,

if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
2	ALY	B	312	2	10,11,12	0.41±0.01	0±0 (0±0%)
2	ALY	B	315	2	10,11,12	0.42±0.02	0±0 (0±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
2	ALY	B	312	2	7,12,14	0.49±0.03	0±0 (0±0%)
2	ALY	B	315	2	7,12,14	0.49±0.03	0±0 (0±0%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ALY	B	315	2	-	0±0,9,10,12	-
2	ALY	B	312	2	-	0±0,9,10,12	-

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation i

The completeness of assignment taking into account all chemical shift lists is 87% for the well-defined parts and 84% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping i

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1615
Number of shifts mapped to atoms	1565
Number of unparsed shifts	0
Number of shifts with mapping errors	50
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	2

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

- No matching atom found in the structure. First 5 (of 50) occurrences are reported below.

List ID	Chain	Res	Type	Atom	Shift Data		
					Value	Uncertainty	Ambiguity
1	UNMAPPED	5	ALY	H	8.028	0.006	1
1	UNMAPPED	5	ALY	HA	4.163	0.003	1
1	UNMAPPED	5	ALY	HB2	1.675	0.013	2
1	UNMAPPED	5	ALY	HB3	1.84	0.018	2
1	UNMAPPED	5	ALY	HE2	3.36	0.000	2
1	UNMAPPED	5	ALY	HG2	1.285	0.019	2
1	UNMAPPED	5	ALY	HG3	1.378	0.003	2
1	UNMAPPED	5	ALY	HZ1	8.006	0.011	1
1	UNMAPPED	5	ALY	HZ2	8.006	0.011	1
1	UNMAPPED	5	ALY	HZ3	8.006	0.011	1
1	UNMAPPED	5	ALY	HH31	2.094	0.010	2
1	UNMAPPED	5	ALY	HH32	2.094	0.010	2
1	UNMAPPED	5	ALY	HH33	2.094	0.010	2
1	UNMAPPED	6	GLY	H	8.594	0.016	1

Continued on next page...

Continued from previous page...

List ID	Chain	Res	Type	Atom	Shift Data		
					Value	Uncertainty	Ambiguity
1	UNMAPPED	6	GLY	HA2	3.875	0.005	2
1	UNMAPPED	7	LYS	H	8.395	0.006	1
1	UNMAPPED	7	LYS	HA	4.418	0.004	1
1	UNMAPPED	7	LYS	HB2	1.801	0.008	2
1	UNMAPPED	7	LYS	HB3	1.902	0.008	2
1	UNMAPPED	7	LYS	HG2	1.494	0.000	2
1	UNMAPPED	8	ALY	H	7.867	0.009	1
1	UNMAPPED	8	ALY	HA	4.357	0.002	1
1	UNMAPPED	8	ALY	HB2	1.767	0.009	2
1	UNMAPPED	8	ALY	HB3	1.878	0.011	2
1	UNMAPPED	8	ALY	HD2	1.485	0.018	2
1	UNMAPPED	8	ALY	HD3	1.636	0.005	2
1	UNMAPPED	8	ALY	HE2	3.133	0.019	2
1	UNMAPPED	8	ALY	HG2	1.22	0.001	2
1	UNMAPPED	8	ALY	HG3	1.287	0.015	2
1	UNMAPPED	8	ALY	HZ1	7.955	0.015	1
1	UNMAPPED	8	ALY	HZ2	7.955	0.015	1
1	UNMAPPED	8	ALY	HZ3	7.955	0.015	1
1	UNMAPPED	8	ALY	HM1	2.072	0.013	2
1	UNMAPPED	9	LEU	H	8.353	0.004	1
1	UNMAPPED	9	LEU	HA	4.347	0.000	1
1	UNMAPPED	9	LEU	HB2	1.784	0.006	2
1	UNMAPPED	9	LEU	HB3	1.853	0.000	2
1	UNMAPPED	10	ARG	H	8.456	0.002	1
1	UNMAPPED	10	ARG	HA	4.293	0.005	1
1	UNMAPPED	10	ARG	HB2	1.751	0.000	2
1	UNMAPPED	10	ARG	HB3	1.83	0.000	2
1	UNMAPPED	10	ARG	HD2	3.231	0.002	2
1	UNMAPPED	10	ARG	HE	7.25	0.007	1
1	UNMAPPED	10	ARG	HG2	1.563	0.000	2
1	UNMAPPED	13	ASN	H	8.183	0.001	1
1	UNMAPPED	13	ASN	HA	4.498	0.002	1
1	UNMAPPED	13	ASN	HB2	2.79	0.003	2
1	UNMAPPED	13	ASN	HB3	2.689	0.001	2
1	UNMAPPED	13	ASN	HD21	6.831	0.010	2
1	UNMAPPED	13	ASN	HD22	7.528	0.010	2

7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	127	-0.52 ± 0.19	Should be checked
$^{13}\text{C}_\beta$	123	0.30 ± 0.13	None needed (< 0.5 ppm)
$^{13}\text{C}'$	124	-0.02 ± 0.17	None needed (< 0.5 ppm)
^{15}N	117	0.32 ± 0.39	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 87%, i.e. 1516 atoms were assigned a chemical shift out of a possible 1747. 0 out of 17 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	590/596 (99%)	236/238 (99%)	241/244 (99%)	113/114 (99%)
Sidechain	811/987 (82%)	525/639 (82%)	267/310 (86%)	19/38 (50%)
Aromatic	115/164 (70%)	49/79 (62%)	61/79 (77%)	5/6 (83%)
Overall	1516/1747 (87%)	810/956 (85%)	569/633 (90%)	137/158 (87%)

7.1.4 Statistically unusual chemical shifts [i](#)


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	52	LYS	HD2	3.08	0.58 – 2.64	7.2
1	A	130	GLU	HB3	0.65	0.95 – 3.05	-6.4

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:



Random coil index (RCI) for chain UNMAPPED:

