

Full wwPDB X-ray Structure Validation Report (i)

Dec 10, 2022 – 02:30 PM EST

1L8D
Rad50 coiled-coil Zn hook
Hopfner, K.P.; Tainer, J.A.
2002-03-20
2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.31.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# {\rm Entries})$	$(\# { m Entries}, { m resolution} { m range}({ m \AA}))$
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain			
1	А	112	62%	27%	• 8%	
1	В	112	57%	30%	•• 9%	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	CIT	А	507	-	Х	-	-



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 1857 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called DNA double-strand break repair rad50 ATPase.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Δ	103	Total	С	Ν	0	S	-91	0	0
	1 A	105	850	525	156	166	3	21	0	0
1	р	102	Total	С	Ν	0	S	53	0	0
1	D	102	839	518	152	165	4		0	U

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
А	395	MET	-	initiating methionine	UNP P58301
В	395	MET	-	initiating methionine	UNP P58301

• Molecule 2 is MERCURY (II) ION (three-letter code: HG) (formula: Hg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	Total Hg 1 1	0	0


• Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O_4P).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
3	А	1	Total O P 5 4 1	0	0
3	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
3	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
3	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
3	В	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	А	1	Total 13	C 6	O 7	0	0

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	74	Total O 74 74	0	0
5	В	50	$\begin{array}{cc} {\rm Total} & {\rm O} \\ 50 & 50 \end{array}$	0	0




3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: DNA double-strand break repair rad50 ATPase



• Molecule 1: DNA double-strand break repair rad50 ATPase





4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	31.96Å 77.87Å 53.33Å	Depositor
a, b, c, α , β , γ	90.00° 91.55° 90.00°	Depositor
Resolution (Å)	8.00 - 2.20	Depositor
% Data completeness	(Not available) (8.00-2.20)	Depositor
(in resolution range)	(1101 available) (0.00 2.20)	Depositor
R_{merge}	(Not available)	Depositor
R _{sym}	(Not available)	Depositor
Refinement program	CNS	Depositor
R, R_{free}	0.225 , 0.278	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	1857	wwPDB-VP
Average B, all atoms $(Å^2)$	38.0	wwPDB-VP



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, HG, CIT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles		
	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.52	0/853	0.70	1/1132~(0.1%)	
1	В	0.50	0/842	0.95	3/1117~(0.3%)	
All	All	0.51	0/1695	0.83	4/2249~(0.2%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	В	0	1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
1	В	489	ARG	NE-CZ-NH1	19.43	130.01	120.30
1	В	489	ARG	NE-CZ-NH2	-12.91	113.84	120.30
1	В	489	ARG	NH1-CZ-NH2	-7.06	111.64	119.40
1	А	497	ARG	NE-CZ-NH2	-6.64	116.98	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	В	489	ARG	Sidechain



5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	850	0	901	21	0
1	В	839	0	886	30	0
2	А	1	0	0	0	0
3	А	25	0	0	0	0
3	В	5	0	0	1	0
4	А	13	0	5	0	0
5	А	74	0	0	6	0
5	В	50	0	0	5	0
All	All	1857	0	1792	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:398:LEU:HA	5:B:20:HOH:O	1.58	1.01
1:B:492:ASP:HB2	5:B:123:HOH:O	1.64	0.97
1:A:405:LYS:HE3	1:A:409:ILE:HD11	1.52	0.89
1:A:437:LEU:HD21	1:A:459:LEU:HB3	1.62	0.81
1:B:452:THR:HG21	5:B:118:HOH:O	1.85	0.75
1:B:489:ARG:HA	5:B:123:HOH:O	1.86	0.75
1:A:399:LEU:HG	1:A:403:GLU:OE1	1.88	0.72
1:A:454:GLU:HG3	5:A:88:HOH:O	1.88	0.72
1:A:474:LEU:O	1:A:478:ILE:HG12	1.95	0.66
1:B:467:LEU:O	1:B:471:LYS:HG3	1.96	0.65
1:A:459:LEU:HD21	1:B:447:CYS:HA	1.78	0.64
1:A:467:LEU:O	1:A:471:LYS:HG3	2.00	0.61
1:B:487:GLU:O	1:B:491:ILE:HG12	2.01	0.60
1:A:413:ARG:CZ	5:A:93:HOH:O	2.49	0.59
1:B:452:THR:HG23	1:B:454:GLU:H	1.68	0.58
1:B:412:GLU:O	1:B:416:ILE:HD13	2.05	0.57
1:B:469:ASN:O	1:B:472:ASN:HB3	2.04	0.56
1:B:437:LEU:HD11	1:B:459:LEU:HB3	1.89	0.55
1:B:413:ARG:NH2	1:B:414:ASN:OD1	2.39	0.55

Continued on next page...



A 1 - 1		Interatomic	Clash
Atom-1	Atom-2	distance (\AA)	overlap (Å)
1:B:406:LYS:HE2	1:B:492:ASP:OD1	2.07	0.55
1:B:409:ILE:HD13	1:B:487:GLU:HG2	1.89	0.54
1:B:437:LEU:HD22	1:B:456:ARG:HG3	1.88	0.54
1:B:451:LEU:CD1	1:B:456:ARG:HB2	2.40	0.51
1:B:416:ILE:HG21	1:B:481:LYS:HB2	1.90	0.51
1:B:451:LEU:HD11	1:B:456:ARG:HB2	1.93	0.50
1:B:419:ARG:NH2	1:B:477:LEU:HD21	2.26	0.50
1:B:443:LYS:HD3	1:B:450:GLU:HA	1.93	0.50
1:A:396:LYS:N	5:A:23:HOH:O	2.44	0.49
1:B:423:LEU:HD21	1:B:470:SER:O	2.14	0.48
1:B:489:ARG:O	1:B:493:MET:HG3	2.14	0.48
1:B:428:GLY:O	1:B:432:THR:HG23	2.13	0.47
1:A:413:ARG:NH2	5:A:93:HOH:O	2.46	0.47
1:B:409:ILE:CG2	1:B:488:LEU:HG	2.44	0.47
1:A:405:LYS:O	1:A:409:ILE:HG12	2.14	0.47
1:A:406:LYS:HE2	5:A:60:HOH:O	2.15	0.47
1:A:418:GLN:NE2	5:A:46:HOH:O	2.48	0.46
1:B:419:ARG:NH2	5:B:62:HOH:O	2.32	0.45
1:B:452:THR:HG22	1:B:455:HIS:H	1.82	0.45
1:B:419:ARG:CZ	1:B:477:LEU:HD21	2.47	0.44
1:A:453:ASP:O	1:A:457:GLU:HG2	2.19	0.43
1:A:491:ILE:O	1:A:495:ILE:HG13	2.19	0.43
1:A:423:LEU:O	1:A:427:ILE:HG13	2.19	0.42
1:A:406:LYS:O	1:A:410:GLU:HG3	2.20	0.42
1:B:449:ARG:NH2	3:B:301:PO4:O2	2.48	0.42
1:A:409:ILE:HD12	1:A:487:GLU:OE1	2.21	0.41
1:A:430:LEU:HD23	1:A:430:LEU:HA	1.88	0.41
1:B:409:ILE:HG22	1:B:488:LEU:HG	2.01	0.41
1:B:491:ILE:O	1:B:495:ILE:HG13	2.21	0.41
1:A:465:LEU:HD23	1:A:465:LEU:HA	1.93	0.40
1:A:489:ARG:NH2	1:A:493:MET:SD	2.95	0.40

Continued from previous page...

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	А	101/112~(90%)	97~(96%)	4 (4%)	0	100	100
1	В	100/112~(89%)	97~(97%)	3~(3%)	0	100	100
All	All	201/224~(90%)	194 (96%)	7 (4%)	0	100	100

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Analysed Rotameric Outliers		Percentiles		
1	А	96/104~(92%)	88~(92%)	8 (8%)	11 11		
1	В	95/104 (91%)	90~(95%)	5 (5%)	22 27		
All	All	191/208~(92%)	178 (93%)	13 (7%)	16 17		

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	402	LEU
1	А	423	LEU
1	А	467	LEU
1	А	469	ASN
1	А	477	LEU
1	А	485	GLU
1	А	488	LEU
1	А	489	ARG
1	В	397	LYS
1	В	402	LEU
1	В	413	ARG
1	В	437	LEU
1	В	452	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:



Mol	Chain	Res	Type
1	А	418	GLN
1	А	468	ASN
1	А	472	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 8 ligands modelled in this entry, 1 is monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Turne	Chain	Res	Link	Bo	ond leng	ths	Bond angles		
IVIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PO4	А	306	-	4,4,4	1.52	0	$6,\!6,\!6$	0.44	0
3	PO4	А	303	-	4,4,4	1.53	0	$6,\!6,\!6$	0.44	0
3	PO4	А	302	-	4,4,4	1.40	0	6,6,6	0.44	0
4	CIT	А	507	-	12,12,12	2.48	8 (66%)	17,17,17	2.17	<mark>6 (35%)</mark>
3	PO4	А	305	-	4,4,4	1.37	0	6,6,6	0.47	0
3	PO4	А	304	-	4,4,4	1.31	0	$6,\!6,\!6$	0.45	0
3	PO4	В	301	-	4,4,4	1.75	1 (25%)	$6,\!6,\!6$	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.



'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	CIT	А	507	-	-	8/16/16/16	-

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	А	507	CIT	O1-C1	-3.63	1.10	1.22
4	А	507	CIT	O2-C1	-3.42	1.19	1.30
4	А	507	CIT	O3-C5	3.18	1.32	1.22
4	А	507	CIT	C4-C3	3.08	1.57	1.53
4	А	507	CIT	O4-C5	-3.00	1.20	1.30
4	А	507	CIT	C2-C3	2.91	1.57	1.53
4	А	507	CIT	O7-C3	2.07	1.47	1.43
3	В	301	PO4	P-O3	-2.05	1.48	1.54
4	А	507	CIT	O5-C6	2.02	1.28	1.22

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
4	А	507	CIT	O4-C5-C4	4.08	127.47	114.35
4	А	507	CIT	O5-C6-C3	-3.77	116.91	122.25
4	А	507	CIT	O3-C5-C4	-3.26	113.43	122.94
4	А	507	CIT	C3-C2-C1	3.11	121.34	113.81
4	А	507	CIT	O6-C6-C3	3.10	118.43	113.05
4	А	507	CIT	O2-C1-O1	2.43	129.35	123.30

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	А	507	CIT	C3-C4-C5-O3
4	А	507	CIT	C3-C4-C5-O4
4	А	507	CIT	O7-C3-C6-O5
4	А	507	CIT	C2-C3-C6-O6
4	А	507	CIT	C4-C3-C6-O5
4	А	507	CIT	C4-C3-C6-O6
4	А	507	CIT	C2-C3-C6-O5
4	А	507	CIT	O7-C3-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	В	301	PO4	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

