



Full wwPDB NMR Structure Validation Report ⓘ

Jun 4, 2023 – 06:02 PM EDT

PDB ID : 2LBK
BMRB ID : 17564
Title : Glycyl-tRNA(UCC)1B anticodon stem-loop from Staphylococcus epidermidis
Authors : Chang, A.T.; Nikonowicz, E.P.
Deposited on : 2011-03-31

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
BMRB Restraints Analysis : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.33


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 58%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
RNA backbone	4643	676

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	17	 76% 12% 12%

2 Ensemble composition and analysis

This entry contains 8 models. This entry does not contain polypeptide chains, therefore identification of well-defined residues and clustering analysis are not possible. All residues are included in the validation scores.

3 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 539 atoms, of which 185 are hydrogens and 0 are deuteriums.

- Molecule 1 is a RNA chain called RNA (5'-R(*GP*GP*GP*AP*CP*CP*UP*UP*CP*CP*CP*GP*GP*UP*CP*UP*C)-3').


Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		P
1	A	17	539	159	185	59	120	16	0

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: RNA (5'-R(*GP*GP*GP*AP*CP*CP*UP*UP*CP*CP*CP*GP*GP*UP*CP*UP*C)-3')



4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

- Molecule 1: RNA (5'-R(*GP*GP*GP*AP*CP*CP*UP*UP*CP*CP*CP*GP*GP*UP*CP*UP*C)-3')



4.2.2 Score per residue for model 2

- Molecule 1: RNA (5'-R(*GP*GP*GP*AP*CP*CP*UP*UP*CP*CP*CP*GP*GP*UP*CP*UP*C)-3')




4.2.3 Score per residue for model 3

- Molecule 1: RNA (5'-R(*GP*GP*GP*AP*CP*CP*UP*UP*CP*CP*CP*GP*GP*UP*CP*UP*C)-3')




4.2.4 Score per residue for model 4

- Molecule 1: RNA (5'-R(*GP*GP*GP*AP*CP*CP*UP*UP*CP*CP*CP*GP*GP*UP*CP*UP*C)-3')




4.2.5 Score per residue for model 5

- Molecule 1: RNA (5'-R(*GP*GP*GP*AP*CP*CP*UP*UP*CP*CP*CP*GP*GP*UP*CP*UP*C)-3')




4.2.6 Score per residue for model 6

- Molecule 1: RNA (5'-R(*GP*GP*GP*AP*CP*CP*UP*UP*CP*CP*CP*GP*GP*UP*CP*UP*C)-3')




4.2.7 Score per residue for model 7

- Molecule 1: RNA (5'-R(*GP*GP*GP*AP*CP*CP*UP*UP*CP*CP*CP*GP*GP*UP*CP*UP*C)-3')



4.2.8 Score per residue for model 8

- Molecule 1: RNA (5'-R(*GP*GP*GP*AP*CP*CP*UP*UP*CP*CP*CP*GP*GP*UP*CP*UP*C)-3')



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 50 calculated structures, 8 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	structure solution	
X-PLOR NIH	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	194
Number of shifts mapped to atoms	194
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	58%

6 Model quality i

6.1 Standard geometry i

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	354	185	185	4±1
All	All	2832	1480	1480	34

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:11:C:H2'	1:A:12:G:O4'	0.71	1.85	3	8
1:A:8:U:O2'	1:A:9:C:H3'	0.64	1.91	2	7
1:A:11:C:O2'	1:A:12:G:H5'	0.61	1.95	4	3
1:A:10:C:H2'	1:A:11:C:O4'	0.58	1.98	1	1
1:A:16:U:O2'	1:A:17:C:H5'	0.56	2.01	3	3
1:A:10:C:C4	1:A:11:C:C6	0.51	2.98	7	1
1:A:7:U:O2'	1:A:8:U:H5'	0.51	2.05	4	1
1:A:2:G:O2'	1:A:3:G:H5'	0.47	2.09	3	2
1:A:10:C:N4	1:A:11:C:C2	0.44	2.85	7	1
1:A:7:U:O4	1:A:8:U:C4	0.43	2.71	2	1
1:A:2:G:H2'	1:A:3:G:O4'	0.43	2.14	3	2
1:A:6:C:C5	1:A:7:U:C4	0.42	3.07	6	2
1:A:6:C:C4	1:A:7:U:C4	0.41	3.09	6	1
1:A:8:U:H6	1:A:8:U:O5'	0.40	2.00	8	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

6.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

6.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers	Suiteness
1	A	16/17 (94%)	2±1 (12±4%)	0±0 (0±0%)	0.25±0.02
All	All	128/136 (94%)	15 (12%)	0 (0%)	0.25

The overall RNA backbone suiteness is 0.25.

All unique RNA backbone outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
1	A	9	C	8
1	A	11	C	4
1	A	8	U	2
1	A	7	U	1

There are no RNA pucker outliers to report.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 58% for the well-defined parts and 58% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	194
Number of shifts mapped to atoms	194
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 58%, i.e. 186 atoms were assigned a chemical shift out of a possible 320. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Sugar	145/187 (78%)	81/102 (79%)	64/85 (75%)	0/0 (—%)
Base	41/133 (31%)	22/82 (27%)	19/29 (66%)	0/22 (0%)
Overall	186/320 (58%)	103/184 (56%)	83/114 (73%)	0/22 (0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 58%, i.e. 186 atoms were assigned a chemical shift out of a possible 320. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Sugar	145/187 (78%)	81/102 (79%)	64/85 (75%)	0/0 (—%)
Base	41/133 (31%)	22/82 (27%)	19/29 (66%)	0/22 (0%)
Overall	186/320 (58%)	103/184 (56%)	83/114 (73%)	0/22 (0%)

7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

No *random coil index*(RCI) plot could be generated from the current chemical shift list. RCI is only applicable to proteins

8 NMR restraints analysis

8.1 Conformationally restricting restraints

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	237
Intra-residue ($ i-j =0$)	110
Sequential ($ i-j =1$)	94
Medium range ($ i-j >1$ and $ i-j <5$)	0
Long range ($ i-j \geq 5$)	33
Inter-chain	0
Hydrogen bond restraints	0
Disulfide bond restraints	0
Total dihedral-angle restraints	144
Number of unmapped restraints	0
Number of restraints per residue	22.4
Number of long range restraints per residue ¹	1.9

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	3.8	0.17
0.2-0.5 (Medium)	2.0	0.27
>0.5 (Large)	None	None

8.2.2 Average number of dihedral-angle violations per model

Dihedral-angle violations less than 1° are not included in the calculation. There are no dihedral-angle violations

9 Distance violation analysis [i](#)


9.1 Summary of distance violations [i](#)

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Restrains type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
Intra-residue (i-j =0)	110	46.4	8	7.3	3.4	2	1.8	0.8
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	110	46.4	8	7.3	3.4	2	1.8	0.8
Sequential (i-j =1)	94	39.7	1	1.1	0.4	1	1.1	0.4
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	94	39.7	1	1.1	0.4	1	1.1	0.4
Medium range (i-j >1 & i-j <5)	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Long range (i-j ≥5)	33	13.9	1	3.0	0.4	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	33	13.9	1	3.0	0.4	0	0.0	0.0
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	237	100.0	10	4.2	4.2	3	1.3	1.3
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	237	100.0	10	4.2	4.2	3	1.3	1.3

¹ percentage calculated with respect to the total number of distance restraints, ² percentage calculated with respect to the number of restraints in a particular restraint category, ³ violated in at least one model, ⁴ violated in all the models

9.1.1 Bar chart : Distribution of distance restraints and violations [i](#)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfid bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model [i](#)


The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Model ID	Number of violations						Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total				
1	5	1	0	1	0	7	0.16	0.23	0.05	0.15
2	5	1	0	0	0	6	0.18	0.27	0.05	0.16
3	5	1	0	1	0	7	0.16	0.23	0.05	0.15
4	4	1	0	1	0	6	0.16	0.23	0.05	0.15
5	3	1	0	1	0	5	0.19	0.26	0.05	0.17
6	4	1	0	1	0	6	0.17	0.23	0.05	0.16
7	3	1	0	0	0	4	0.2	0.24	0.04	0.2
8	3	1	0	1	0	5	0.18	0.23	0.04	0.16

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints,

⁵Inter-chain restraints, ⁶Standard deviation

9.2.1 Bar graph : Distance Violation statistics for each model [i](#)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right

9.3 Distance violation statistics for the ensemble [i](#)

Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 227(IR:102, SQ:93, MR:0, LR:32, IC:0) restraints are not violated in the ensemble.

Number of violated restraints						Fraction of the ensemble	
IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total	Count ⁶	%
1	0	0	0	0	1	1	12.5
4	0	0	0	0	4	2	25.0
0	0	0	0	0	0	3	37.5
0	0	0	0	0	0	4	50.0
0	0	0	0	0	0	5	62.5
0	0	0	1	0	1	6	75.0


Continued on next page...

Continued from previous page...

Number of violated restraints						Fraction of the ensemble	
IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total	Count ⁶	%
1	0	0	0	0	1	7	87.5
2	1	0	0	0	3	8	100.0

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints, ⁵Inter-chain restraints, ⁶ Number of models with violations


9.3.1 Bar graph : Distance violation statistics for the ensemble [i](#)



9.4 Most violated distance restraints in the ensemble [i](#)

9.4.1 Histogram : Distribution of mean distance violations [i](#)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble



9.4.2 Table: Most violated distance restraints [i](#)

The following table provides the mean and the standard deviation of the violation for each restraint sorted by number of violated models and the mean value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.


Key	Atom-1	Atom-2	Models ¹	Mean (Å)	SD ¹ (Å)	Median (Å)
(1,195)	1:A:8:U:H5	1:A:8:U:H4'	8	0.24	0.02	0.23
(1,172)	1:A:13:G:H1'	1:A:12:G:H3'	8	0.22	0.01	0.22
(1,152)	1:A:5:C:H1'	1:A:5:C:H5''	8	0.17	0.0	0.17
(1,166)	1:A:9:C:H1'	1:A:9:C:H5''	7	0.15	0.0	0.15
(2,31)	1:A:6:C:O2	1:A:12:G:H21	6	0.12	0.01	0.12
(1,185)	1:A:5:C:H5	1:A:5:C:H2'	2	0.12	0.0	0.12
(1,196)	1:A:9:C:H5	1:A:9:C:H2'	2	0.12	0.0	0.12
(1,204)	1:A:16:U:H5	1:A:16:U:H2'	2	0.12	0.0	0.12
(1,192)	1:A:7:U:H5	1:A:7:U:H2'	2	0.11	0.0	0.11

¹Number of violated models, ²Standard deviation

9.5 All violated distance restraints [i](#)

9.5.1 Histogram : Distribution of distance violations [i](#)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



9.5.2 Table : All distance violations [i](#)

The following table lists the absolute value of the violation for each restraint in the ensemble sorted by its value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,195)	1:A:8:U:H5	1:A:8:U:H4'	2	0.27
(1,195)	1:A:8:U:H5	1:A:8:U:H4'	5	0.26
(1,195)	1:A:8:U:H5	1:A:8:U:H4'	7	0.24
(1,195)	1:A:8:U:H5	1:A:8:U:H4'	1	0.23
(1,195)	1:A:8:U:H5	1:A:8:U:H4'	3	0.23
(1,195)	1:A:8:U:H5	1:A:8:U:H4'	4	0.23
(1,195)	1:A:8:U:H5	1:A:8:U:H4'	6	0.23
(1,195)	1:A:8:U:H5	1:A:8:U:H4'	8	0.23
(1,172)	1:A:13:G:H1'	1:A:12:G:H3'	4	0.23
(1,172)	1:A:13:G:H1'	1:A:12:G:H3'	5	0.23

Continued on next page...

Continued from previous page...

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,172)	1:A:13:G:H1'	1:A:12:G:H3'	6	0.23
(1,172)	1:A:13:G:H1'	1:A:12:G:H3'	1	0.22
(1,172)	1:A:13:G:H1'	1:A:12:G:H3'	2	0.22
(1,172)	1:A:13:G:H1'	1:A:12:G:H3'	3	0.22
(1,172)	1:A:13:G:H1'	1:A:12:G:H3'	7	0.22
(1,172)	1:A:13:G:H1'	1:A:12:G:H3'	8	0.21
(1,152)	1:A:5:C:H1'	1:A:5:C:H5''	1	0.17
(1,152)	1:A:5:C:H1'	1:A:5:C:H5''	4	0.17
(1,152)	1:A:5:C:H1'	1:A:5:C:H5''	5	0.17
(1,152)	1:A:5:C:H1'	1:A:5:C:H5''	6	0.17
(1,152)	1:A:5:C:H1'	1:A:5:C:H5''	7	0.17
(1,166)	1:A:9:C:H1'	1:A:9:C:H5''	2	0.16
(1,166)	1:A:9:C:H1'	1:A:9:C:H5''	5	0.16
(1,166)	1:A:9:C:H1'	1:A:9:C:H5''	8	0.16
(1,152)	1:A:5:C:H1'	1:A:5:C:H5''	2	0.16
(1,152)	1:A:5:C:H1'	1:A:5:C:H5''	3	0.16
(1,152)	1:A:5:C:H1'	1:A:5:C:H5''	8	0.16
(1,166)	1:A:9:C:H1'	1:A:9:C:H5''	1	0.15
(1,166)	1:A:9:C:H1'	1:A:9:C:H5''	3	0.15
(1,166)	1:A:9:C:H1'	1:A:9:C:H5''	6	0.15
(1,166)	1:A:9:C:H1'	1:A:9:C:H5''	7	0.15
(2,31)	1:A:6:C:O2	1:A:12:G:H21	8	0.13
(2,31)	1:A:6:C:O2	1:A:12:G:H21	3	0.12
(2,31)	1:A:6:C:O2	1:A:12:G:H21	4	0.12
(1,204)	1:A:16:U:H5	1:A:16:U:H2'	1	0.12
(1,196)	1:A:9:C:H5	1:A:9:C:H2'	2	0.12
(1,185)	1:A:5:C:H5	1:A:5:C:H2'	2	0.12
(1,185)	1:A:5:C:H5	1:A:5:C:H2'	3	0.12
(2,31)	1:A:6:C:O2	1:A:12:G:H21	1	0.11
(2,31)	1:A:6:C:O2	1:A:12:G:H21	5	0.11
(2,31)	1:A:6:C:O2	1:A:12:G:H21	6	0.11
(1,204)	1:A:16:U:H5	1:A:16:U:H2'	3	0.11
(1,196)	1:A:9:C:H5	1:A:9:C:H2'	6	0.11
(1,193)	1:A:8:U:H5	1:A:8:U:H2'	4	0.11
(1,192)	1:A:7:U:H5	1:A:7:U:H2'	1	0.11
(1,192)	1:A:7:U:H5	1:A:7:U:H2'	4	0.11

10 Dihedral-angle violation analysis [i](#)


10.1 Summary of dihedral-angle violations [i](#)

The following table provides the summary of dihedral-angle violations in different dihedral-angle types. Violations less than 1° are not included in the calculation.

Angle type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
ZETA	16	11.1	0	0.0	0.0	0	0.0	0.0
ALPHA	16	11.1	0	0.0	0.0	0	0.0	0.0
DELTA	17	11.8	0	0.0	0.0	0	0.0	0.0
GAMMA	12	8.3	0	0.0	0.0	0	0.0	0.0
.	51	35.4	0	0.0	0.0	0	0.0	0.0
BETA	16	11.1	0	0.0	0.0	0	0.0	0.0
EPSILON	16	11.1	0	0.0	0.0	0	0.0	0.0
Total	144	100.0	0	0.0	0.0	0	0.0	0.0

¹ percentage calculated with respect to total number of dihedral-angle restraints, ² percentage calculated with respect to number of restraints in a particular dihedral-angle type, ³ violated in at least one model, ⁴ violated in all the models

10.1.1 Bar chart : Distribution of dihedral-angles and violations [i](#)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories

10.2 Dihedral-angle violation statistics for each model [i](#)

No violations found

10.3 Dihedral-angle violation statistics for the ensemble [i](#)

No violations found

10.4 Most violated dihedral-angle restraints in the ensemble [i](#)

No violations found

10.5 All violated dihedral-angle restraints [i](#)

No violations found