



wwPDB NMR Structure Validation Summary Report ⓘ

Apr 21, 2024 – 09:17 AM EDT

PDB ID : 2LBS
BMRB ID : 17574
Title : Solution structure of double-stranded RNA binding domain of *S. cerevisiae* RNase III (Rnt1p) in complex with AAGU tetraloop hairpin
Authors : Wang, Z.; Hartman, E.; Roy, K.; Chanfreau, G.; Feigon, J.
Deposited on : 2011-04-06

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 51%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428
RNA backbone	4643	676

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	32	 41% 47% 12%
2	B	90	 7% 56% 31% 7%

2 Ensemble composition and analysis i

This entry contains 16 models. Model 8 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	B:365-B:448 (84)	1.01	8

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 15, 16
2	3, 14
Single-model clusters	2

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2434 atoms, of which 1066 are hydrogens and 0 are deuteriums.

- Molecule 1 is a RNA chain called RNA (32-MER).

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	P	
1	A	32	1025	305	345	121	223	31	0

- Molecule 2 is a protein called Ribonuclease 3.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
2	B	90	1409	429	721	129	127	3	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	364	GLY	-	expression tag	UNP Q02555
B	365	SER	-	expression tag	UNP Q02555

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: RNA (32-MER)

Chain A: 



- Molecule 2: Ribonuclease 3

Chain B: 



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 8. Colouring as in section 4.1 above.


- Molecule 1: RNA (32-MER)

Chain A: 



- Molecule 2: Ribonuclease 3

Chain B: 



E427	I425	R426	A427	A428	E429	M430	A431	L432	R433	D434	K435	K436	M437	L438	D439	F440	Y441	A442	K443	Q444	R445	A446	A447	I448	F449	R450	S451	E452	S453
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics*.

Of the 100 calculated structures, 16 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	structure solution	2.24
X-PLOR NIH	refinement	2.24

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	952
Number of shifts mapped to atoms	952
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	51%

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.13±0.01	0±0/760 (0.0± 0.0%)	1.98±0.01	42±1/1183 (3.5± 0.1%)
2	B	0.24±0.01	0±0/652 (0.0± 0.0%)	0.41±0.01	0±0/879 (0.0± 0.0%)
All	All	0.84	0/22592 (0.0%)	1.53	664/32992 (2.0%)

There are no bond-length outliers.

5 of 45 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	25	G	N7-C8-N9	9.72	117.96	113.10	8	16
1	A	2	G	N7-C8-N9	9.66	117.93	113.10	4	16
1	A	11	G	N7-C8-N9	9.40	117.80	113.10	16	16
1	A	3	G	N7-C8-N9	9.38	117.79	113.10	9	16
1	A	26	G	N7-C8-N9	9.33	117.77	113.10	6	16

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	680	345	345	86±11
2	B	644	680	679	131±14
All	All	21184	16400	16384	3393

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 90.

5 of 1104 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
2:B:412:LEU:HD22	2:B:437:MET:HE2	1.04	1.28	11	7
1:A:15:A:N3	1:A:16:A:H2'	1.00	1.69	16	1
2:B:370:ALA:HB1	2:B:428:ALA:HB1	0.99	1.28	13	8
2:B:390:VAL:O	2:B:401:ILE:HG22	0.97	1.57	9	13
2:B:404:CYS:C	2:B:411:VAL:HG23	0.97	1.80	8	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	84/90 (93%)	66±2 (79±3%)	14±2 (16±3%)	4±1 (5±2%)	4	27
All	All	1344/1440 (93%)	1062 (79%)	219 (16%)	63 (5%)	4	27

5 of 14 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	B	365	SER	8
2	B	399	ASN	8
2	B	407	GLY	7
2	B	420	ILE	6
2	B	398	PRO	6

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	B	67/72 (93%)	36±3 (54±5%)	31±3 (46±5%)	0 2
All	All	1072/1152 (93%)	577 (54%)	495 (46%)	0 2

5 of 61 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	B	402	VAL	16
2	B	444	GLN	16
2	B	371	LYS	15
2	B	373	GLN	15
2	B	377	LEU	15

6.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers	Suiteness
1	A	31/32 (97%)	6±1 (21±3%)	2±1 (7±2%)	0.14±0.02
All	All	496/512 (97%)	104 (21%)	35 (7%)	0.14

The overall RNA backbone suiteness is 0.14.

5 of 13 unique RNA backbone outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
1	A	17	G	16
1	A	18	U	16
1	A	19	G	16
1	A	22	C	15
1	A	21	A	11

5 of 9 unique RNA pucker outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
1	A	18	U	16
1	A	17	G	7
1	A	21	A	4
1	A	24	U	2
1	A	16	A	2

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 51% for the well-defined parts and 52% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	952
Number of shifts mapped to atoms	952
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	2

7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	88	-0.34 ± 0.14	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	81	0.20 ± 0.14	None needed (< 0.5 ppm)
$^{13}\text{C}'$	79	-0.21 ± 0.14	None needed (< 0.5 ppm)
^{15}N	82	-0.16 ± 0.31	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 51%, i.e. 896 atoms were assigned a chemical shift out of a possible 1762. 0 out of 15 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	403/423 (95%)	167/173 (97%)	158/168 (94%)	78/82 (95%)
Sidechain	476/686 (69%)	322/448 (72%)	149/204 (73%)	5/34 (15%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	17/54 (31%)	17/25 (68%)	0/27 (0%)	0/2 (0%)
Sugar	0/352 (0%)	0/192 (0%)	0/160 (0%)	0/0 (—%)
Base	0/247 (0%)	0/151 (0%)	0/55 (0%)	0/41 (0%)
Overall	896/1762 (51%)	506/989 (51%)	307/614 (50%)	83/159 (52%)

7.1.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	B	448	ILE	CG1	16.40	19.24 – 36.26	-6.7
1	B	420	ILE	CG1	18.29	19.24 – 36.26	-5.6

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain B:

