



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 23, 2024 – 02:59 AM EDT

PDB ID : 6LBR  
Title : Crystal structure of yeast Cdc13 and ssDNA  
Authors : Ge, Y.; Wu, Z.; Wu, J.; Lei, M.  
Deposited on : 2019-11-14  
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.20.1  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	551	 4% 77% 15% • 6%
1	B	551	 2% 32% 5% • 62%
2	C	25	 12% 84% • 12%
2	D	25	 16% 76% 16% 8%

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7028 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called KLLA0F20922p.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	519	4190	2702	687	788	13	0	0	0
1	B	208	1671	1078	280	307	6	0	0	0

- Molecule 2 is a DNA chain called Telomere single-strand DNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	C	22	460	219	84	135	22	0	0	0
2	D	23	480	229	86	142	23	0	0	0

- Molecule 3 is water.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	148	Total 148	O 148	0	0
3	B	35	Total 35	O 35	0	0
3	C	24	Total 24	O 24	0	0
3	D	20	Total 20	O 20	0	0





GLU  
GLU  
LEU  
THR  
ALA  
CYS  
ILE  
GLY  
SER  
ILE  
VAL  
ASP  
PHE  
THR  
LEU  
ILE  
LEU  
PRO  
ARG  
VAL  
LEU  
ARG  
VAL  
ASN  
GLU  
TYR  
LEU  
TYR  
ILE  
LYS  
ILE  
ILE  
TRP  
CYS  
PRO  
ILE  
TYR  
ALA  
THR  
LEU  
GLU  
SER  
LEU  
ILE  
HIS  
SER  
SER  
ARG  
GLU  
TYR  
ASP  
ASN  
ASP  
THR  
LEU  
LYS  
TYR  
GLU  
ASN  
LEU  
SER

ASP  
ILE  
ASP

- Molecule 2: Telomere single-strand DNA



- Molecule 2: Telomere single-strand DNA



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	75.17Å 106.95Å 173.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.84 – 2.50 45.84 – 2.50	Depositor EDS
% Data completeness (in resolution range)	93.8 (45.84-2.50) 93.7 (45.84-2.50)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.65 (at 2.51Å)	Xtrriage
Refinement program	PHENIX 1.15_3459	Depositor
R, $R_{free}$	0.202 , 0.252 0.203 , 0.252	Depositor DCC
$R_{free}$ test set	2219 reflections (4.80%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	37.5	Xtrriage
Anisotropy	0.136	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 53.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	7028	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.19% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.41	0/4278	0.69	8/5802 (0.1%)
1	B	0.41	0/1706	0.62	1/2310 (0.0%)
2	C	0.82	0/516	1.02	0/797
2	D	0.87	0/538	1.05	0/831
All	All	0.50	0/7038	0.74	9/9740 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	702	ARG	N-CA-C	14.85	151.10	111.00
1	A	703	MET	CB-CA-C	12.26	134.92	110.40
1	A	704	SER	N-CA-CB	-11.40	93.40	110.50
1	A	702	ARG	CB-CA-C	-8.14	94.11	110.40
1	A	740	GLU	CB-CA-C	-7.21	95.98	110.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	617	CYS	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4190	0	4252	53	0
1	B	1671	0	1719	23	0
2	C	460	0	251	1	0
2	D	480	0	263	6	0
3	A	148	0	0	2	0
3	B	35	0	0	0	0
3	C	24	0	0	0	0
3	D	20	0	0	0	0
All	All	7028	0	6485	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 76 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:656:THR:HG22	1:B:658:THR:H	1.31	0.94
1:A:759:THR:HG21	1:A:779:LYS:HA	1.64	0.78
1:A:562:ASP:HB3	1:A:564:THR:HB	1.68	0.76
1:A:656:THR:HG22	1:A:658:THR:H	1.51	0.74
1:A:860:ILE:HG23	1:A:878:ILE:HD11	1.70	0.72

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	515/551 (94%)	497 (96%)	18 (4%)	0	100	100
1	B	204/551 (37%)	198 (97%)	6 (3%)	0	100	100
All	All	719/1102 (65%)	695 (97%)	24 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	477/507 (94%)	449 (94%)	28 (6%)	19	37
1	B	192/507 (38%)	185 (96%)	7 (4%)	35	61
All	All	669/1014 (66%)	634 (95%)	35 (5%)	23	44

5 of 35 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	509	LEU
1	B	564	THR
1	B	612	MET
1	A	690	LEU
1	A	678	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	810	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	519/551 (94%)	-0.11	23 (4%) 34 37	17, 34, 81, 143	0
1	B	208/551 (37%)	-0.04	11 (5%) 26 28	25, 40, 73, 107	0
2	C	22/25 (88%)	0.01	3 (13%) 3 2	30, 47, 100, 112	0
2	D	23/25 (92%)	0.46	4 (17%) 1 1	32, 60, 89, 120	0
All	All	772/1152 (67%)	-0.07	41 (5%) 26 28	17, 37, 80, 143	0

The worst 5 of 41 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	703	MET	10.1
1	A	704	SER	7.5
1	A	701	ASN	6.0
1	A	702	ARG	6.0
1	B	508	ALA	4.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.