

Full wwPDB X-ray Structure Validation Report (i)

Sep 19, 2020 - 09:16 AM BST

PDB ID	:	6LBS
Title	:	Crystal structure of yeast Stn1
Authors	:	Ge, Y.; Wu, Z.; Wu, J.; Lei, M.
Deposited on	:	2019-11-14
Resolution	:	2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as 541 be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.14.6
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
$\operatorname{CCP4}$:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.14.6

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	130704	$3163 \ (2.60-2.60)$
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455(2.60-2.60)
Sidechain outliers	138945	3455(2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain			
1	А	166	2% 71 %	17%		11%
	_	100	2%	1170		1170
1	В	166	64%	20%	•	12%
1	С	166	65%	20%	·	13%
1	D	166	69%	18%	•	10%
1	Е	166	^{2%} 70%	17%	·	11%
1	F	166	3% 69%	16%	• •	11%



2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 7463 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues			Atom	IS			ZeroOcc	AltConf	Trace
1	Λ	148	Total	С	Ν	Ο	\mathbf{S}	Se	0	0	0
L T	Л	140	1227	806	199	213	4	5	0	0	0
1	В	146	Total	С	Ν	Ο	S	Se	0	0	0
1	D	140	1217	799	198	211	4	5	0	0	0
1	C	144	Total	С	Ν	Ο	\mathbf{S}	Se	0	0	0
L L	U	144	1191	782	194	206	4	5			
1	п	150	Total	С	Ν	Ο	\mathbf{S}	Se	0	0	0
1	D	100	1245	818	203	215	4	5	0	0	U
1	F	148	Total	С	Ν	Ο	\mathbf{S}	Se	0	0	0
	Ľ	140	1228	806	200	213	4	5	0	0	0
1	F	148	Total	C	Ν	Ō	\mathbf{S}	Se	0	0	0
	Ľ	140	1229	807	200	213	4	5	U	U	U

• Molecule 1 is a protein called KLLA0C11825p.

• Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	13	Total O 13 13	0	0
2	В	13	Total O 13 13	0	0
2	С	10	Total O 10 10	0	0
2	D	29	Total O 29 29	0	0
2	Ε	25	$\begin{array}{cc} \text{Total} & \text{O} \\ 25 & 25 \end{array}$	0	0
2	F	36	$\begin{array}{cc} {\rm Total} & {\rm O} \\ 36 & 36 \end{array}$	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: KLLA0C11825p



• Molecule 1: KLLA0C11825p





4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	158.43Å 94.75 Å 97.50 Å	Depositor
a, b, c, α , β , γ	90.00° 96.26° 90.00°	Depositor
$\mathbf{Bosolution} \left(\overset{\wedge}{\mathbf{A}} \right)$	37.95 - 2.60	Depositor
	40.04 - 2.59	EDS
% Data completeness	97.2 (37.95-2.60)	Depositor
(in resolution range)	96.7(40.04-2.59)	EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.64 (at 2.58 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.15_3459	Depositor
B B.	0.183 , 0.235	Depositor
n, n_{free}	0.182 , 0.233	DCC
R_{free} test set	2114 reflections $(4.88%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	38.4	Xtriage
Anisotropy	0.062	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.35 , 49.3	EDS
L-test for twinning ²	$ < L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7463	wwPDB-VP
Average B, all atoms $(Å^2)$	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 7.10% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bo	nd lengths	Bond angles		
	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.40	0/1249	0.58	0/1678	
1	В	0.42	0/1238	0.63	0/1662	
1	С	0.42	0/1211	0.68	1/1626~(0.1%)	
1	D	0.47	0/1267	0.59	0/1700	
1	Е	0.42	0/1250	0.64	0/1678	
1	F	0.48	1/1250~(0.1%)	0.60	0/1678	
All	All	0.44	1/7465~(0.0%)	0.62	1/10022~(0.0%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	$\operatorname{Ideal}(\operatorname{\AA})$
1	F	356	CYS	CB-SG	-6.29	1.71	1.82

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	С	420	THR	N-CA-C	5.73	126.47	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	432	SER	Peptide



5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1227	0	1277	18	0
1	В	1217	0	1268	24	0
1	С	1191	0	1240	20	0
1	D	1245	0	1303	25	0
1	Е	1228	0	1279	17	0
1	F	1229	0	1282	18	0
2	А	13	0	0	1	0
2	В	13	0	0	1	0
2	С	10	0	0	2	0
2	D	29	0	0	0	0
2	Е	25	0	0	0	0
2	F	36	0	0	0	0
All	All	7463	0	7649	116	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (116) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:292:MSE:HE1	1:F:354:HIS:HA	1.56	0.88
1:D:292:MSE:HE1	1:D:354:HIS:HA	1.65	0.79
1:B:292:MSE:HE1	1:B:354:HIS:HA	1.65	0.78
1:C:360:ILE:HD12	1:C:399:MSE:HE2	1.69	0.74
1:B:287:LEU:HD22	1:B:350:MSE:HE1	1.69	0.73
1:E:287:LEU:HD22	1:E:350:MSE:HE1	1.77	0.66
1:E:344:ASP:HB3	1:E:346:PHE:H	1.61	0.66
1:D:421:SER:HB2	1:D:424:ARG:H	1.64	0.63
1:A:393:LYS:NZ	1:A:422:GLU:OE1	2.34	0.61
1:D:360:ILE:HD12	1:D:399:MSE:HE2	1.83	0.60
1:B:382:ILE:HD12	1:B:382:ILE:H	1.66	0.60
1:D:432:SER:O	1:D:433:LYS:HB2	2.03	0.59
1:D:287:LEU:HB3	1:D:350:MSE:HE1	1.85	0.59
1:D:300:TYR:HD1	1:D:304:LEU:HD22	1.68	0.58
1:F:299:ILE:HG13	1:F:346:PHE:O	2.04	0.57



		Interatomic	Clash
Atom-1	Atom-2	distance $(Å)$	overlap (Å)
1:F:291:SER:OG	1:F:292:MSE:HE2	2.06	0.56
1:C:363:ARG:HH21	1:C:377:ILE:HG12	1.70	0.56
1:F:287:LEU:HB3	1:F:350:MSE:HE1	1.89	0.55
1:C:376:THR:HG23	1:C:426:LYS:HG2	1.88	0.55
1:B:393:LYS:NZ	1:B:422:GLU:O	2.27	0.55
1:F:356:CYS:HB3	1:F:399:MSE:HE1	1.89	0.55
1:D:290:LYS:HE3	1:D:301:GLU:OE2	2.06	0.54
1:A:304:LEU:HD12	1:B:300:TYR:HB3	1.89	0.54
1:C:287:LEU:HD22	1:C:350:MSE:CE	2.38	0.54
1:A:299:ILE:HD11	1:A:344:ASP:O	2.08	0.53
1:A:304:LEU:HD11	1:B:299:ILE:HG22	1.91	0.53
1:D:413:ILE:HG21	1:D:432:SER:HB3	1.91	0.52
1:B:370:VAL:HG12	1:B:372:ILE:HG12	1.91	0.52
1:B:416:TRP:HZ2	1:C:430:THR:HG1	1.57	0.52
1:A:298:LYS:HB2	1:A:301:GLU:HG3	1.91	0.52
1:A:367:LEU:HD23	1:A:372:ILE:HD13	1.92	0.52
1:C:303:LEU:HD23	1:D:304:LEU:HD21	1.92	0.51
1:E:344:ASP:HB3	1:E:347:THR:H	1.76	0.51
1:E:342:THR:OG1	1:E:349:ASN:HB3	2.11	0.51
1:A:421:SER:HB3	1:A:424:ARG:HB2	1.93	0.50
1:D:288:ILE:HD11	1:D:399:MSE:HG2	1.93	0.50
1:C:338:MSE:SE	1:C:392:SER:HB3	2.61	0.50
1:B:365:GLN:NE2	1:B:409:ILE:HB	2.26	0.50
1:C:292:MSE:HE1	1:C:354:HIS:HA	1.93	0.50
1:C:367:LEU:HD21	1:C:375:ILE:HD13	1.94	0.49
1:E:421:SER:HB2	1:E:424:ARG:H	1.77	0.49
1:B:416:TRP:CZ3	1:B:418:ASP:HB2	2.47	0.49
1:D:287:LEU:HD22	1:D:350:MSE:HE1	1.94	0.48
1:D:413:ILE:HD13	1:D:432:SER:HB3	1.95	0.48
1:A:292:MSE:HE1	1:A:354:HIS:HA	1.94	0.48
1:B:391:LEU:HD22	1:B:395:LEU:HD23	1.95	0.48
1:E:340:ILE:HD12	1:E:391:LEU:HD21	1.95	0.48
1:F:415:TRP:HA	1:F:428:PHE:O	2.13	0.48
1:C:417:VAL:HA	1:C:427:VAL:HG12	1.96	0.48
1:A:420:THR:OG1	1:A:421:SER:N	2.46	0.48
1:F:375:ILE:HG13	1:F:376:THR:N	2.29	0.48
1:F:371:GLN:HA	1:F:431:TYR:CZ	2.49	0.47
1:D:292:MSE:HE3	1:D:357:TYR:HB3	1.96	0.47
1:A:307:ASN:O	1:A:308:LEU:HD23	2.15	0.47
1:E:382:ILE:H	1:E:382:ILE:HD12	1.78	0.47
1:F:275:SER:HB3	1:F:278:GLN:OE1	2.15	0.47



		Interatomic	Clash
Atom-1	Atom-2	distance $(Å)$	overlan (Å)
1.D.291.SEB.OG	$1 \cdot D \cdot 292 \cdot MSE \cdot HE2$	2.15	0.47
1:F:402:HIS:O	1:F:406:THR:HB	$\frac{2.10}{2.16}$	0.46
1:E:416:TBP:CZ3	1:E:418:ASP:HB2	2.51	0.46
1:C:344:ASP:HB2	1:C:347:THR:OG1	2.01	0.46
1:B:322:VAL:N	2:B:501:HOH:O	2.49	0.46
1:B:336:ALA:HA	1:B:341:ILE:O	2.16	0.46
1:A:299:ILE:HG21	1:A:299:ILE:HD13	1.59	0.46
1:B:350:MSE:HB3	1:B:350:MSE:HE2	1.81	0.46
1:C:304:LEU:HD12	1:C:304:LEU:HA	1.74	0.45
1:A:378:LYS:HE3	1:A:424:ARG:HH21	1.81	0.45
1:C:412:LEU:CD1	1:C:429:PHE:HB3	2.46	0.45
1:E:307:ASN:OD1	1:E:322:VAL:HB	2.15	0.45
1:E:375:ILE:HG13	1:E:376:THR:N	2.32	0.45
1:E:421:SER:HB3	1:E:423:GLU:H	1.82	0.45
1:C:412:LEU:HD11	1:C:429:PHE:HB3	1.98	0.45
1:B:360:ILE:HD12	1:B:399:MSE:HE2	2.00	0.44
1:F:358:ARG:HH21	1:F:361:ILE:HG21	1.82	0.44
1:C:363:ARG:NH2	1:C:377:ILE:HG12	2.32	0.44
1:D:421:SER:HB2	1:D:423:GLU:H	1.83	0.44
1:F:420:THR:HG21	1:F:426:LYS:HE3	2.00	0.44
1:A:376:THR:OG1	1:A:426:LYS:HG2	2.18	0.44
1:D:300:TYR:CD1	1:D:304:LEU:HD22	2.51	0.44
1:B:372:ILE:HD13	1:B:372:ILE:HA	1.72	0.43
1:D:371:GLN:HA	1:D:431:TYR:CZ	2.53	0.43
1:F:361:ILE:O	1:F:365:GLN:HB2	2.17	0.43
1:E:287:LEU:HB3	1:E:350:MSE:HE1	2.00	0.43
1:C:364:PHE:HE2	1:C:427:VAL:HG21	1.83	0.43
1:D:292:MSE:CE	1:D:354:HIS:HA	2.42	0.43
1:F:350:MSE:HB3	1:F:350:MSE:HE2	1.93	0.43
1:E:371:GLN:HA	1:E:431:TYR:CZ	2.54	0.43
1:E:304:LEU:HB3	1:F:325:LEU:CD1	2.49	0.43
1:B:298:LYS:HB2	1:B:301:GLU:HG3	2.00	0.42
1:B:338:MSE:CE	1:B:395:LEU:HD22	2.49	0.42
1:F:279:MSE:HE1	1:F:322:VAL:HG12	2.00	0.42
1:F:350:MSE:HE2	1:F:353:LEU:HD23	2.00	0.42
1:D:421:SER:HB2	1:D:424:ARG:N	2.34	0.42
1:B:303:LEU:HD21	1:B:325:LEU:HD23	2.00	0.42
1:C:430:THR:C	2:C:505:HOH:O	2.57	0.42
1:D:287:LEU:CB	1:D:350:MSE:HE1	2.49	0.42
1:D:350:MSE:HE2	1:D:350:MSE:HB3	1.72	0.42
1:C:368:ILE:C	1:C:370:VAL:H	2.21	0.42



Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	$distance ({ m \AA})$	overlap (Å)
1:A:279:MSE:HE1	1:A:322:VAL:HG12	2.01	0.42
1:B:357:TYR:CE2	1:B:361:ILE:HD11	2.55	0.42
1:F:299:ILE:H	1:F:299:ILE:HG13	1.46	0.41
1:A:299:ILE:HG13	1:A:346:PHE:C	2.40	0.41
1:B:376:THR:OG1	1:B:426:LYS:HE2	2.19	0.41
1:C:372:ILE:HG23	1:C:373:PRO:HD2	2.03	0.41
1:E:370:VAL:HG12	1:E:372:ILE:HG12	2.02	0.41
1:D:279:MSE:O	1:D:283:ILE:HG13	2.20	0.41
1:E:332:LEU:HD23	1:E:332:LEU:HA	1.84	0.41
1:D:404:LEU:HD23	1:D:404:LEU:HA	1.78	0.41
1:E:299:ILE:HD11	1:E:345:SER:C	2.39	0.41
1:D:304:LEU:HD12	1:D:304:LEU:HA	1.82	0.41
1:B:362:THR:O	1:B:366:SER:HB3	2.20	0.41
1:A:344:ASP:HB3	2:A:512:HOH:O	2.21	0.40
1:B:397:LEU:HD23	1:B:397:LEU:HA	1.81	0.40
1:A:299:ILE:HA	1:A:348:ILE:HG13	2.04	0.40
1:A:308:LEU:HD11	1:B:303:LEU:HD21	2.02	0.40
1:D:335:LEU:HA	1:D:338:MSE:HE3	2.04	0.40
1:C:371:GLN:OE1	2:C:501:HOH:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	\mathbf{ntiles}
1	А	144/166~(87%)	138~(96%)	6~(4%)	0	100	100
1	В	142/166~(86%)	135~(95%)	7~(5%)	0	100	100
1	С	140/166~(84%)	128~(91%)	12 (9%)	0	100	100
1	D	146/166~(88%)	141 (97%)	5(3%)	0	100	100
1	Ε	144/166~(87%)	140 (97%)	4(3%)	0	100	100



001000							
Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	F	144/166~(87%)	138~(96%)	6 (4%)	0	100 100	
All	All	860/996~(86%)	820 (95%)	40~(5%)	0	100 100	

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	\mathbf{ntiles}
1	А	142/152~(93%)	134~(94%)	8 (6%)	21	42
1	В	141/152~(93%)	129~(92%)	12 (8%)	10	21
1	С	137/152~(90%)	128~(93%)	9~(7%)	16	33
1	D	144/152~(95%)	131 (91%)	13 (9%)	9	18
1	Е	142/152~(93%)	135~(95%)	7(5%)	25	48
1	F	142/152~(93%)	128 (90%)	14 (10%)	8	15
All	All	848/912~(93%)	785~(93%)	63 (7%)	13	28

All (63) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	281	ARG
1	А	284	LEU
1	А	294	SER
1	А	304	LEU
1	А	358	ARG
1	А	366	SER
1	А	371	GLN
1	А	405	ASN
1	В	275	SER
1	В	279	MSE
1	В	284	LEU
1	В	292	MSE
1	В	298	LYS



Mol	Chain	Res	Type
1	В	326	LYS
1	В	342	THR
1	В	350	MSE
1	В	358	ARG
1	В	366	SER
1	В	370	VAL
1	В	422	GLU
1	С	284	LEU
1	С	358	ARG
1	С	365	GLN
1	С	375	ILE
1	С	376	THR
1	С	394	LYS
1	С	408	HIS
1	С	416	TRP
1	С	420	THR
1	D	282	GLU
1	D	284	LEU
1	D	301	GLU
1	D	304	LEU
1	D	326	LYS
1	D	340	ILE
1	D	346	PHE
1	D	350	MSE
1	D	358	ARG
1	D	363	ARG
1	D	370	VAL
1	D	387	LYS
1	D	424	ARG
1	Е	275	SER
1	Е	284	LEU
1	E	299	ILE
1	Е	304	LEU
1	E	350	MSE
1	Е	358	ARG
1	E	387	LYS
1	F	274	LEU
1	F	275	SER
1	F	284	LEU
1	F	299	ILE
1	F	324	HIS
1	F	350	MSE



Continued from previous page						
Mol	Chain	\mathbf{Res}	Type			
1	F	356	CYS			
1	F	358	ARG			
1	F	365	GLN			
1	F	370	VAL			
1	F	375	ILE			
1	F	394	LYS			
1	F	406	THR			
1	F	432	SER			

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$<$ RSRZ $>$	#RSRZ $>$ 2	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q < 0.9
1	А	143/166~(86%)	-0.33	4 (2%) 53 46	24,43,83,114	0
1	В	141/166~(84%)	-0.33	3 (2%) 63 58	25, 41, 82, 123	0
1	С	139/166~(83%)	0.14	11 (7%) 12 9	26, 56, 111, 136	0
1	D	145/166~(87%)	-0.55	0 100 100	20,33,66,105	0
1	Ε	143/166~(86%)	-0.41	4 (2%) 53 46	23,39,70,109	0
1	F	143/166~(86%)	-0.51	5 (3%) 44 36	17,33,78,130	0
All	All	854/996~(85%)	-0.33	27 (3%) 47 40	17, 40, 89, 136	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	С	307	ASN	6.0
1	С	323	LEU	5.1
1	С	420	THR	4.9
1	В	274	LEU	4.7
1	С	276	PRO	3.9
1	А	408	HIS	3.7
1	F	309	GLY	3.6
1	А	409	ILE	3.5
1	Е	409	ILE	3.5
1	С	274	LEU	3.2
1	Е	309	GLY	3.2
1	С	409	ILE	3.0
1	С	308	LEU	2.8
1	С	376	THR	2.7
1	С	309	GLY	2.7
1	F	308	LEU	2.6
1	A	413	ILE	2.6
1	С	408	HIS	2.5
1	F	408	HIS	2.4



Mol	Chain	Res	Type	RSRZ
1	F	409	ILE	2.2
1	А	432	SER	2.2
1	Е	408	HIS	2.2
1	В	410	GLY	2.1
1	Е	406	THR	2.1
1	В	408	HIS	2.1
1	С	277	LYS	2.1
1	F	304	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.

