



# wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 14, 2023 – 11:15 AM EDT

PDB ID : 1LE5  
Title : Crystal structure of a NF-kB heterodimer bound to an IFN $\beta$ -kB  
Authors : Berkowitz, B.; Huang, D.B.; Chen-Park, F.E.; Sigler, P.B.; Ghosh, G.  
Deposited on : 2002-04-09  
Resolution : 2.75 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1277 (2.78-2.74)
Ramachandran outliers	138981	1257 (2.78-2.74)
Sidechain outliers	138945	1257 (2.78-2.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	C	12	17% 83%
1	G	12	50% 33% 17%
2	D	12	25% 75%
2	H	12	25% 8% 67%
3	A	274	31% 57% 12% .
3	E	274	31% 57% 10% .
4	B	313	28% 61% 10%
4	F	313	34% 52% 13% .

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 10439 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(\*TP\*GP\*GP\*GP\*AP\*AP\*AP\*TP\*TP\*CP\*CP\*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	C	12	Total 244	C 118	N 44	O 71	P 11	0	0	0
1	G	12	Total 244	C 118	N 44	O 71	P 11	0	0	0

- Molecule 2 is a DNA chain called 5'-D(\*AP\*AP\*GP\*GP\*AP\*AP\*TP\*TP\*TP\*CP\*CP\*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	D	12	Total 242	C 117	N 45	O 69	P 11	0	0	0
2	H	12	Total 242	C 117	N 45	O 69	P 11	0	0	0

- Molecule 3 is a protein called Nuclear factor NF-kappa-B p65 subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	A	274	Total 2184	C 1361	N 402	O 409	S 12	0	0	0
3	E	274	Total 2184	C 1361	N 402	O 409	S 12	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	18	MET	-	cloning artifact	UNP Q04207
A	19	ALA	-	cloning artifact	UNP Q04207
E	18	MET	-	cloning artifact	UNP Q04207
E	19	ALA	-	cloning artifact	UNP Q04207

- Molecule 4 is a protein called Nuclear factor NF-kappa-B p50 subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	B	313	Total	C	N	O	S	0	0	0
			2462	1559	429	461	13			
4	F	313	Total	C	N	O	S	0	0	0
			2462	1559	429	461	13			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	38	MET	-	initiating methionine	UNP P25799
F	38	MET	-	initiating methionine	UNP P25799

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	4	Total	O	0	0
			4	4		
5	D	9	Total	O	0	0
			9	9		
5	G	6	Total	O	0	0
			6	6		
5	H	8	Total	O	0	0
			8	8		
5	A	33	Total	O	0	0
			33	33		
5	B	32	Total	O	0	0
			32	32		
5	E	32	Total	O	0	0
			32	32		
5	F	51	Total	O	0	0
			51	51		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: 5'-D(\*TP\*GP\*GP\*GP\*AP\*AP\*AP\*TP\*TP\*CP\*CP\*T)-3'

Chain C: 




- Molecule 1: 5'-D(\*TP\*GP\*GP\*GP\*AP\*AP\*AP\*TP\*TP\*CP\*CP\*T)-3'

Chain G: 




- Molecule 2: 5'-D(\*AP\*AP\*GP\*GP\*AP\*AP\*TP\*TP\*TP\*CP\*CP\*C)-3'

Chain D: 





- Molecule 2: 5'-D(\*AP\*AP\*GP\*GP\*AP\*AP\*TP\*TP\*TP\*CP\*CP\*C)-3'


Chain H: 








- Molecule 3: Nuclear factor NF-kappa-B p65 subunit

Chain A: 









• Molecule 3: Nuclear factor NF-kappa-B p65 subunit



• Molecule 4: Nuclear factor NF-kappa-B p50 subunit



• Molecule 4: Nuclear factor NF-kappa-B p50 subunit



M38	H107	S171	D235	N320
Y41	A108	D172	A236	I321
L42	H109	L173	Y238	T322
Q43	S110	A174	D239	K323
I44	L111	Y175	S240	V327
L45	V112	L176	M244	F328
E46	G113	Q177	A285	V329
Q47	K114	A178	S246	
P48	H115	E179	M247	R332
K49	C116	G182	L248	R333
Q50	E117	D183	K249	K334
	D118	R184	I250	S335
F55	C121	Q185	D254	D336
R56	T122	L186	R255	L337
Y57	V123	T187	D254	E338
V58	V124	D188	R255	T339
	A125	R189	S340	S340
H64	G126	E190	C259	E341
G65	P127	K191	Y267	F342
G66		E192	L268	K343
L67	M130	I193	L268	F344
P68	V131	I194	D271	F345
G69	V132	R195	K272	L346
A70	G133	Q196	V273	
S71	F134	A197	Q274	F349
S72	A135	A198	K275	E350
E73	M136	V199	Q279	
K74	L137	Q200	I280	
N75	G138	T202	R281	
K76	I139	K203	F282	
K77	L140	E204	Y283	
S78	H141	M205	E284	
	V142	V209	E285	
Q81	T143	V210	E286	
V82	K144	M213	E287	
K83	K145	F214	M288	
I84	K146	T215	G289	
C85	V147	A216	V291	
N86	F148	F217	F298	
Y87	E149	L218	S299	
V88	T150	P219	P300	
G89	L151	D220	V303	
P90	E152	S221	R304	
A91	A153	T222	R305	
K92	M155	Q223	Q306	
V93	M156	S224	F307	
I94	T156	F225	A308	
V95	E157	T226	I309	
Q96		R227	V310	
L97	I160	R228	Y316	
V98	R161	L229	K317	
T99	G162	E230	D318	
N100	Y163	P231	V319	
G101	M164	V232		
K102	P165	S234		
M103	G166			
I104	L167			
H105	H170			

## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	137.53Å 138.01Å 89.32Å 90.00° 97.25° 90.00°	Depositor
Resolution (Å)	20.00 – 2.75	Depositor
% Data completeness (in resolution range)	88.0 (20.00-2.75)	Depositor
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	CNS 0.9	Depositor
R, $R_{free}$	0.260 , 0.293	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	10439	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	68.0	wwPDB-VP



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	C	0.32	0/273	0.74	0/420
1	G	0.69	0/273	1.26	3/420 (0.7%)
2	D	0.32	0/271	0.78	0/416
2	H	0.80	0/271	1.31	2/416 (0.5%)
3	A	0.39	0/2236	0.67	0/3031
3	E	0.39	0/2236	0.69	0/3031
4	B	0.35	0/2514	0.60	0/3394
4	F	0.41	0/2514	0.66	0/3394
All	All	0.41	0/10588	0.71	5/14522 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	G	0	5
2	H	0	6
All	All	0	11

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	8	DT	O4'-C4'-C3'	-5.71	102.22	104.50
2	H	17	DA	O4'-C1'-C2'	5.32	110.16	105.90
1	G	7	DA	O4'-C1'-C2'	5.27	110.11	105.90
1	G	5	DA	N9-C1'-C2'	5.21	122.49	112.60
2	H	15	DG	N9-C1'-C2'	5.10	122.29	112.60

There are no chirality outliers.

5 of 11 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	G	11	DC	Sidechain
1	G	4	DG	Sidechain
1	G	5	DA	Sidechain
1	G	6	DA	Sidechain
1	G	7	DA	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	244	0	138	28	0
1	G	244	0	138	33	0
2	D	242	0	137	23	0
2	H	242	0	137	14	0
3	A	2184	0	2146	239	0
3	E	2184	0	2146	245	0
4	B	2462	0	2458	301	0
4	F	2462	0	2458	250	0
5	A	33	0	0	5	0
5	B	32	0	0	4	0
5	C	4	0	0	3	0
5	D	9	0	0	1	0
5	E	32	0	0	7	0
5	F	51	0	0	5	0
5	G	6	0	0	2	0
5	H	8	0	0	1	0
All	All	10439	0	9758	1074	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 54.

The worst 5 of 1074 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:1:DT:H3'	4:B:65:GLY:HA2	1.30	1.11
4:F:250:ILE:HG23	4:F:268:LEU:HD21	1.29	1.06
4:B:104:ILE:HG22	4:B:211:ARG:HH22	1.23	1.03

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:273:ARG:HD2	3:E:280:LEU:HD11	1.41	1.02
4:B:134:PHE:HB3	4:B:137:LEU:HD11	1.42	1.02

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	272/274 (99%)	204 (75%)	49 (18%)	19 (7%)	1	1
3	E	272/274 (99%)	201 (74%)	50 (18%)	21 (8%)	1	0
4	B	311/313 (99%)	236 (76%)	54 (17%)	21 (7%)	1	1
4	F	311/313 (99%)	240 (77%)	44 (14%)	27 (9%)	1	0
All	All	1166/1174 (99%)	881 (76%)	197 (17%)	88 (8%)	1	0

5 of 88 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	A	81	PRO
3	A	83	HIS
3	A	115	ASN
3	A	204	GLY
4	B	101	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	243/243 (100%)	218 (90%)	25 (10%)	7	12
3	E	243/243 (100%)	215 (88%)	28 (12%)	5	9
4	B	269/269 (100%)	244 (91%)	25 (9%)	9	15
4	F	269/269 (100%)	239 (89%)	30 (11%)	6	10
All	All	1024/1024 (100%)	916 (90%)	108 (10%)	7	11

5 of 108 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	E	81	PRO
3	E	236	ARG
4	F	227	ARG
3	E	104	LEU
3	E	154	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 52 such sidechains are listed below:

Mol	Chain	Res	Type
3	E	29	GLN
3	E	186	ASN
4	F	247	ASN
3	E	111	HIS
3	E	138	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

### 6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.