

wwPDB NMR Structure Validation Summary Report (i)

Jun 4, 2023 – 12:45 PM EDT

PDB ID	:	2LFH
BMRB ID	:	17753
Title	:	Solution NMR Structure of the Helix-loop-Helix Domain of Human ID3 Pro-
		tein, Northeast Structural Genomics Consortium Target HR3111A
Authors	:	Eletsky, A.; Wang, D.; Kohan, E.; Janjua, H.; Acton, T.B.; Xiao, R.; Ev-
		erett, J.K.; Montelione, G.T.; Szyperski, T.; Northeast Structural Genomics
		Consortium (NESG)
Deposited on	:	2011-06-30

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/NMRValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
BMRB Restraints Analysis	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.33

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment is 44%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# Entries)$	${f NMR} \ {f archive} \ (\#{f Entries})$
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain			
1	А	68	57%	•	41%	
1	В	68	60%	•	38%	



2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 5 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues				
Well-defined core Residue range (total) Backbone RMSD (Å) Medoid model				
1	A:27-A:66, B:27-B:68 (82)	1.10	5	

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 5 single-model clusters were found.

Cluster number	Models
1	2, 5, 8, 9, 10, 19, 20
2	7, 15, 16, 17
3	11, 14
4	1, 13
Single-model clusters	3; 4; 6; 12; 18



3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 2118 atoms, of which 1048 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called DNA-binding protein inhibitor ID-3.

Mol	Chain	Residues		Atoms					Trace
1	٨	69	Total	С	Н	Ν	Ο	S	0
	A	68	1059	332	524	102	97	4	0
1	В	69	Total	С	Η	Ν	0	S	0
	D	68	1059	332	524	102	97	4	0

There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	1	MET	-	expression tag	UNP Q02535
А	2	GLY	-	expression tag	UNP Q02535
А	3	HIS	-	expression tag	UNP Q02535
А	4	HIS	-	expression tag	UNP Q02535
А	5	HIS	-	expression tag	UNP Q02535
А	6	HIS	-	expression tag	UNP Q02535
A	7	HIS	-	expression tag	UNP Q02535
А	8	HIS	-	expression tag	UNP Q02535
А	9	SER	-	expression tag	UNP Q02535
A	10	HIS	-	expression tag	UNP Q02535
А	11	MET	-	expression tag	UNP Q02535
A	12	GLY	-	expression tag	UNP Q02535
A	13	GLY	-	expression tag	UNP Q02535
В	1	MET	-	expression tag	UNP Q02535
В	2	GLY	-	expression tag	UNP Q02535
В	3	HIS	-	expression tag	UNP Q02535
В	4	HIS	-	expression tag	UNP Q02535
В	5	HIS	-	expression tag	UNP Q02535
В	6	HIS	-	expression tag	UNP Q02535
В	7	HIS	-	expression tag	UNP Q02535
В	8	HIS	-	expression tag	UNP Q02535
В	9	SER	-	expression tag	UNP Q02535
В	10	HIS	-	expression tag	UNP Q02535
В	11	MET	-	expression tag	UNP Q02535
В	12	GLY	-	expression tag	UNP Q02535
В	13	GLY	-	expression tag	UNP Q02535



4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: DNA-binding protein inhibitor ID-3

Chain A:	57%	·	41%	
M G G C G C G C G C G C G C G C G C G C G	414 K115 F17 A18 A18 A18 A18 A18 A18 A18 A18 A18 A18	V68		
• Molecule 1: DN	A-binding protein inhib	oitor ID-3		
Chain B:	60%	·	38%	
555110 555110 53210	614 615 717 717 718 719 723 723 723 723 723 723 723 723 723 723			

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 5. Colouring as in section 4.1 above.

• Molecule 1: DNA-binding protein inhibitor ID-3

Chain A:	51%	7%	41%
M1 62 62 63 64 84 84 84 84 85 85 612 613 613 613 613 613 613 715	A18 A19 E20 E21 P22 L25 L25 L25 L26 L26 D27 D28 M29 M29 M29	V62 R57 V67 V68	
• Molecule 1: DNA-bir	ding protein inhibit	or ID-3	
Chain B:	53%	7% •	38%
M C2 C2 C2 C2 C2 C2 C2 C2 C2 C2 C2 C2 C2	A18 820 821 822 824 824 824 824 824 844 844	N52 V52 V67 V68	



5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: target function.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
TALOS+	geometry optimization	1.2009.0721.18
CNS	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	676
Number of shifts mapped to atoms	676
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	44%



6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	А	327	329	329	2 ± 1
1	В	342	347	347	2±1
All	All	13380	13520	13520	48

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

5 of 26 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total
1:B:27:ASP:HB3	1:B:30:ASN:HB3	0.55	1.77	13	2
1:A:36:LEU:HD11	1:B:55:LEU:HD22	0.55	1.79	10	1
1:A:62:ILE:HG21	1:B:39:LEU:HB3	0.50	1.84	12	2
1:B:40:VAL:HG11	1:B:57:ARG:HB3	0.49	1.83	9	1
1:B:43:VAL:HG22	1:B:57:ARG:HG2	0.49	1.85	5	1

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	40/68~(59%)	$37\pm2(92\pm5\%)$	$2\pm2~(6\pm4\%)$	$1\pm1~(2\pm2\%)$	9 46
1	В	41/68~(60%)	$38\pm2(91\pm4\%)$	$3\pm2~(6\pm4\%)$	$1\pm1~(2\pm2\%)$	10 49
All	All	1620/2720~(60%)	1484 (92%)	99~(6%)	37~(2%)	9 48

entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

5 of 14 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	А	49	LEU	5
1	В	67	VAL	4
1	А	47	THR	4
1	В	27	ASP	4
1	А	27	ASP	4

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	38/59~(64%)	$37 \pm 1 (97 \pm 2\%)$	$1\pm1 (3\pm2\%)$	49 91
1	В	40/59~(68%)	$39 \pm 1 (98 \pm 2\%)$	$1\pm1~(2\pm2\%)$	53 92
All	All	1560/2360~(66%)	1522 (98%)	38 (2%)	51 92

5 of 16 unique residues with a non-rotameric side chain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	А	57	ARG	7
1	А	35	ARG	4
1	В	57	ARG	4
1	В	35	ARG	3
1	В	53	GLU	3



6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 44% for the well-defined parts and 36% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: assigned_chem_shift_list_1

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	676
Number of shifts mapped to atoms	676
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	${\rm Correction}\pm{\rm precision},ppm$	Suggested action
$^{13}C_{\alpha}$	52	-0.45 ± 0.20	None needed (< 0.5 ppm)
$^{13}C_{\beta}$	50	0.35 ± 0.07	None needed (< 0.5 ppm)
$^{13}C'$	49	-0.32 ± 0.12	None needed (< 0.5 ppm)
¹⁵ N	44	0.62 ± 0.23	Should be applied

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 44%, i.e. 520 atoms were assigned a chemical shift out of a possible 1193. 0 out of 22 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathrm{H}$	$^{13}\mathrm{C}$	$^{15}\mathbf{N}$
Backbone	187/406~(46%)	76/164~(46%)	77/164~(47%)	34/78~(44%)
Sidechain	311/736~(42%)	212/478~(44%)	94/224~(42%)	5/34~(15%)

Continued on next page...



Continueu	jioni piceious puge	····		
	Total	$^{1}\mathbf{H}$	$^{13}\mathrm{C}$	$^{15}\mathbf{N}$
Aromatic	22/51~(43%)	10/24~(42%)	10/24~(42%)	2/3~(67%)
Overall	520/1193 (44%)	298/666~(45%)	181/412~(44%)	41/115 (36%)

Continued from previous page...


7.1.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:





8 NMR restraints analysis (i)

8.1 Conformationally restricting restraints (i)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	631
Intra-residue (i-j =0)	262
Sequential (i-j =1)	180
Medium range ($ i-j >1$ and $ i-j <5$)	131
Long range $(i-j \ge 5)$	58
Inter-chain	0
Hydrogen bond restraints	0
Disulfide bond restraints	0
Total dihedral-angle restraints	88
Number of unmapped restraints	0
Number of restraints per residue	5.3
Number of long range restraints per residue ¹	0.4

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations (i)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model (i)

Distance violations less than 0.1 Å are not included in the calculation. There are no distance violations

8.2.2 Average number of dihedral-angle violations per model (i)

Dihedral-angle violations less than 1° are not included in the calculation.

Bins $(^{\circ})$	Average number of violations per model	Max $(^{\circ})$
1.0-10.0 (Small)	2.4	7.5

Continued on next page...



Continued from previous page...

Bins ($^{\circ}$)	Average number of violations per model	Max ($^{\circ}$)
10.0-20.0 (Medium)	None	None
>20.0 (Large)	None	None



9 Distance violation analysis (i)


9.1 Summary of distance violations (i)

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Destroints type	Count	$\%^1$	Vio	lated	3	Consis	tently	y Violated ⁴
Restraints type	Count	70-	Count	$\%^2$	$\%^1$	Count	$\%^2$	$\%^1$
Intra-residue (i-j =0)	262	41.5	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	217	34.4	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	45	7.1	0	0.0	0.0	0	0.0	0.0
Sequential (i-j =1)	180	28.5	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	30	4.8	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	122	19.3	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	28	4.4	0	0.0	0.0	0	0.0	0.0
Medium range ($ i-j > 1 \& i-j < 5$)	131	20.8	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	29	4.6	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	59	9.4	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	43	6.8	0	0.0	0.0	0	0.0	0.0
Long range $(i-j \ge 5)$	58	9.2	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	13	2.1	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	45	7.1	0	0.0	0.0	0	0.0	0.0
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	631	100.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	59	9.4	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	411	65.1	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	161	25.5	0	0.0	0.0	0	0.0	0.0

 1 percentage calculated with respect to the total number of distance restraints, 2 percentage calculated with respect to the number of restraints in a particular restraint category, 3 violated in at least one model, 4 violated in all the models





9.1.1 Bar chart : Distribution of distance restraints and violations (i)

Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model (i)

No violations found

9.3 Distance violation statistics for the ensemble (i)

No violations found

9.4 Most violated distance restraints in the ensemble (i)

No violations found

9.5 All violated distance restraints (i)

No violations found



10 Dihedral-angle violation analysis (i)


10.1 Summary of dihedral-angle violations (i)

The following table provides the summary of dihedral-angle violations in different dihedral-angle types. Violations less than 1° are not included in the calculation.

Angle type	Count	Count $\%^1$					Consistently Violated ⁴		
Angle type	Count	70	Count	$\%^2$	$\%^1$	Count	$\%^2$	$\%^1$	
PSI	44	50.0	19	43.2	21.6	0	0.0	0.0	
PHI	44	50.0	9	20.5	10.2	0	0.0	0.0	
Total	88	100.0	28	31.8	31.8	0	0.0	0.0	

 1 percentage calculated with respect to total number of dihedral-angle restraints, 2 percentage calculated with respect to number of restraints in a particular dihedral-angle type, 3 violated in at least one model, 4 violated in all the models

10.1.1 Bar chart : Distribution of dihedral-angles and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories





10.2 Dihedral-angle violation statistics for each model (i)

The following table provides the dihedral-angle violation statistics for each model in the ensemble. Violations less than 1° are not included in the statistics.

Model ID	Nun	nber o	of violations	\mathbf{M}_{oon} (°)	M_{ov} (°)	SD (°)	Median ($^{\circ}$)	
Model ID	PSI	PHI	Total	Mean $(^{\circ})$	$Max (^{\circ})$	SD ()		
1	3	2	5	1.96	3.4	0.75	1.7	
2	4	1	5	2.6	5.7	1.56	1.8	
3	0	1	1	1.4	1.4	0.0	1.4	
4	2	1	3	3.33	7.5	2.95	1.3	
5	1	1	2	1.85	2.4	0.55	1.85	
6	2	2	4	1.2	1.3	0.07	1.2	
7	0	2	2	2.15	2.3	0.15	2.15	
8	1	0	1	1.1	1.1	0.0	1.1	
9	2	1	3	1.93	2.9	0.74	1.8	
10	3	0	3	1.83	2.8	0.69	1.5	
11	3	1	4	2.0	3.3	0.77	1.7	
12	0	0	0	0.0	0.0	0.0	0.0	
13	1	1	2	2.0	2.6	0.6	2.0	
14	1	1	2	2.85	3.8	0.95	2.85	
15	0	0	0	0.0	0.0	0.0	0.0	
16	0	0	0	0.0	0.0	0.0	0.0	
17	1	1	2	3.45	4.7	1.25	3.45	
18	3	2	5	2.18	3.1	0.78	1.9	
19	1	1	2	1.6	1.9	0.3	1.6	
20	0	1	1	1.3	1.3	0.0	1.3	







The mean(dot), median(x) and the standard deviation are shown in blue with respect to the y axis on the right

10.3 Dihedral-angle violation statistics for the ensemble (i)

Violation analysis may find that some restraints are violated in very few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of ensemble.

Nun	nber o	f violated restraints					
PSI	PHI	Total	Count^1	%			
13	4	17	1	5.0			
5	2	7	2	10.0			
0	2	2	3	15.0			
0	0	0	4	20.0			
1	1	2	5	25.0			
0	0	0	6	30.0			
0	0	0	7	35.0			
0	0	0	8	40.0			
0	0	0	9	45.0			
0	0	0	10	50.0			
0	0	0	11	55.0			


Continued on next page...



Nun	nber o	of violated restraints	Fraction of the ensemble					
PSI	PHI	Total	Count^1	%				
0	0	0	12	60.0				
0	0	0	13	65.0				
0	0	0	14	70.0				
0	0	0	15	75.0				
0	0	0	16	80.0				
0	0	0	17	85.0				
0	0	0	18	90.0				
0	0	0	19	95.0				
0	0	0	20	100.0				


Continued from previous page...

 1 Number of models with violations



PSI

PHI




10.4 Most violated dihedral-angle restraints in the ensemble (i)

10.4.1 Histogram : Distribution of mean dihedral-angle violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models



in the ensemble



10.4.2 Table: Most violated dihedral-angle restraints (i)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	$Models^1$	Mean	\mathbf{SD}^2	Median
(1,56)	1:B:56:GLN:N	1:B:56:GLN:CA	1:B:56:GLN:C	1:B:57:ARG:N	5	2.32	0.89	1.8
(1,37)	1:A:51:GLN:C	1:A:52:VAL:N	1:A:52:VAL:CA	1:A:52:VAL:C	5	1.98	0.82	1.8
(1,57)	1:A:56:GLN:C	1:A:57:ARG:N	1:A:57:ARG:CA	1:A:57:ARG:C	3	2.5	1.56	1.5
(1,2)	1:B:27:ASP:C	1:B:28:ASP:N	1:B:28:ASP:CA	1:B:28:ASP:C	3	1.53	0.26	1.4
(1,36)	1:B:36:LEU:N	1:B:36:LEU:CA	1:B:36:LEU:C	1:B:37:ARG:N	2	5.3	2.2	5.3
(1,52)	1:B:55:LEU:N	1:B:55:LEU:CA	1:B:55:LEU:C	1:B:56:GLN:N	2	2.6	0.7	2.6
(1,55)	1:A:56:GLN:N	1:A:56:GLN:CA	1:A:56:GLN:C	1:A:57:ARG:N	2	2.5	0.3	2.5
(1,38)	1:B:51:GLN:C	1:B:52:VAL:N	1:B:52:VAL:CA	1:B:52:VAL:C	2	2.45	0.65	2.45
(1,73)	1:A:60:ASP:C	1:A:61:TYR:N	1:A:61:TYR:CA	1:A:61:TYR:C	2	1.8	0.6	1.8
(1,12)	1:B:30:ASN:N	1:B:30:ASN:CA	1:B:30:ASN:C	1:B:31:HIS:N	2	1.65	0.05	1.65


¹ Number of violated models, ²Standard deviation, All angle values are in degree (°)



10.5 All violated dihedral-angle restraints (i)

10.5.1 Histogram : Distribution of violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



10.5.2 Table: All violated dihedral-angle restraints (i)

The following table provides the list of violations for the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	Model ID	Violation (°)
(1, 36)	1:B:36:LEU:N	1:B:36:LEU:CA	1:B:36:LEU:C	1:B:37:ARG:N	4	7.5
(1,35)	1:A:36:LEU:N	1:A:36:LEU:CA	1:A:36:LEU:C	1:A:37:ARG:N	2	5.7
(1,57)	1:A:56:GLN:C	1:A:57:ARG:N	1:A:57:ARG:CA	1:A:57:ARG:C	17	4.7
(1,56)	1:B:56:GLN:N	1:B:56:GLN:CA	1:B:56:GLN:C	1:B:57:ARG:N	14	3.8
(1,37)	1:A:51:GLN:C	1:A:52:VAL:N	1:A:52:VAL:CA	1:A:52:VAL:C	1	3.4
(1,52)	1:B:55:LEU:N	1:B:55:LEU:CA	1:B:55:LEU:C	1:B:56:GLN:N	11	3.3
(1,38)	1:B:51:GLN:C	1:B:52:VAL:N	1:B:52:VAL:CA	1:B:52:VAL:C	18	3.1
(1,36)	1:B:36:LEU:N	1:B:36:LEU:CA	1:B:36:LEU:C	1:B:37:ARG:N	18	3.1
(1,56)	1:B:56:GLN:N	1:B:56:GLN:CA	1:B:56:GLN:C	1:B:57:ARG:N	9	2.9
(1,55)	1:A:56:GLN:N	1:A:56:GLN:CA	1:A:56:GLN:C	1:A:57:ARG:N	10	2.8

