

wwPDB X-ray Structure Validation Summary Report (i)

Sep 11, 2023 – 07:26 PM EDT

PDB ID	:	4LG2
Title	:	Crystal structure of Reston Ebola virus VP35 RNA binding domain bound to
		12-bp dsRNA
Authors	:	Bale, S.; Julien, JP.; Bornholdt, Z.A.; Krois, A.S.; Wilson, I.A.; Saphire,
		E.O.
Deposited on	:	2013-06-27
Resolution	:	2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019) Refmac : 5.8.0158 CCP4 : 7.0.044 (Gargrove) Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)	MolProbity Xtriage (Phenix) EDS	:	
CCP4 : 7.0.044 (Gargrove) Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)	Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)	CCP4	:	7.0.044 (Gargrove)
Validation Pipeline (wwPDB-VP) : 2.35.1	Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)
RNA backbone	3102	1159 (3.00-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain						
1	А	146	% 65%	17%	·	16%			
1	В	146	68%	14%	•	16%			
1	С	146	61%	21%	•	16%			
1	D	146	68%	15%	•	16%			



Mol		Length	Quality of chain					
2	Е	12	75%		2	25%		
2	F	12	58%	17%	2	25%		
2	Ι	12	8%		2	25%		
2	J	12	42% 67%		17%	17%		



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 4703 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	А	122	Total	С	Ν	Ο	S	0	0	0
	A	122	952	609	169	168	6	0	0	0
1	В	122	Total	С	Ν	0	S	0	0	0
	D	122	952	609	169	168	6	0	0	0
1	С	122	Total	С	Ν	0	S	0	0	0
	U	122	952	609	169	168	6	0	0	0
1	П	122	Total	С	Ν	0	S	0	0	0
	D	122	952	609	169	168	6	0	0	U

• Molecule 1 is a protein called Polymerase cofactor.

There are 84 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	184	MET	-	expression tag	UNP Q8JPY0
А	185	ALA	-	expression tag	UNP Q8JPY0
А	186	HIS	-	expression tag	UNP Q8JPY0
A	187	HIS	-	expression tag	UNP Q8JPY0
А	188	HIS	-	expression tag	UNP Q8JPY0
A	189	HIS	-	expression tag	UNP Q8JPY0
А	190	HIS	-	expression tag	UNP Q8JPY0
A	191	HIS	-	expression tag	UNP Q8JPY0
А	192	VAL	-	expression tag	UNP Q8JPY0
А	193	ASP	-	expression tag	UNP Q8JPY0
А	194	ASP	-	expression tag	UNP Q8JPY0
А	195	ASP	-	expression tag	UNP Q8JPY0
А	196	ASP	-	expression tag	UNP Q8JPY0
А	197	LYS	-	expression tag	UNP Q8JPY0
А	198	GLU	-	expression tag	UNP Q8JPY0
А	199	ASN	-	expression tag	UNP Q8JPY0
А	200	LEU	-	expression tag	UNP Q8JPY0
А	201	TYR	-	expression tag	UNP Q8JPY0
А	202	PHE	-	expression tag	UNP Q8JPY0
А	203	GLN	-	expression tag	UNP Q8JPY0
А	204	SER	-	expression tag	UNP Q8JPY0



Chain	Residue	Modelled	Actual	Comment	Reference
В	184	MET	-	expression tag	UNP Q8JPY0
В	185	ALA	-	expression tag	UNP Q8JPY0
В	186	HIS	-	expression tag	UNP Q8JPY0
В	187	HIS	-	expression tag	UNP Q8JPY0
В	188	HIS	-	expression tag	UNP Q8JPY0
В	189	HIS	-	expression tag	UNP Q8JPY0
В	190	HIS	-	expression tag	UNP Q8JPY0
В	191	HIS	-	expression tag	UNP Q8JPY0
В	192	VAL	-	expression tag	UNP Q8JPY0
В	193	ASP	-	expression tag	UNP Q8JPY0
В	194	ASP	-	expression tag	UNP Q8JPY0
В	195	ASP	-	expression tag	UNP Q8JPY0
В	196	ASP	-	expression tag	UNP Q8JPY0
В	197	LYS	-	expression tag	UNP Q8JPY0
В	198	GLU	-	expression tag	UNP Q8JPY0
В	199	ASN	-	expression tag	UNP Q8JPY0
В	200	LEU	-	expression tag	UNP Q8JPY0
В	201	TYR	-	expression tag	UNP Q8JPY0
В	202	PHE	-	expression tag	UNP Q8JPY0
В	203	GLN	-	expression tag	UNP Q8JPY0
В	204	SER	-	expression tag	UNP Q8JPY0
С	184	MET	-	expression tag	UNP Q8JPY0
С	185	ALA	-	expression tag	UNP Q8JPY0
С	186	HIS	-	expression tag	UNP Q8JPY0
С	187	HIS	-	expression tag	UNP Q8JPY0
С	188	HIS	-	expression tag	UNP Q8JPY0
С	189	HIS	-	expression tag	UNP Q8JPY0
С	190	HIS	-	expression tag	UNP Q8JPY0
С	191	HIS	-	expression tag	UNP Q8JPY0
С	192	VAL	-	expression tag	UNP Q8JPY0
С	193	ASP	-	expression tag	UNP Q8JPY0
С	194	ASP	-	expression tag	UNP Q8JPY0
С	195	ASP	-	expression tag	UNP Q8JPY0
С	196	ASP	-	expression tag	UNP Q8JPY0
С	197	LYS	-	expression tag	UNP Q8JPY0
С	198	GLU	-	expression tag	UNP Q8JPY0
С	199	ASN	-	expression tag	UNP Q8JPY0
С	200	LEU	-	expression tag	UNP Q8JPY0
U		TYR	_	expression tag	UNP Q8JPY0
C	201	1 1 1 1			
	$\begin{array}{r} 201 \\ 202 \end{array}$	PHE	_	expression tag	UNP Q8JPY0
С					-



Chain	Residue	Modelled	Actual	Comment	Reference
D	184	MET	-	expression tag	UNP Q8JPY0
D	185	ALA	-	expression tag	UNP Q8JPY0
D	186	HIS	-	expression tag	UNP Q8JPY0
D	187	HIS	-	expression tag	UNP Q8JPY0
D	188	HIS	-	expression tag	UNP Q8JPY0
D	189	HIS	-	expression tag	UNP Q8JPY0
D	190	HIS	-	expression tag	UNP Q8JPY0
D	191	HIS	-	expression tag	UNP Q8JPY0
D	192	VAL	-	expression tag	UNP Q8JPY0
D	193	ASP	-	expression tag	UNP Q8JPY0
D	194	ASP	-	expression tag	UNP Q8JPY0
D	195	ASP	-	expression tag	UNP Q8JPY0
D	196	ASP	-	expression tag	UNP Q8JPY0
D	197	LYS	-	expression tag	UNP Q8JPY0
D	198	GLU	-	expression tag	UNP Q8JPY0
D	199	ASN	-	expression tag	UNP Q8JPY0
D	200	LEU	-	expression tag	UNP Q8JPY0
D	201	TYR	-	expression tag	UNP Q8JPY0
D	202	PHE	-	expression tag	UNP Q8JPY0
D	203	GLN	-	expression tag	UNP Q8JPY0
D	204	SER	-	expression tag	UNP Q8JPY0

• Molecule 2 is a RNA chain called dsRNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	Е	0	Total	С	Ν	Ο	Р	0	Ο	Ο
2	Ľ	3	187	85	33	61	8	0	0	0
2	F	9	Total	С	Ν	Ο	Р	0	0	0
2	Г	3	193	86	35	63	9	0	0	0
2	T	12	Total	С	Ν	Ο	Р	0	0	Ο
2	J	12	255	114	45	84	12	0	0	0
2	T	12	Total	С	Ν	Ο	Р	0	0	0
	1	12	252	114	45	82	11	0	0	0

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	Total O 1 1	0	0
3	В	5	Total O 5 5	0	0
3	С	1	Total O 1 1	0	0




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	J	1	Total O 1 1	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Polymerase cofactor

 \bullet Molecule 2: dsRNA

Chain E:	75%		25%
<mark>స్ రి</mark> ఎ ళ ల			
• Molecule 2: dsRNA			
Chain F:	58%	17%	25%
0 A 6 4 0 7 0 12 0 12 0 12			
\bullet Molecule 2: dsRNA			
Chain J:	67%	17%	17%
C1 U2 A3 A5 A5 C6 C12 C12 C12			
\bullet Molecule 2: dsRNA			
Chain I:	75%		25%
<mark>6 8 8 6 6</mark>			



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	55.09Å 72.87Å 175.55Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.30 - 2.70	Depositor
Resolution (A)	39.30 - 2.70	EDS
% Data completeness	99.5 (39.30-2.70)	Depositor
(in resolution range)	99.5 (39.30-2.70)	EDS
R _{merge}	0.08	Depositor
R _{sym}	0.09	Depositor
$< I/\sigma(I) > 1$	2.09 (at 2.69 Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
D D.	0.209 , 0.262	Depositor
R, R_{free}	0.212 , 0.264	DCC
R_{free} test set	1016 reflections (5.07%)	wwPDB-VP
Wilson B-factor $(Å^2)$	71.1	Xtriage
Anisotropy	0.393	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.36 , 42.8	EDS
L-test for twinning ²	$< L >=0.51, < L^2>=0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4703	wwPDB-VP
Average B, all atoms $(Å^2)$	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.99% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond	angles
	Ullaili	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.40	0/974	0.59	0/1317
1	В	0.39	0/974	0.60	0/1317
1	С	0.37	0/974	0.57	0/1317
1	D	0.38	0/974	0.59	0/1317
2	Е	0.33	0/208	0.78	0/322
2	F	0.46	0/215	0.79	0/333
2	Ι	0.39	0/281	0.85	0/436
2	J	0.40	0/284	0.90	0/440
All	All	0.39	0/4884	0.65	0/6799

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	952	0	984	18	0
1	В	952	0	984	18	0
1	С	952	0	984	19	0
1	D	952	0	984	8	0
2	Е	187	0	99	0	0
2	F	193	0	98	1	0
2	Ι	252	0	131	2	0
2	J	255	0	130	3	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	А	1	0	0	0	0
3	В	5	0	0	0	0
3	С	1	0	0	0	0
3	J	1	0	0	0	0
All	All	4703	0	4394	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 65 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:308:LYS:HB2	1:B:311:ARG:HG3	1.67	0.77
1:B:318:LYS:NZ	1:B:322:GLY:O	2.22	0.69
1:D:231:LEU:HD12	1:D:314:VAL:HG22	1.74	0.68
1:B:242:ASN:HD21	1:B:275:ILE:HG12	1.61	0.66
1:C:308:LYS:HB2	1:C:311:ARG:HG3	1.79	0.65

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	120/146~(82%)	120 (100%)	0	0	100	100
1	В	120/146~(82%)	119~(99%)	1 (1%)	0	100	100
1	\mathbf{C}	120/146~(82%)	119~(99%)	1 (1%)	0	100	100
1	D	120/146~(82%)	120 (100%)	0	0	100	100
All	All	480/584~(82%)	478 (100%)	2~(0%)	0	100	100

There are no Ramachandran outliers to report.



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	А	106/129~(82%)	99~(93%)	7~(7%)	16	38
1	В	106/129~(82%)	100 (94%)	6~(6%)	20	44
1	С	106/129~(82%)	97~(92%)	9~(8%)	10	24
1	D	106/129~(82%)	97~(92%)	9~(8%)	10	24
All	All	424/516~(82%)	393~(93%)	31 (7%)	14	33

5 of 31 residues with a non-rotameric side chain are listed below:

Mol	Chain	\mathbf{Res}	Type
1	С	242	ASN
1	D	261	SER
1	С	247	THR
1	D	287	ARG
1	D	242	ASN

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such side chains are listed below:

Mol	Chain	Res	Type
1	В	249	HIS

5.3.3 RNA (i)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	Е	8/12~(66%)	0	0
2	F	8/12~(66%)	0	0
2	Ι	$11/12 \ (91\%)$	1 (9%)	0
2	J	$11/12 \ (91\%)$	3~(27%)	1 (9%)
All	All	38/48~(79%)	4 (10%)	1 (2%)

All (4) RNA backbone outliers are listed below:



Mol	Chain	Res	Type
2	J	4	G
2	J	5	А
2	J	6	С
2	Ι	4	G

All (1) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	J	5	A

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	Q < 0.9
1	А	122/146~(83%)	0.26	1 (0%) 86 87	18, 29, 45, 53	0
1	В	122/146~(83%)	0.28	0 100 100	16, 33, 54, 63	0
1	С	122/146~(83%)	0.85	18 (14%) 2 1	33, 50, 66, 77	0
1	D	122/146~(83%)	0.59	12 (9%) 7 5	31, 51, 64, 68	0
2	Е	9/12~(75%)	0.43	0 100 100	27, 30, 76, 82	0
2	F	9/12~(75%)	0.55	2(22%) 0 0	17, 44, 94, 96	0
2	Ι	12/12~(100%)	0.32	1 (8%) 11 9	45, 51, 83, 84	0
2	J	12/12~(100%)	1.70	5~(41%) 0 0	37, 65, 98, 100	0
All	All	530/632~(83%)	0.52	39 (7%) 14 12	16, 44, 66, 100	0

The worst 5 of 39 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	219	ASP	6.2
1	D	224	PHE	5.7
1	С	224	PHE	4.5
1	С	284	ILE	4.5
1	С	320	GLN	4.4

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.

