



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 28, 2023 – 02:15 PM EDT

PDB ID : 3LHA  
Title : Crystal structure of mouse VPS26B(R240S/G241A/E242S) in spacegroup P41 21 2  
Authors : Collins, B.; Shaw, D.; Norwood, S.  
Deposited on : 2010-01-21  
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	340	 3% 56% 22% • 17%
1	B	340	 4% 56% 22% •• 17%

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 4855 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Vacuolar protein sorting-associated protein 26B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	283	2351	1506	396	443	6	0	1	0
1	B	282	2352	1507	396	443	6	0	1	0

There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	MET	-	expression tag	UNP Q8C0E2
A	-2	GLY	-	expression tag	UNP Q8C0E2
A	-1	SER	-	expression tag	UNP Q8C0E2
A	0	HIS	-	expression tag	UNP Q8C0E2
A	1	HIS	-	expression tag	UNP Q8C0E2
A	2	HIS	-	expression tag	UNP Q8C0E2
A	3	HIS	-	expression tag	UNP Q8C0E2
A	4	HIS	-	expression tag	UNP Q8C0E2
A	5	HIS	-	expression tag	UNP Q8C0E2
A	6	MET	-	expression tag	UNP Q8C0E2
A	197	SER	LEU	engineered mutation	UNP Q8C0E2
A	199	GLU	ARG	engineered mutation	UNP Q8C0E2
A	242	SER	GLU	engineered mutation	UNP Q8C0E2
B	-3	MET	-	expression tag	UNP Q8C0E2
B	-2	GLY	-	expression tag	UNP Q8C0E2
B	-1	SER	-	expression tag	UNP Q8C0E2
B	0	HIS	-	expression tag	UNP Q8C0E2
B	1	HIS	-	expression tag	UNP Q8C0E2
B	2	HIS	-	expression tag	UNP Q8C0E2
B	3	HIS	-	expression tag	UNP Q8C0E2
B	4	HIS	-	expression tag	UNP Q8C0E2
B	5	HIS	-	expression tag	UNP Q8C0E2
B	6	MET	-	expression tag	UNP Q8C0E2
B	197	SER	LEU	engineered mutation	UNP Q8C0E2
B	199	GLU	ARG	engineered mutation	UNP Q8C0E2

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	242	SER	GLU	engineered mutation	UNP Q8C0E2


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	93	Total O 93 93	0	0
2	B	59	Total O 59 59	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Vacuolar protein sorting-associated protein 26B



- Molecule 1: Vacuolar protein sorting-associated protein 26B



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	118.14Å 118.14Å 193.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.85 – 2.80 40.85 – 2.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (40.85-2.80) 100.0 (40.85-2.80)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.92 (at 2.81Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.5_2)	Depositor
R, $R_{free}$	0.252 , 0.284 0.256 , 0.288	Depositor DCC
$R_{free}$ test set	1734 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	52.3	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 53.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.88	EDS
Total number of atoms	4855	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	51.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.62% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.51	0/2401	0.65	0/3229
1	B	0.50	0/2402	0.68	2/3231 (0.1%)
All	All	0.51	0/4803	0.66	2/6460 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	60	ARG	CB-CG-CD	5.15	124.99	111.60
1	B	60	ARG	CG-CD-NE	5.13	122.57	111.80

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2351	0	2345	76	0
1	B	2352	0	2350	83	2
2	A	93	0	0	10	1
2	B	59	0	0	5	0
All	All	4855	0	4695	156	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 156 close contacts within the same asymmetric unit are listed below, sorted by

their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:179:ASN:ND2	1:B:190:VAL:HB	1.72	1.05
1:B:179:ASN:HD22	1:B:190:VAL:HB	1.22	1.04
1:A:179:ASN:ND2	1:A:190:VAL:HB	1.74	1.02
1:A:179:ASN:HD22	1:A:190:VAL:HB	1.25	0.99
1:B:171:CYS:HA	1:B:199:GLU:HB2	1.58	0.86

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:95:ARG:NH2	2:A:366:HOH:O[6_444]	2.11	0.09
1:B:196:LEU:O	1:B:196:LEU:O[7_555]	2.14	0.06

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	278/340 (82%)	256 (92%)	17 (6%)	5 (2%)	8	28
1	B	279/340 (82%)	254 (91%)	19 (7%)	6 (2%)	6	22
All	All	557/680 (82%)	510 (92%)	36 (6%)	11 (2%)	7	24

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	168	ILE
1	A	171	CYS
1	A	235	ASP
1	B	168	ILE
1	B	171	CYS



### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	261/308 (85%)	233 (89%)	28 (11%)	6	20
1	B	262/308 (85%)	236 (90%)	26 (10%)	8	23
All	All	523/616 (85%)	469 (90%)	54 (10%)	7	21

5 of 54 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	9	SER
1	B	54	LEU
1	B	230	LYS
1	B	10	VAL
1	B	31	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	123	ASN
1	B	179	ASN
1	B	288	GLN
1	B	273	ASN
1	A	288	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	283/340 (83%)	0.14	9 (3%) 47 37	26, 45, 93, 133	0
1	B	282/340 (82%)	0.16	14 (4%) 28 19	26, 45, 92, 132	0
All	All	565/680 (83%)	0.15	23 (4%) 37 27	26, 45, 92, 133	0

The worst 5 of 23 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	235	ASP	5.7
1	B	58	ASN	5.2
1	A	234	MET	4.9
1	B	99	ILE	4.0
1	B	234	MET	4.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.