



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 14, 2020 – 03:54 AM BST

PDB ID : 6LIY
Title : SeMet CRL Protein of Arabidopsis
Authors : Wang, F.F.; Guan, K.L.; Sun, P.K.; Xing, W.M.
Deposited on : 2019-12-13
Resolution : 1.76 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : **FAILED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.14.4.dev1

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 3412 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chromophore lyase CRL, chloroplastic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	S				Se
1	A	195	Total	C	N	O	S	Se	0	0	0
			1555	972	279	293	8	3			
1	B	196	Total	C	N	O	S	Se	0	0	0
			1549	967	278	293	8	3			

- Molecule 2 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
2	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
2	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	139	Total 139	O 139	0	0
3	B	145	Total 145	O 145	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS failed to run properly.


- Molecule 1: Chromophore lyase CRL, chloroplastic

Chain A: 



- Molecule 1: Chromophore lyase CRL, chloroplastic

Chain B: 



4 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	38.29Å 73.54Å 67.15Å 90.00° 97.16° 90.00°	Depositor
Resolution (Å)	32.19 – 1.76	Depositor
% Data completeness (in resolution range)	97.1 (32.19-1.76)	Depositor
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.96 (at 1.76Å)	Xtrriage
Refinement program	PHENIX 1.14_3260	Depositor
R, R_{free}	0.207 , 0.244	Depositor
Wilson B-factor (Å ²)	26.7	Xtrriage
Anisotropy	0.544	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	3412	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.44	0/1592	0.56	0/2152
1	B	0.44	0/1587	0.57	0/2149
All	All	0.44	0/3179	0.57	0/4301

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1555	0	1463	14	0
1	B	1549	0	1445	21	0
2	A	12	0	12	2	0
2	B	12	0	12	1	0
3	A	139	0	0	6	0
3	B	145	0	0	5	1
All	All	3412	0	2932	38	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:101:ARG:NH1	3:B:401:HOH:O	1.91	0.95
2:A:301:MES:O1S	3:A:401:HOH:O	1.98	0.82
1:B:143:ASP:OD1	3:B:402:HOH:O	2.04	0.75
1:B:133:ARG:CD	1:B:158:ASP:O	2.35	0.75
1:B:133:ARG:NE	1:B:159:ILE:O	2.22	0.71
1:B:133:ARG:HD3	1:B:158:ASP:O	1.90	0.71
1:A:139:LYS:HE3	3:A:438:HOH:O	1.92	0.68
1:B:117:CYS:SG	1:B:123:CYS:HB2	2.33	0.67
1:A:176:ARG:O	3:A:402:HOH:O	2.13	0.66
1:A:206:LYS:HD3	3:A:402:HOH:O	1.96	0.65
1:A:129:SER:HG	1:A:164:THR:HG1	1.41	0.64
1:A:86:MSE:SE	1:A:156:ILE:HD12	2.49	0.61
1:B:133:ARG:HD2	1:B:158:ASP:O	2.00	0.61
1:B:86:MSE:HG3	1:B:90:SER:HB2	1.87	0.56
1:B:54:ARG:HH22	1:B:82:PRO:HG3	1.72	0.55
1:B:133:ARG:HE	1:B:160:GLY:HA2	1.71	0.54
1:A:206:LYS:H	1:A:206:LYS:HD3	1.71	0.54
1:A:90:SER:OG	1:A:113:MSE:HE3	2.09	0.53
1:B:92:VAL:HG22	1:B:113:MSE:HG2	1.89	0.53
1:B:53:SER:O	1:B:57:THR:HG23	2.10	0.52
1:A:92:VAL:HG22	1:A:113:MSE:HG2	1.92	0.51
1:A:206:LYS:HE2	3:A:421:HOH:O	2.13	0.49
1:B:87:VAL:O	3:B:403:HOH:O	2.20	0.49
1:A:86:MSE:HB2	1:A:90:SER:HB3	1.94	0.48
1:A:53:SER:O	1:A:57:THR:HG23	2.14	0.48
1:A:75:ASN:HB3	1:A:100:TRP:CD1	2.50	0.46
1:B:86:MSE:CG	1:B:90:SER:HB2	2.46	0.44
1:B:219:GLU:HG3	3:B:476:HOH:O	2.18	0.43
1:A:86:MSE:HE3	1:A:86:MSE:HB3	1.94	0.43
2:B:301:MES:H32	2:B:301:MES:H82	1.46	0.43
1:B:133:ARG:HD3	1:B:158:ASP:C	2.39	0.42
1:B:50:ARG:O	1:B:54:ARG:HG2	2.19	0.42
1:A:101:ARG:NH1	3:A:405:HOH:O	2.43	0.42
1:B:206:LYS:HA	1:B:206:LYS:HD2	1.78	0.42
2:A:301:MES:H31	2:A:301:MES:H82	1.75	0.42
1:B:228:LYS:HG3	3:B:430:HOH:O	2.20	0.42
1:B:223:GLN:HB3	1:B:228:LYS:HE3	2.03	0.40
1:B:175:LYS:HB3	1:B:175:LYS:HE3	1.78	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:465:HOH:O	3:B:497:HOH:O[1_655]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	MES	A	301	-	12,12,12	1.92	2 (16%)	14,16,16	2.46	7 (50%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	MES	B	301	-	12,12,12	1.85	2 (16%)	14,16,16	3.33	7 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MES	A	301	-	-	4/6/14/14	0/1/1/1
2	MES	B	301	-	-	5/6/14/14	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	MES	C8-S	-5.87	1.69	1.77
2	B	301	MES	C8-S	-5.52	1.69	1.77
2	B	301	MES	O2S-S	2.45	1.52	1.45
2	A	301	MES	O2S-S	2.37	1.52	1.45

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	301	MES	O1S-S-C8	8.96	117.71	106.92
2	B	301	MES	O2S-S-C8	-4.58	101.39	106.92
2	A	301	MES	C5-N4-C3	3.70	117.16	108.83
2	B	301	MES	C2-C3-N4	-3.68	104.52	110.10
2	A	301	MES	C2-C3-N4	-3.53	104.75	110.10
2	A	301	MES	O1S-S-C8	3.38	110.98	106.92
2	B	301	MES	C5-N4-C3	3.36	116.38	108.83
2	A	301	MES	C6-C5-N4	-3.35	105.02	110.10
2	A	301	MES	C7-N4-C5	3.13	119.24	111.23
2	B	301	MES	O1-C6-C5	2.84	118.04	111.80
2	A	301	MES	O2S-S-C8	2.70	110.17	106.92
2	A	301	MES	O3S-S-C8	2.57	109.93	105.77
2	B	301	MES	C6-C5-N4	-2.57	106.21	110.10
2	B	301	MES	C7-N4-C3	2.56	117.79	111.23

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	MES	C7-C8-S-O1S
2	A	301	MES	C7-C8-S-O2S
2	A	301	MES	C7-C8-S-O3S
2	B	301	MES	C8-C7-N4-C3
2	B	301	MES	C7-C8-S-O1S
2	B	301	MES	C7-C8-S-O2S
2	B	301	MES	C7-C8-S-O3S
2	A	301	MES	C8-C7-N4-C5
2	B	301	MES	C8-C7-N4-C5

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	MES	2	0
2	B	301	MES	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS failed to run properly - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS failed to run properly - this section is therefore empty.

6.3 Carbohydrates

EDS failed to run properly - this section is therefore empty.

6.4 Ligands

EDS failed to run properly - this section is therefore empty.

6.5 Other polymers

EDS failed to run properly - this section is therefore empty.