

# wwPDB NMR Structure Validation Summary Report (i)

Jun 4, 2023 – 06:18 AM EDT

PDB ID : 2LNW BMRB ID : 18182

Title : Identification and structural basis for a novel interaction between Vav2 and

Arap3

Authors : Wu, B.; Zhang, J.; Wu, J.; Shi, Y.

Deposited on : 2012-01-05

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

wwPDB-RCI : v 1n 11 5 13 A (Berjanski et al., 2005)

PANAV : Wang et al. (2010)

 $\begin{array}{ccc} wwPDB\text{-}ShiftChecker &:& v1.2\\ BMRB \ Restraints \ Analysis &:& v1.2 \end{array}$ 

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.33

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $SOLUTION\ NMR$ 

The overall completeness of chemical shifts assignment is 86%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$egin{array}{c} { m NMR \ archive} \ { m (\#Entries)} \end{array}$		
Clashscore	158937	12864		
Ramachandran outliers	154571	11451		
Sidechain outliers	154315	11428		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain					
1	A	122	69%	14%	10%	7%		
2	В	9	44%	56%				



# 2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 8 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues							
Well-defined core	Residue range	(total)	Backbone RMSD (Å)	Medoid model			
1	A:668-A:768,	B:1409-	0.69	8			
	B:1412 (105)						

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 2 single-model clusters were found.

Cluster number Models							
1	2, 3, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 19, 20						
2	1, 4						
Single-model clusters	17; 18						



# 3 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 2027 atoms, of which 997 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called Guanine nucleotide exchange factor VAV2.

Mol	Chain	Residues	Atoms					Trace	
1	Λ	119	Total	С	Н	N	О	S	0
1 A	113	1887	607	935	163	180	2	U	

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	650	MET	-	expression tag	UNP P52735
A	651	GLY	-	expression tag	UNP P52735
A	652	HIS	-	expression tag	UNP P52735
A	653	HIS	-	expression tag	UNP P52735
A	654	HIS	-	expression tag	UNP P52735
A	655	HIS	-	expression tag	UNP P52735
A	656	HIS	-	expression tag	UNP P52735
A	657	HIS	-	expression tag	UNP P52735
A	658	MET	-	expression tag	UNP P52735

• Molecule 2 is a protein called Arf-GAP with Rho-GAP domain, ANK repeat and PH domain-containing protein 3.

Mol	Chain	Residues	Atoms				Trace		
9	D	0	Total	С	Н	N	О	Р	0
$\begin{array}{ c c c c c c c c c c c c c c c c c c c$	9	140	46	62	9	22	1		




# 4 Residue-property plots (i)

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: Guanine nucleotide exchange factor VAV2



 $\bullet$  Molecule 2: Arf-GAP with Rho-GAP domain, ANK repeat and PH domain-containing protein 3


Chain B: 44% 56%

E1404 E1405 P1406 V1407 Y1408 G1412

# 4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 8. Colouring as in section 4.1 above.

• Molecule 1: Guanine nucleotide exchange factor VAV2



E771

• Molecule 2: Arf-GAP with Rho-GAP domain, ANK repeat and PH domain-containing protein 3

Chain B: 44% 56%







#### 5 Refinement protocol and experimental data overview (i)



The models were refined using the following method: simulated annealing.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: structures with the lowest energy.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version	
CNS	refinement		

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1450
Number of shifts mapped to atoms	1450
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	86%



# 6 Model quality (i)

### 6.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PTR

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	852	840	836	9±3
2	В	30	24	24	0±1
All	All	17640	17280	17200	179

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

5 of 86 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total
1:A:752:LEU:HD13	1:A:759:LEU:HD23	0.81	1.51	4	1
1:A:673:TRP:CH2	1:A:744:VAL:HG21	0.74	2.17	3	2
1:A:673:TRP:CE3	1:A:697:ILE:HD12	0.69	2.23	15	3
1:A:737:PHE:CE2	1:A:743:LEU:HD23	0.68	2.23	12	2
1:A:756:PHE:CE2	2:B:1411:VAL:HG22	0.68	2.24	7	1



# 6.3 Torsion angles (i)

#### 6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	101/122 (83%)	93±3 (92±3%)	7±2 (7±2%)	1±1 (1±1%)	26	73
2	В	3/9~(33%)	3±0 (87±16%)	0±0 (13±16%)	0±0 (0±0%)	100	100
All	All	$2080/2620 \ (79\%)$	1909 (92%)	156 (8%)	15 (1%)	26	73

5 of 7 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	701	PRO	5
1	A	726	ASP	3
1	A	677	ASN	2
1	A	672	PRO	2
1	A	703	GLU	1

#### 6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed Rotameric		Outliers	Perce	entiles
1	A	92/112 (82%)	75±5 (82±5%)	17±5 (18±5%)	4	37
2	В	3/7 (43%)	3±0 (92±14%)	0±0 (8±14%)	15	62
All	All	1900/2380 (80%)	1558 (82%)	342 (18%)	4	38

5 of 58 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	682	GLN	18
1	A	680	ARG	13

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	753	LYS	13
1	A	686	LEU	11
1	A	725	LYS	11

#### 6.3.3 RNA (i)

There are no RNA molecules in this entry.

### 6.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with |Z| > 2 is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mal	Type	Chain	Dec	Tiple	Link Counts RMSZ		${ m gths}$
MIOI	туре	Chain	nes	Link	Counts	RMSZ	#Z>2
2	PTR	В	1408	2	15,16,17	$1.27 \pm 0.01$	1±0 (6±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with |Z| > 2 is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Dec	Link		Bond ang	eles
MIOI	туре	Chain	nes	Link	Counts	RMSZ	#Z>2
2	PTR	В	1408	2	19,22,24	$0.52 \pm 0.02$	0±0 (0±0%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PTR	В	1408	2	-	$0\pm0,10,11,13$	$0\pm0,1,1,1$

All unique bond outliers are listed below.

Mol	Chain	Pos	Type	Atoms	$\mathbf{z}$	$Observed(\AA)$	Ideal(Å)	Mod	dels
MIOI	Chain	nes	Type	Atoms	L	Observed(A)	Ideal(A)	Worst	Total
2	В	1408	PTR	P-O1P	3.48	1.61	1.50	12	20

There are no bond-angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

# 6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

# 6.6 Ligand geometry (i)

There are no ligands in this entry.

# 6.7 Other polymers (i)

There are no such molecules in this entry.

### 6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 86% for the well-defined parts and 84% for the entire structure.

#### 7.1 Chemical shift list 1

File name: working cs.cif

Chemical shift list name: assigned\_chem\_shift\_list\_1

#### 7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1450
Number of shifts mapped to atoms	1450
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	4

### 7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, $ppm$	Suggested action
$^{13}\mathrm{C}_{\alpha}$	111	$-0.37 \pm 0.14$	None needed ( $< 0.5 \text{ ppm}$ )
$^{13}C_{\beta}$	108	$0.08 \pm 0.17$	None needed (< 0.5 ppm)
<sup>13</sup> C′	110	$0.40 \pm 0.13$	None needed ( $< 0.5 \text{ ppm}$ )
$^{15}N$	107	$0.36 \pm 0.37$	None needed (< 0.5 ppm)

### 7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 86%, i.e. 1290 atoms were assigned a chemical shift out of a possible 1499. 0 out of 14 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathbf{H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Backbone	500/522~(96%)	204/210 (97%)	199/210 (95%)	97/102 (95%)
Sidechain	719/808 (89%)	502/520 (97%)	213/257 (83%)	4/31 (13%)

Continued on next page...



Continued from previous page...

	Total	$^{1}\mathbf{H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Aromatic	71/169 (42%)	71/81 (88%)	0/78 (0%)	0/10 (0%)
Overall	1290/1499~(86%)	777/811 (96%)	412/545 (76%)	101/143 (71%)

#### 7.1.4 Statistically unusual chemical shifts (i)


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	761	THR	HG1	6.28	0.08 - 2.19	24.4
1	В	1410	GLU	HG2	3.56	1.24 - 3.30	6.3
1	В	1410	GLU	HG3	3.56	1.20 - 3.30	6.3
1	A	743	LEU	HA	2.03	2.04 - 6.55	-5.0

#### 7.1.5 Random Coil Index (RCI) plots (i)


The image below reports random coil index values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:



Random coil index (RCI) for chain B:







# 8 NMR restraints analysis (i)

# 8.1 Conformationally restricting restraints (i)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	1794
Intra-residue ( i-j =0)	559
Sequential ( i-j =1)	428
Medium range ( $ i-j >1$ and $ i-j <5$ )	243
Long range ( i-j ≥5)	447
Inter-chain	71
Hydrogen bond restraints	46
Disulfide bond restraints	0
Total dihedral-angle restraints	119
Number of unmapped restraints	0
Number of restraints per residue	14.6
Number of long range restraints per residue <sup>1</sup>	3.7

<sup>&</sup>lt;sup>1</sup>Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

## 8.2 Residual restraint violations (i)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

# 8.2.1 Average number of distance violations per model (i)

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	0.2	0.19
0.2-0.5 (Medium)	None	None
>0.5 (Large)	None	None



### 8.2.2 Average number of dihedral-angle violations per model (i)

Dihedral-angle violations less than  $1^{\circ}$  are not included in the calculation. There are no dihedral-angle violations



# 9 Distance violation analysis (i)

# 9.1 Summary of distance violations (i)


The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Dostusints toma	C	<b>%</b> <sup>1</sup>	Vio	lated	3	Consis	stently Violated <sup>4</sup>		
Restraints type	Count	70	Count	$\%^2$	$\%^1$	Count	$\%^2$	$\%^1$	
Intra-residue ( i-j =0)	559	31.2	2	0.4	0.1	0	0.0	0.0	
Backbone-Backbone	104	5.8	0	0.0	0.0	0	0.0	0.0	
Backbone-Sidechain	432	24.1	1	0.2	0.1	0	0.0	0.0	
Sidechain-Sidechain	23	1.3	1	4.3	0.1	0	0.0	0.0	
Sequential ( i-j =1)	428	23.9	1	0.2	0.1	0	0.0	0.0	
Backbone-Backbone	177	9.9	1	0.6	0.1	0	0.0	0.0	
Backbone-Sidechain	232	12.9	0	0.0	0.0	0	0.0	0.0	
Sidechain-Sidechain	19	1.1	0	0.0	0.0	0	0.0	0.0	
Medium range ( $ i-j >1 \&  i-j <5$ )	243	13.5	0	0.0	0.0	0	0.0	0.0	
Backbone-Backbone	69	3.8	0	0.0	0.0	0	0.0	0.0	
Backbone-Sidechain	100	5.6	0	0.0	0.0	0	0.0	0.0	
Sidechain-Sidechain	74	4.1	0	0.0	0.0	0	0.0	0.0	
Long range ( $ i-j  \ge 5$ )	447	24.9	0	0.0	0.0	0	0.0	0.0	
Backbone-Backbone	68	3.8	0	0.0	0.0	0	0.0	0.0	
Backbone-Sidechain	174	9.7	0	0.0	0.0	0	0.0	0.0	
Sidechain-Sidechain	205	11.4	0	0.0	0.0	0	0.0	0.0	
Inter-chain	71	4.0	0	0.0	0.0	0	0.0	0.0	
Backbone-Backbone	4	0.2	0	0.0	0.0	0	0.0	0.0	
Backbone-Sidechain	22	1.2	0	0.0	0.0	0	0.0	0.0	
Sidechain-Sidechain	45	2.5	0	0.0	0.0	0	0.0	0.0	
Hydrogen bond	46	2.6	0	0.0	0.0	0	0.0	0.0	
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0	
Total	1794	100.0	3	0.2	0.2	0	0.0	0.0	
Backbone-Backbone	468	26.1	1	0.2	0.1	0	0.0	0.0	
Backbone-Sidechain	960	53.5	1	0.1	0.1	0	0.0	0.0	
Sidechain-Sidechain	366	20.4	1	0.3	0.1	0	0.0	0.0	

 $<sup>^1</sup>$  percentage calculated with respect to the total number of distance restraints,  $^2$  percentage calculated with respect to the number of restraints in a particular restraint category,  $^3$  violated in at least one model,  $^4$  violated in all the models



#### 9.1.1 Bar chart: Distribution of distance restraints and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

### 9.2 Distance violation statistics for each model (i)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

MadalID	Number of violations						M (8)	M (Å)	${ m SD}^6$ (Å)	Madian (Å)
Model ID	$IR^1$	$SQ^2$	$MR^3$	$LR^4$	$IC^5$	Total	Mean (Å)	Max (Å)	$SD^*(A)$	Median (Å)
1	0	0	0	0	0	0	0.0	0.0	0.0	0.0
2	0	0	0	0	0	0	0.0	0.0	0.0	0.0
3	1	0	0	0	0	1	0.19	0.19	0.0	0.19
4	0	0	0	0	0	0	0.0	0.0	0.0	0.0
5	0	1	0	0	0	1	0.11	0.11	0.0	0.11
6	0	0	0	0	0	0	0.0	0.0	0.0	0.0
7	0	0	0	0	0	0	0.0	0.0	0.0	0.0
8	0	0	0	0	0	0	0.0	0.0	0.0	0.0
9	0	0	0	0	0	0	0.0	0.0	0.0	0.0
10	1	0	0	0	0	1	0.16	0.16	0.0	0.16
11	0	0	0	0	0	0	0.0	0.0	0.0	0.0

Continued on next page...




Continued from previous page...

Model ID	Number of violations						Mean (Å)	Max (Å)	$SD^6$ (Å)	Median (Å)
Model 1D	$IR^1$	$SQ^2$	$ m MR^3$	$LR^4$	$IC^5$	Total	Mean (A)	Max (A)	SD'(A)	Median (A)
12	0	0	0	0	0	0	0.0	0.0	0.0	0.0
13	0	0	0	0	0	0	0.0	0.0	0.0	0.0
14	0	0	0	0	0	0	0.0	0.0	0.0	0.0
15	0	0	0	0	0	0	0.0	0.0	0.0	0.0
16	0	0	0	0	0	0	0.0	0.0	0.0	0.0
17	0	0	0	0	0	0	0.0	0.0	0.0	0.0
18	0	0	0	0	0	0	0.0	0.0	0.0	0.0
19	0	0	0	0	0	0	0.0	0.0	0.0	0.0
20	1	0	0	0	0	1	0.14	0.14	0.0	0.14

 $<sup>^1{\</sup>rm Intra-residue}$  restraints,  $^2{\rm Sequential}$  restraints,  $^3{\rm Medium}$  range restraints,  $^4{\rm Long}$  range restraints,  $^5{\rm Inter-chain}$  restraints,  $^6{\rm Standard}$  deviation

### 9.2.1 Bar graph: Distance Violation statistics for each model (i)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right



#### 9.3 Distance violation statistics for the ensemble (i)


Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 1745(IR:557, SQ:427, MR:243, LR:447, IC:71) restraints are not violated in the ensemble.

Nu	mber	of vio	lated	restr	aints	Fraction of the ensemble			
$IR^1$	$SQ^2$	$MR^3$	$LR^4$	$IC^5$	Total	Count <sup>6</sup>	%		
1	1	0	0	0	2	1	5.0		
1	0	0	0	0	1	2	10.0		
0	0	0	0	0	0	3	15.0		
0	0	0	0	0	0	4	20.0		
0	0	0	0	0	0	5	25.0		
0	0	0	0	0	0	6	30.0		
0	0	0	0	0	0	7	35.0		
0	0	0	0	0	0	8	40.0		
0	0	0	0	0	0	9	45.0		
0	0	0	0	0	0	10	50.0		
0	0	0	0	0	0	11	55.0		
0	0	0	0	0	0	12	60.0		
0	0	0	0	0	0	13	65.0		
0	0	0	0	0	0	14	70.0		
0	0	0	0	0	0	15	75.0		
0	0	0	0	0	0	16	80.0		
0	0	0	0	0	0	17	85.0		
0	0	0	0	0	0	18	90.0		
0	0	0	0	0	0	19	95.0		
0	0	0	0	0	0	20	100.0		

 $<sup>^1</sup>$ Intra-residue restraints,  $^2$ Sequential restraints,  $^3$ Medium range restraints,  $^4$ Long range restraints,  $^5$ Inter-chain restraints,  $^6$  Number of models with violations



#### 9.3.1 Bar graph: Distance violation statistics for the ensemble (i)




### 9.4 Most violated distance restraints in the ensemble (i)

# 9.4.1 Histogram : Distribution of mean distance violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





#### 9.4.2 Table: Most violated distance restraints (i)

The following table provides the mean and the standard deviation of the violation for each restraint sorted by number of violated models and the mean value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	${f Models^1}$	Mean (Å)	$\mathbf{SD}^1$ (Å)	Median (Å)
(1,1664)	2:B:1407:VAL:HA	2:B:1407:VAL:HG21	2	0.15	0.01	0.15
(1,1664)	2:B:1407:VAL:HA	2:B:1407:VAL:HG22	2	0.15	0.01	0.15
(1,1664)	2:B:1407:VAL:HA	2:B:1407:VAL:HG23	2	0.15	0.01	0.15


<sup>&</sup>lt;sup>1</sup>Number of violated models, <sup>2</sup>Standard deviation

# 9.5 All violated distance restraints (i)

# 9.5.1 Histogram : Distribution of distance violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.





#### 9.5.2 Table: All distance violations (i)

The following table lists the absolute value of the violation for each restraint in the ensemble sorted by its value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,659)	1:A:709:ILE:HD11	1:A:709:ILE:HB	3	0.19
(1,659)	1:A:709:ILE:HD12	1:A:709:ILE:HB	3	0.19
(1,659)	1:A:709:ILE:HD13	1:A:709:ILE:HB	3	0.19
(1,1664)	2:B:1407:VAL:HA	2:B:1407:VAL:HG21	10	0.16
(1,1664)	2:B:1407:VAL:HA	2:B:1407:VAL:HG22	10	0.16
(1,1664)	2:B:1407:VAL:HA	2:B:1407:VAL:HG23	10	0.16
(1,1664)	2:B:1407:VAL:HA	2:B:1407:VAL:HG21	20	0.14
(1,1664)	2:B:1407:VAL:HA	2:B:1407:VAL:HG22	20	0.14
(1,1664)	2:B:1407:VAL:HA	2:B:1407:VAL:HG23	20	0.14
(1,399)	1:A:691:ALA:H	1:A:690:HIS:HA	5	0.11



# 10 Dihedral-angle violation analysis (i)


# 10.1 Summary of dihedral-angle violations (i)

The following table provides the summary of dihedral-angle violations in different dihedral-angle types. Violations less than 1° are not included in the calculation.

Angle type	Count	071				Consistently Violated <sup>4</sup>		
Angle type	Count	70	Count	$\%^2$	$\%^1$	Count	$\%^2$	<b>%</b> ¹
PHI	60	50.4	0	0.0	0.0	0	0.0	0.0
PSI	59	49.6	0	0.0	0.0	0	0.0	0.0
Total	119	100.0	0	0.0	0.0	0	0.0	0.0

 $<sup>^1</sup>$  percentage calculated with respect to total number of dihedral-angle restraints,  $^2$  percentage calculated with respect to number of restraints in a particular dihedral-angle type,  $^3$  violated in at least one model,  $^4$  violated in all the models

#### 10.1.1 Bar chart: Distribution of dihedral-angles and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories



10.2	Dihedral-angle	violation	statistics	for	each i	model	i
10.4	Dincular-angle	VIOIGUIOII	SUGUISUICS	101	Cacii .	mouci	

No violations found

Dihedral-angle violation statistics for the ensemble (i) 10.3

No violations found

Most violated dihedral-angle restraints in the ensemble (i)

No violations found

All violated dihedral-angle restraints (i) 10.5

No violations found

