



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 16, 2023 – 10:25 AM JST

PDB ID : 6LNM
Title : Crystal structure of CASK-CaMK in complex with Mint1-CID
Authors : Cai, Q.; Zhang, M.
Deposited on : 2019-12-31
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	351	 73% 17% 7%
1	C	351	 2% 67% 21% 8%
1	E	351	 2% 70% 18% 8%
2	B	50	 54% 26% 16%
2	D	50	 2% 48% 22% 14% 16%
2	F	50	 2% 52% 24% 10% 14%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 8985 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peripheral plasma membrane protein CASK.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	325	2602	1670	448	467	17	0	1	0
1	C	322	2580	1656	445	462	17	0	0	0
1	E	322	2589	1664	445	463	17	0	1	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	GLY	-	expression tag	UNP Q62915
A	-4	PRO	-	expression tag	UNP Q62915
A	-3	GLY	-	expression tag	UNP Q62915
A	-2	SER	-	expression tag	UNP Q62915
A	-1	GLU	-	expression tag	UNP Q62915
A	0	PHE	-	expression tag	UNP Q62915
C	-5	GLY	-	expression tag	UNP Q62915
C	-4	PRO	-	expression tag	UNP Q62915
C	-3	GLY	-	expression tag	UNP Q62915
C	-2	SER	-	expression tag	UNP Q62915
C	-1	GLU	-	expression tag	UNP Q62915
C	0	PHE	-	expression tag	UNP Q62915
E	-5	GLY	-	expression tag	UNP Q62915
E	-4	PRO	-	expression tag	UNP Q62915
E	-3	GLY	-	expression tag	UNP Q62915
E	-2	SER	-	expression tag	UNP Q62915
E	-1	GLU	-	expression tag	UNP Q62915
E	0	PHE	-	expression tag	UNP Q62915

- Molecule 2 is a protein called Amyloid-beta A4 precursor protein-binding family A member 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	42	Total	C	N	O	S	0	1	0
			352	225	60	66	1			
2	D	42	Total	C	N	O	S	0	0	0
			344	220	57	66	1			
2	F	43	Total	C	N	O	S	0	0	0
			351	225	58	67	1			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	345	GLY	-	expression tag	UNP B2RUJ5
B	346	PRO	-	expression tag	UNP B2RUJ5
B	347	GLY	-	expression tag	UNP B2RUJ5
B	348	SER	-	expression tag	UNP B2RUJ5
B	349	GLU	-	expression tag	UNP B2RUJ5
B	350	PHE	-	expression tag	UNP B2RUJ5
D	345	GLY	-	expression tag	UNP B2RUJ5
D	346	PRO	-	expression tag	UNP B2RUJ5
D	347	GLY	-	expression tag	UNP B2RUJ5
D	348	SER	-	expression tag	UNP B2RUJ5
D	349	GLU	-	expression tag	UNP B2RUJ5
D	350	PHE	-	expression tag	UNP B2RUJ5
F	345	GLY	-	expression tag	UNP B2RUJ5
F	346	PRO	-	expression tag	UNP B2RUJ5
F	347	GLY	-	expression tag	UNP B2RUJ5
F	348	SER	-	expression tag	UNP B2RUJ5
F	349	GLU	-	expression tag	UNP B2RUJ5
F	350	PHE	-	expression tag	UNP B2RUJ5


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	43	Total	O	0	0
			43	43		
3	B	5	Total	O	0	0
			5	5		
3	C	64	Total	O	0	0
			64	64		
3	D	5	Total	O	0	0
			5	5		
3	E	44	Total	O	0	0
			44	44		
3	F	6	Total	O	0	0
			6	6		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Peripheral plasma membrane protein CASK




- Molecule 1: Peripheral plasma membrane protein CASK



- Molecule 1: Peripheral plasma membrane protein CASK





- Molecule 2: Amyloid-beta A4 precursor protein-binding family A member 1



- Molecule 2: Amyloid-beta A4 precursor protein-binding family A member 1



- Molecule 2: Amyloid-beta A4 precursor protein-binding family A member 1



4 Data and refinement statistics i

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, α , β , γ	151.42Å 151.42Å 49.45Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	41.40 – 2.40 41.40 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.5 (41.40-2.40) 99.5 (41.40-2.40)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.57 (at 2.39Å)	Xtrriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.171 , 0.213 0.160 , 0.200	Depositor DCC
R_{free} test set	2384 reflections (4.81%)	wwPDB-VP
Wilson B-factor (Å ²)	31.0	Xtrriage
Anisotropy	0.080	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 9.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.38$, $\langle L^2 \rangle = 0.21$	Xtrriage
Estimated twinning fraction	0.127 for -h,-k,l 0.129 for h,-h-k,-l 0.416 for -k,-h,-l	Xtrriage
Reported twinning fraction	0.570 for H, K, L 0.430 for -K, -H, -L	Depositor
Outliers	0 of 49551 reflections	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8985	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.71	1/2669 (0.0%)	0.86	0/3602
1	C	0.71	1/2644 (0.0%)	0.88	2/3568 (0.1%)
1	E	0.70	0/2657	0.87	0/3586
2	B	0.70	0/362	0.99	0/488
2	D	0.74	0/351	0.94	0/474
2	F	0.67	0/359	0.98	0/485
All	All	0.71	2/9042 (0.0%)	0.88	2/12203 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	E	0	1
All	All	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	76	GLU	CD-OE2	-5.72	1.19	1.25
1	C	92	GLU	CD-OE2	-5.03	1.20	1.25

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	28	ARG	NE-CZ-NH1	-5.77	117.42	120.30
1	C	32	ARG	CG-CD-NE	5.41	123.17	111.80

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	334	ASP	Peptide
1	E	334	ASP	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2602	0	2595	44	1
1	C	2580	0	2572	64	0
1	E	2589	0	2581	62	0
2	B	352	0	364	9	0
2	D	344	0	351	22	0
2	F	351	0	359	13	0
3	A	43	0	0	5	0
3	B	5	0	0	0	0
3	C	64	0	0	9	0
3	D	5	0	0	3	0
3	E	44	0	0	4	0
3	F	6	0	0	3	0
All	All	8985	0	8822	194	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 194 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:228:GLU:OE1	1:E:232:LYS:HE2	1.48	1.12
1:E:174:LEU:HD12	1:E:195:PRO:HB2	1.45	0.99
1:E:95:ASP:HB3	1:E:151:SER:HA	1.46	0.96
1:A:174:LEU:HD12	1:A:195:PRO:HB2	1.48	0.94
1:C:155:SER:HB2	3:C:423:HOH:O	1.69	0.92

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:153:GLU:OE2	1:A:228:GLU:OE2[1_554]	2.12	0.08

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	322/351 (92%)	317 (98%)	4 (1%)	1 (0%)	41	55
1	C	318/351 (91%)	312 (98%)	6 (2%)	0	100	100
1	E	319/351 (91%)	313 (98%)	6 (2%)	0	100	100
2	B	41/50 (82%)	38 (93%)	1 (2%)	2 (5%)	2	1
2	D	40/50 (80%)	37 (92%)	3 (8%)	0	100	100
2	F	41/50 (82%)	39 (95%)	1 (2%)	1 (2%)	6	6
All	All	1081/1203 (90%)	1056 (98%)	21 (2%)	4 (0%)	41	48

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	387[A]	ARG
2	B	387[B]	ARG
2	F	387	ARG
1	A	329	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	282/299 (94%)	258 (92%)	24 (8%)	10	16
1	C	280/299 (94%)	245 (88%)	35 (12%)	4	5
1	E	281/299 (94%)	254 (90%)	27 (10%)	8	12
2	B	40/46 (87%)	32 (80%)	8 (20%)	1	1
2	D	39/46 (85%)	30 (77%)	9 (23%)	1	1
2	F	40/46 (87%)	31 (78%)	9 (22%)	1	1
All	All	962/1035 (93%)	850 (88%)	112 (12%)	5	7

5 of 112 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	232	LYS
2	F	386	MET
2	D	365	VAL
2	F	378	GLU
1	E	315	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	318	ASN
1	E	241	GLN
1	C	315	HIS
1	E	238	ASN
1	C	294	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	325/351 (92%)	-0.67	0 100 100	15, 28, 55, 72	4 (1%)
1	C	322/351 (91%)	-0.74	2 (0%) 89 88	15, 29, 53, 88	4 (1%)
1	E	322/351 (91%)	-0.71	3 (0%) 84 82	18, 29, 50, 87	4 (1%)
2	B	42/50 (84%)	-0.36	0 100 100	19, 37, 70, 101	0
2	D	42/50 (84%)	-0.35	1 (2%) 59 57	19, 34, 89, 125	0
2	F	43/50 (86%)	-0.24	1 (2%) 60 58	23, 41, 88, 117	0
All	All	1096/1203 (91%)	-0.66	7 (0%) 89 88	15, 30, 56, 125	12 (1%)

The worst 5 of 7 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	378	GLU	4.0
1	C	334	ASP	3.0
1	C	328	LEU	3.0
2	F	378	GLU	2.9
1	E	329	PRO	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.