



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 12, 2024 – 07:53 PM EDT

PDB ID : 3LOM
Title : CRYSTAL STRUCTURE OF GERANYLTRANSFERASE FROM *Legionella pneumophila*
Authors : Patskovsky, Y.; Toro, R.; Rutter, M.; Sauder, J.M.; Burley, S.K.; Almo, S.C.;
New York SGX Research Center for Structural Genomics (NYSGXRC)
Deposited on : 2010-02-04
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 2.36.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	313	 13% 80% 11% • 8%
1	B	313	 10% 80% 10% • 9%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PO4	B	3	-	X	-	-

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 4641 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Geranyltranstransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	288	2254	1432	384	426	12	0	2	0
1	B	285	2237	1423	379	421	14	0	4	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	5	MET	-	expression tag	UNP Q5ZT35
A	6	SER	-	expression tag	UNP Q5ZT35
A	310	GLU	-	expression tag	UNP Q5ZT35
A	311	GLY	-	expression tag	UNP Q5ZT35
A	312	HIS	-	expression tag	UNP Q5ZT35
A	313	HIS	-	expression tag	UNP Q5ZT35
A	314	HIS	-	expression tag	UNP Q5ZT35
A	315	HIS	-	expression tag	UNP Q5ZT35
A	316	HIS	-	expression tag	UNP Q5ZT35
A	317	HIS	-	expression tag	UNP Q5ZT35
B	5	MET	-	expression tag	UNP Q5ZT35
B	6	SER	-	expression tag	UNP Q5ZT35
B	310	GLU	-	expression tag	UNP Q5ZT35
B	311	GLY	-	expression tag	UNP Q5ZT35
B	312	HIS	-	expression tag	UNP Q5ZT35
B	313	HIS	-	expression tag	UNP Q5ZT35
B	314	HIS	-	expression tag	UNP Q5ZT35
B	315	HIS	-	expression tag	UNP Q5ZT35
B	316	HIS	-	expression tag	UNP Q5ZT35
B	317	HIS	-	expression tag	UNP Q5ZT35

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	67	Total	O	0	0
			67	67		
3	B	68	Total	O	0	0
			68	68		

4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	94.78Å 94.78Å 148.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 2.30 36.80 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.6 (40.00-2.30) 99.6 (36.80-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.72 (at 2.29Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.221 , 0.267 0.216 , 0.257	Depositor DCC
R_{free} test set	976 reflections (3.18%)	wwPDB-VP
Wilson B-factor (Å ²)	40.8	Xtrriage
Anisotropy	0.877	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 48.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4641	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/2291	0.59	0/3097
1	B	0.40	0/2279	0.60	0/3080
All	All	0.40	0/4570	0.60	0/6177

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2254	0	2315	15	0
1	B	2237	0	2304	13	0
2	A	5	0	0	0	0
2	B	10	0	0	0	0
3	A	67	0	0	2	0
3	B	68	0	0	0	0
All	All	4641	0	4619	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:208:GLN:HB2	1:A:211:VAL:HG22	1.76	0.68
1:A:40:PRO:HA	1:A:43:ARG:HD3	1.79	0.62
1:B:270:LEU:O	1:B:274:ILE:HG12	2.03	0.59
1:A:270:LEU:O	1:A:274:ILE:HG12	2.02	0.58
1:A:180:VAL:HB	1:A:259:THR:HG21	1.86	0.57
1:B:135:LEU:HD22	1:B:151:ILE:HD11	1.86	0.56
1:A:132:ILE:HD13	1:B:132:ILE:HG13	1.89	0.54
1:B:147:GLN:O	1:B:151:ILE:HG23	2.07	0.54
1:A:9:SER:HA	1:A:12:MET:HB2	1.92	0.52
1:B:137[A]:MET:HG3	1:B:138:ARG:HG2	1.94	0.50
1:B:45:ALA:HB3	1:B:123:VAL:HG21	1.94	0.49
1:A:272:GLU:O	1:A:276:VAL:HG23	2.13	0.48
1:B:49:ALA:HB1	1:B:92:ILE:HD11	1.97	0.46
1:B:189:HIS:CG	1:B:231:GLN:HG2	2.51	0.45
1:A:45:ALA:HB3	1:A:123:VAL:HG21	1.99	0.44
1:A:31:LEU:HD11	1:A:134:VAL:HG22	1.99	0.44
1:B:52:SER:HA	1:B:108:LYS:HE2	2.00	0.44
1:A:32:LEU:HD11	1:A:50:LEU:HD12	2.00	0.43
1:A:52:SER:H	1:A:56[B]:ARG:HH22	1.66	0.43
1:A:164:MET:HE3	3:A:323:HOH:O	2.17	0.43
1:B:135:LEU:O	1:B:139:LEU:HB2	2.19	0.43
1:A:41:GLN:HG2	3:A:336:HOH:O	2.19	0.42
1:B:58:ARG:O	1:B:62:VAL:HG23	2.19	0.42
1:A:237:LEU:O	1:A:242:PRO:HD2	2.20	0.42
1:B:35:ILE:O	1:B:43:ARG:NH1	2.47	0.41
1:A:266:ASN:HD21	1:A:268:GLN:HG2	1.85	0.41
1:B:78:ILE:H	1:B:78:ILE:HG12	1.63	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	286/313 (91%)	278 (97%)	8 (3%)	0	100	100
1	B	285/313 (91%)	277 (97%)	8 (3%)	0	100	100
All	All	571/626 (91%)	555 (97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	247/266 (93%)	228 (92%)	19 (8%)	13	16
1	B	246/266 (92%)	229 (93%)	17 (7%)	15	20
All	All	493/532 (93%)	457 (93%)	36 (7%)	14	18

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26	LEU
1	A	35	ILE
1	A	43	ARG
1	A	55	LYS
1	A	56[A]	ARG
1	A	56[B]	ARG
1	A	83	LEU
1	A	89	TYR
1	A	105	ARG
1	A	143	LEU
1	A	153	GLN
1	A	156	VAL
1	A	211	VAL
1	A	259	THR
1	A	266	ASN
1	A	268	GLN
1	A	279	GLN
1	A	287	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	305	ARG
1	B	9	SER
1	B	32	LEU
1	B	50	LEU
1	B	71	VAL
1	B	78	ILE
1	B	89	TYR
1	B	138	ARG
1	B	147	GLN
1	B	151	ILE
1	B	153	GLN
1	B	156	VAL
1	B	198	LEU
1	B	259	THR
1	B	268	GLN
1	B	279	GLN
1	B	282	MET
1	B	287	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	11	GLN
1	A	30	GLN
1	A	33	ASN
1	A	101	ASN
1	A	157	ASN
1	A	268	GLN
1	A	279	GLN
1	A	301	GLN
1	A	303	GLN
1	B	101	ASN
1	B	269	GLN
1	B	279	GLN
1	B	303	GLN

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PO4	B	3	-	4,4,4	2.17	3 (75%)	6,6,6	1.55	1 (16%)
2	PO4	A	2	-	4,4,4	0.90	0	6,6,6	0.55	0
2	PO4	B	1	-	4,4,4	1.06	0	6,6,6	0.48	0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	3	PO4	P-O3	-2.43	1.47	1.54
2	B	3	PO4	P-O4	-2.42	1.47	1.54
2	B	3	PO4	P-O2	-2.34	1.47	1.54

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	3	PO4	O3-P-O1	-2.36	102.59	110.95

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	288/313 (92%)	0.73	42 (14%) 2 3	34, 58, 111, 151	0
1	B	285/313 (91%)	0.49	32 (11%) 5 7	33, 53, 90, 116	0
All	All	573/626 (91%)	0.61	74 (12%) 3 5	33, 56, 99, 151	0

All (74) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	308	LEU	7.9
1	B	178	SER	6.9
1	A	11	GLN	6.0
1	A	8	ASN	5.8
1	B	308	LEU	5.7
1	A	242	PRO	5.5
1	B	176	ALA	5.4
1	A	9	SER	5.3
1	A	307	ASN	5.1
1	A	287	LEU	5.1
1	A	13	ASN	5.0
1	A	304	ASN	5.0
1	B	239	LEU	4.9
1	A	10	THR	4.7
1	A	12	MET	4.6
1	B	307	ASN	4.6
1	B	209	HIS	4.5
1	A	310	GLU	4.4
1	A	241	ALA	4.3
1	A	55	LYS	4.3
1	B	8	ASN	4.2
1	B	240	TYR	3.9
1	B	89	TYR	3.5
1	A	69	ILE	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	237	LEU	3.4
1	A	196	LEU	3.4
1	A	197	ILE	3.3
1	A	129	ALA	3.2
1	A	54	GLY	3.2
1	A	309	SER	3.1
1	A	215	ILE	3.1
1	A	210	GLU	2.9
1	A	199	ALA	2.9
1	A	71	VAL	2.8
1	A	211	VAL	2.8
1	A	301	GLN	2.8
1	A	179	SER	2.8
1	A	83	LEU	2.7
1	B	236	TYR	2.7
1	B	196	LEU	2.7
1	A	177	LYS	2.7
1	A	193	THR	2.7
1	A	89	TYR	2.7
1	A	176	ALA	2.7
1	A	56[A]	ARG	2.7
1	B	258	LYS	2.6
1	B	179	SER	2.6
1	B	177	LYS	2.6
1	B	86	THR	2.5
1	A	175	LEU	2.5
1	A	132	ILE	2.5
1	A	164	MET	2.5
1	A	82	ALA	2.5
1	B	128	GLN	2.5
1	B	159	SER	2.4
1	A	209	HIS	2.4
1	A	178	SER	2.4
1	A	127	MET	2.3
1	A	203	MET	2.3
1	B	156	VAL	2.3
1	A	131	ALA	2.3
1	B	127	MET	2.3
1	B	306	SER	2.3
1	B	141	PRO	2.2
1	B	241	ALA	2.2
1	B	267	LYS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	9	SER	2.2
1	B	11	GLN	2.2
1	B	15	GLN	2.1
1	B	197	ILE	2.1
1	B	142	LEU	2.1
1	B	91	LEU	2.0
1	B	143	LEU	2.0
1	B	264	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q < 0.9
2	PO4	A	2	5/5	0.82	0.44	40,53,63,66	5
2	PO4	B	3	5/5	0.96	0.35	44,56,64,65	5
2	PO4	B	1	5/5	0.99	0.15	49,49,59,61	0

6.5 Other polymers [i](#)

There are no such residues in this entry.