



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 19, 2023 – 04:55 PM JST

PDB ID : 6LPD  
Title : Phascolosoma esculenta  
Authors : Su, X.R.; Ming, T.H.  
Deposited on : 2020-01-09  
Resolution : 1.65 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<https://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1827 (1.66-1.66)
Clashscore	141614	1931 (1.66-1.66)
Ramachandran outliers	138981	1891 (1.66-1.66)
Sidechain outliers	138945	1891 (1.66-1.66)
RSRZ outliers	127900	1791 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	174	 83% 13% ..
1	B	174	 84% 12% ..
1	C	174	 83% 12% ..
1	D	174	 83% 13% ..
1	E	174	 83% 15% .
1	F	174	 84% 13% ..

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 10693 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ferritin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	170	1562	972	264	320	6	0	23	0
1	B	170	1550	959	267	317	7	0	20	0
1	C	170	1542	952	268	315	7	0	18	0
1	D	170	1551	958	267	320	6	0	19	0
1	E	170	1521	944	264	306	7	0	17	0
1	F	170	1494	929	256	303	6	0	16	0

- Molecule 2 is FE (II) ION (three-letter code: FE2) (formula: Fe) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total 2	Fe 2	0	0
2	B	3	Total 3	Fe 3	0	0
2	C	2	Total 2	Fe 2	0	0
2	D	2	Total 2	Fe 2	0	0
2	E	2	Total 2	Fe 2	0	0
2	F	2	Total 2	Fe 2	0	0

- Molecule 3 is FE (III) ION (three-letter code: FE) (formula: Fe) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	F	1	Total Fe 1 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	256	Total O 256 256	0	0
4	B	232	Total O 232 232	0	0
4	C	247	Total O 247 247	0	0
4	D	241	Total O 241 241	0	0
4	E	238	Total O 238 238	0	0
4	F	245	Total O 245 245	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ferritin




- Molecule 1: Ferritin




- Molecule 1: Ferritin




- Molecule 1: Ferritin



- Molecule 1: Ferritin



- Molecule 1: Ferritin

Chain F:  % 84% 13% ..



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	153.00Å 153.49Å 153.82Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	108.65 – 1.65 34.25 – 1.65	Depositor EDS
% Data completeness (in resolution range)	96.4 (108.65-1.65) 95.4 (34.25-1.65)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	14.74 (at 1.65Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, $R_{free}$	0.108 , 0.153 0.135 , 0.169	Depositor DCC
$R_{free}$ test set	10411 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	13.3	Xtriage
Anisotropy	0.361	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 49.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.427 for -h,-l,-k 0.399 for l,-k,h 0.419 for -k,-h,-l 0.400 for k,-l,-h 0.400 for -l,h,-k	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	10693	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.94% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FE2, FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	1.35	1/1623 (0.1%)	1.23	10/2179 (0.5%)
1	B	1.33	0/1593	1.19	9/2140 (0.4%)
1	C	1.28	0/1571	1.10	5/2108 (0.2%)
1	D	1.36	0/1589	1.22	11/2134 (0.5%)
1	E	1.40	6/1571 (0.4%)	1.20	9/2108 (0.4%)
1	F	1.37	1/1552 (0.1%)	1.18	8/2084 (0.4%)
All	All	1.35	8/9499 (0.1%)	1.19	52/12753 (0.4%)

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	90[A]	GLU	CD-OE1	-6.84	1.18	1.25
1	E	90[B]	GLU	CD-OE1	-6.84	1.18	1.25
1	E	90[A]	GLU	CA-C	5.36	1.66	1.52
1	E	90[B]	GLU	CA-C	5.36	1.66	1.52
1	F	60	GLU	CD-OE2	-5.30	1.19	1.25

The worst 5 of 52 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	77	ARG	NE-CZ-NH2	-7.68	116.46	120.30
1	F	129[A]	ASP	CA-C-O	6.37	133.47	120.10
1	F	129[B]	ASP	CA-C-O	6.37	133.47	120.10
1	E	85	ARG	NE-CZ-NH1	-6.11	117.24	120.30
1	A	135[A]	PHE	CB-CG-CD2	6.08	125.05	120.80

There are no chirality outliers.

There are no planarity outliers.



## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1562	0	1474	18	0
1	B	1550	0	1446	17	0
1	C	1542	0	1445	28	0
1	D	1551	0	1449	19	0
1	E	1521	0	1455	18	0
1	F	1494	0	1432	24	0
2	A	2	0	0	0	0
2	B	3	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
2	E	2	0	0	0	0
2	F	2	0	0	0	0
3	F	1	0	0	0	0
4	A	256	0	0	10	0
4	B	232	0	0	8	0
4	C	247	0	0	10	0
4	D	241	0	0	9	0
4	E	238	0	0	9	0
4	F	245	0	0	13	0
All	All	10693	0	8701	121	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 121 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:63[C]:HIS:CE1	4:E:306:HOH:O	1.80	1.24
1:F:63[B]:HIS:CE1	4:F:333:HOH:O	1.89	1.22
1:A:63[B]:HIS:CE1	4:A:329:HOH:O	1.93	1.20
1:D:63[B]:HIS:CE1	4:D:330:HOH:O	1.97	1.16
1:B:63[B]:HIS:CE1	4:B:412:HOH:O	1.97	1.15

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	194/174 (112%)	187 (96%)	7 (4%)	0	100	100
1	B	190/174 (109%)	184 (97%)	6 (3%)	0	100	100
1	C	187/174 (108%)	182 (97%)	5 (3%)	0	100	100
1	D	189/174 (109%)	184 (97%)	5 (3%)	0	100	100
1	E	187/174 (108%)	182 (97%)	5 (3%)	0	100	100
1	F	186/174 (107%)	182 (98%)	4 (2%)	0	100	100
All	All	1133/1044 (108%)	1101 (97%)	32 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	173/157 (110%)	171 (99%)	2 (1%)	71	53
1	B	171/157 (109%)	170 (99%)	1 (1%)	86	76
1	C	170/157 (108%)	165 (97%)	5 (3%)	42	16
1	D	171/157 (109%)	170 (99%)	1 (1%)	86	76
1	E	170/157 (108%)	169 (99%)	1 (1%)	86	76
1	F	167/157 (106%)	166 (99%)	1 (1%)	86	76
All	All	1022/942 (108%)	1011 (99%)	11 (1%)	73	57

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	133	GLU
1	D	7	ARG
1	F	126	GLN
1	E	30	TYR
1	C	20	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	149	HIS
1	F	126	GLN
1	D	149	HIS
1	F	149	HIS
1	F	23	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 14 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	170/174 (97%)	-0.32	1 (0%) 89 90	9, 14, 26, 70	0
1	B	170/174 (97%)	-0.31	2 (1%) 79 81	10, 14, 26, 64	0
1	C	170/174 (97%)	-0.29	1 (0%) 89 90	11, 14, 26, 65	0
1	D	170/174 (97%)	-0.37	0 100 100	11, 14, 26, 64	0
1	E	170/174 (97%)	-0.32	1 (0%) 89 90	11, 14, 25, 78	0
1	F	170/174 (97%)	-0.30	2 (1%) 79 81	11, 14, 24, 48	0
All	All	1020/1044 (97%)	-0.32	7 (0%) 87 89	9, 14, 26, 78	0

The worst 5 of 7 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	2	SER	3.7
1	F	2	SER	3.7
1	F	90[A]	GLU	3.3
1	B	82	ASP	2.7
1	C	90[A]	GLU	2.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	FE2	B	303	1/1	0.72	0.12	59,59,59,59	1
2	FE2	F	203	1/1	0.84	0.10	67,67,67,67	1
2	FE2	A	202	1/1	0.90	0.12	60,60,60,60	1
2	FE2	D	202	1/1	0.93	0.20	67,67,67,67	1
2	FE2	C	202	1/1	0.95	0.15	58,58,58,58	1
2	FE2	E	202	1/1	0.96	0.14	66,66,66,66	1
2	FE2	B	301	1/1	0.97	0.06	29,29,29,29	1
3	FE	F	202	1/1	0.97	0.05	31,31,31,31	1
2	FE2	F	201	1/1	0.98	0.08	29,29,29,29	1
2	FE2	A	201	1/1	0.98	0.06	27,27,27,27	1
2	FE2	B	302	1/1	0.98	0.04	28,28,28,28	1
2	FE2	C	201	1/1	0.99	0.04	28,28,28,28	1
2	FE2	E	201	1/1	0.99	0.04	30,30,30,30	1
2	FE2	D	201	1/1	0.99	0.05	28,28,28,28	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around FE2 B 303:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around FE2 F 203:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around FE2 A 202:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around FE2 D 202:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around FE2 C 202:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around FE2 E 202:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around FE2 B 301:**



$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around FE2 F 201:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around FE2 A 201:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







**Electron density around FE2 B 302:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around FE2 C 201:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around FE2 E 201:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers ⓘ

There are no such residues in this entry.